

DOCUMENT RESUME

ED 118 558

SP 009 847

AUTHOR
TITLE

Pyfer, Howard R.; And Others
Cardiac Arrest During Medically-Supervised Exercise
Training: A Report of Fifteen Successful
Defibrillations.

PUB DATE
NOTE

May 75
3p.; Paper presented at the American College of
Sports Medicine Meeting (New Orleans, Louisiana, May
1975)

EDRS PRICE
DESCRIPTORS

MF-\$0.83 HC-\$1.67 Plus Postage
*Cardiovascular System; *Exercise (Physiology); Heart
Rate; Intervention; *Males; Medical Assistants;
Medical Treatment; *Older Adults; Physical Fitness;
*Risk

ABSTRACT

The Cardio-Pulmonary Research Institute conducted an exercise program for men with a history of coronary heart disease. Over 7 years, there were 15 cases of cardiac arrest during exercise (one for every 6,000 man-hours of exercise). Trained medical personnel were present in all cases, and all were resuscitated by electrical defibrillation with no permanent damage. Coronary heart disease patients are at risk of cardiac arrest during exercise, and there seems to be no way of predicting by age, type of exercise, or seeming tolerance to exercise whether or not cardiac arrest will occur. Nor are months of regular exercise a safeguard against an attack. However, exercise, even vigorous exercise, is safe if there are medical personnel present, and the physiological and psychological benefits of exercise in the rehabilitation process for coronary heart disease patients are important. (CD)

* Documents acquired by ERIC include many informal unpublished *
* materials not available from other sources. ERIC makes every effort *
* to obtain the best copy available. Nevertheless, items of marginal *
* reproducibility are often encountered and this affects the quality *
* of the microfiche and hardcopy reproductions ERIC makes available *
* via the ERIC Document Reproduction Service (EDRS). EDRS is not *
* responsible for the quality of the original document. Reproductions *
* supplied by EDRS are the best that can be made from the original. *

CAPRI

CARDIO  PULMONARY
RESEARCH INSTITUTE

U.S. DEPARTMENT OF HEALTH,
EDUCATION & WELFARE
NATIONAL INSTITUTE OF
EDUCATION

"THIS DOCUMENT" HAS BEEN REPRO-
DUCED EXACTLY AS RECEIVED FROM
THE PERSON OR ORGANIZATION ORIGI-
NATING IT. POINTS OF VIEW OR OPINIONS
STATED DO NOT NECESSARILY REPRESENT
OFFICIAL NATIONAL INSTITUTE OF
EDUCATION POSITION OR POLICY.

HONORARY TRUSTEES

ROBERT A. BRUCE, M.D.

BELVIN L. DOANE
Co-founder

HENRY FONDA

SAMUEL M. FOX III, M.D.

ALLEN A. FUNT

ROBERT KING, M.D.

BOARD OF TRUSTEES

RICHARD E. BANGERT
Treasurer

ALFRED J. BARRAN

GEORGE H. BIDDLE

WALLACE R. BUNN

C. C. (JOE) COLE

CHARLES M. DANIEL

JACK M. GOODFELLOW

FREDERICK P. HOLBROOK

WILLIAM T. JACOBSON

HAROLD C. KEAN

RONALD C. KISSUNG

WILLIAM F. MEAD, M.D.

J. THOMAS PAYNE, M.D.

FRANK A. PRITCHARD

HOWARD R. PYFER, M.D.
Co-founder

BROOKS G. RAGEN

W. HUNTER SIMPSON
President

HUGH A. SMITH
Vice President

SAMUEL N. STROUM

MILTON TRAFTON
Vice President

TED ROWE WATKINS
Secretary

T. EVANS WYCKOFF

HOWARD R. PYFER, M.D.
Director of
Medical Services

RICHARD FREDERICK
Managing Director

CARDIAC ARREST DURING MEDICALLY-SUPERVISED EXERCISE TRAINING

A REPORT OF FIFTEEN SUCCESSFUL DEFIBRILLATIONS

Howard R. Pyfer, M.D., William F. Mead, M.D. and Richard C. Frederick, M.S.
Cardio-Pulmonary Research Institute (CAPRI), Seattle, Washington

During the seven years between May, 1968 and May, 1975, cardiac arrest occurred 15 times in 3 cities where medically-supervised CAPRI cardiac exercise training programs were being conducted. Cardiac arrest is defined, here, as the cessation of adequate circulation of blood to sustain life.

CAPRI exercise training sessions meet three times weekly for 45 minutes of walking, jogging and calisthenics. Applicants with coronary heart disease or obstructive lung disease are accepted into the program only by physician referral. Trained medical personnel and emergency equipment are present at all exercise and testing sessions.

Arrests occurred approximately every 6,000 man-exercise hours of medically-supervised exercise training. Two happened within 60 seconds of each other. Fourteen occurred within the first 20 minutes of class; 2 while jogging, 5 while walking after jogging, 4 while standing after jogging, 1 while walking prior to jogging and 2 during calisthenics. One occurred 37 minutes after the beginning of class. All 15 were successfully resuscitated by electrical defibrillation without residual cardiac or brain damage and all were hospitalized for observation. Twelve (80%) returned to exercise training after release from the hospital. Eleven were between 50 and 60 years of age, 1 under 40 and 3 over 60 years. Fourteen had been enrolled in the program more than 6 months and 1 more than 48 months. Twelve (80%) had a history of heart attack, and 3 (20%) had angina pectoris as a primary reason for entering the program. Subsequent coronary angiography was performed on 11 (73%) and coronary artery by-pass surgery on 8 (53%). The exercise tolerance test prior to the cardiac arrest showed that 11 (73%) had electrocardiograph evidence of ischemic heart disease, 10 (67%) had arrhythmia, and 5 (33%) experienced angina pectoris. Two of the 15 died more than 6 months after the initial episode from another cardiac arrest, 1 at home and the other during coronary angiography. The remaining 13 are living and active.

COMMENTS: Coronary heart disease patients are at risk of cardiac arrest during exercise. The number of months that regular exercise training has been performed does not guarantee protection from cardiac arrest.

2



United Way Member Agency

Portland Program

651 S.W. CAPITOL HIGHWAY

PORTLAND, OR 97219

(503) 244-0114

Home Office

914 EAST JEFFERSON - SEATTLE, WASHINGTON 98122 (206) 323-7550

Non-Profit Organization

Yakima Program

5 NORTH NACHES

YAKIMA, WA 98101

(509) 248-1202

Age, exercise tolerance test findings, type or duration of exercise training do not offer any apparent single common denominator for predicting the likelihood of cardiac arrest.

Exercise training therapy for coronary heart disease patients provides positive physiological and psychological changes that contribute to the over-all rehabilitation process. Vigorous exercise, such as jogging and calisthenics, has been shown to be relatively safe if done in a medically-supervised environment.