

DOCUMENT RESUME

ED 244 184

CG 017 456

AUTHOR Mahurin, Kathleen A.; Gatz, Margaret
 TITLE Depression, Health, and Somatic Complaints in Older Adults.
 PUB DATE Aug 83
 NOTE 13p.; Paper presented at the Annual Convention of the American Psychological Association (91st, Anaheim, CA, August 26-30, 1983).
 PUB TYPE Reports - Research/Technical (143) -- Speeches/Conference Papers (150)

EDRS PRICE MF01/PC01 Plus Postage.
 DESCRIPTORS *Depression (Psychology); Factor Analysis; Factor Structure; *Middle Aged Adults; *Older Adults; *Physical Health; Self Evaluation (Individuals)
 IDENTIFIERS *Brief Symptom Inventory

ABSTRACT

Although depression is considered to be common in the elderly, reliable rates of prevalence are lacking. Studies have shown that age differences on measures of depressive symptomatology can be attributed to higher levels of somatic complaints. In order to examine whether the association between somatic and depressive symptoms varies as a function of health status in an older population, a sample of adults aged 50 to 87 (N=111) were surveyed using a self-report health rating, and the Brief Symptom Inventory (BSI) to assess psychiatric symptomatology. Factor analysis of the BSI yielded high loadings for each of the somatic and depressive symptoms, indicating the independence of the two constructs and providing evidence for the validity of the BSI. Data analysis showed that respondents endorsed more somatic complaints than depressive complaints. Of the eight symptoms endorsed most often, six were somatic items. These findings indicate that rates of prevalence for depression in the elderly may be overestimated when measures of depressive symptomatology that include somatic complaints are used. Descriptive statistics for the BSI comparing an elderly sample with normal young, normal adolescent, and psychiatric outpatient samples, along with data analyses, are appended. (LLL)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *

ED244184

CG017456

Depression, Health, and Somatic Complaints in Older Adults

Kathleen A. Mahurin and Margaret Gatz

University of Southern California

U.S. DEPARTMENT OF EDUCATION
NATIONAL INSTITUTE OF EDUCATION
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)
* This document has been reproduced as
received from the person or organization
originating it.
Minor changes have been made to improve
reproduction quality.
• Points of view or opinions stated in this docu-
ment do not necessarily represent official NIE
position or policy.

PERMISSION TO REPRODUCE THIS
MATERIAL HAS BEEN GRANTED BY

Kathleen A. Mahurin

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC)."

Paper presented at American Psychological Association, Anaheim,
CA, August, 1983.

The data presented in this paper were collected as part of a
geriatric health needs assessment survey. The authors would like
to thank John Meyer of St. Vincent Medical Center, Connie
Dessonville, Curt Hileman, Julie Patterson, Judy Zarit, Cynthia
Pearson, Kathy Marshall, Penny Hand, Monica Basco, and the many
others who assisted in data collection and analysis. Finally, we
especially thank Norman Cliff for his consultation concerning
methods of analysis.

INTRODUCTION

Depression is considered to be common in the elderly, however, reliable rates of prevalence are lacking. Based on findings from 3 sources of data: psychiatric case registers, surveys of institutional and clinical populations and general population surveys, the prevalence rates for depression range from 2% to 65% (Gurland, 1976). Investigations of these discrepancies revealed that the number of elderly suffering from depressive symptomatology was much greater than those with an actual "case" of depressive illness (Gurland, 1976; Raskin, 1979; Blazer, 1983). In community studies alone, rates of depressive disorders based on psychiatric interviews were consistently lower than rates that used symptom checklists to identify depressed persons.

Studies have shown that age differences on measures of depressive symptomatology can be attributed to higher levels of somatic complaints (Zung, 1967; Blumenthal, 1975; Zemore & Eames, 1979; Berry & Storandt, 1983). Kay and Bergmann (1980) point out that when scales include multidimensional symptomatology, it becomes difficult to separate psychological symptoms from those with a physical basis. Siegler and Costa (in press) indicate that this difficulty increases with age. Therefore, several researchers suggest that the higher levels of somatic complaints reflect the increased incidence of physical illness in the elderly rather than depression (Blazer, 1983).

The purpose of this study was to test this hypothesis by examining whether the association between somatic and depressive symptoms varied as a function of health status in an older population.

METHODS

Subjects. The sample consisted of 111 community dwelling elderly persons. They were selected by using a multistage probability-proportionate-to-size cluster sampling frame to identify a representative sample of adults aged 50 and older to interview as part of a geriatric health needs assessment survey conducted in a central city area of Los Angeles. Persons ranged in age from 50 to 87 with a mean age of 65.1 years. Ethnic balance was as follows: 52% white, 32% Hispanics, 8% black and 8% Asian, with 34.9% reporting that their primary language was not English. The average educational level was 10 years. Economic status was predominantly at the poverty level, over 68% of the respondents reporting incomes of less than \$7000 per year. Women and men were comparable on age, educational level and income but women slightly outnumbered men by 1.4:1.

Procedures. Health was assessed with a self-report rating of "excellent", "good", "fair", or "poor" and a medical illness checklist. Although these measures cannot be considered substitutes for clinical examinations, they represented economical global assessments which other investigators have found to compare favorably with physician's ratings (LaRue, Bank, Jarvik, & Hetland, 1979) and with other objective measures for assessing the health of community elderly (Fillenbaum, 1979). The Brief Symptom Inventory (BSI) was used to assess psychiatric symptomatology (Derogatis & Spencer, 1982). The BSI provides separate subscales for somatic and depressive complaints and is the most recent version of the Hopkins Symptom Checklist which has been recommended by Waskow and Parloff (1975) for measuring individual distress in adults. However, little is known about the use of this scale with an older population (see Pearson & Gatz, 1982, for scale means and reliabilities for this sample).

RESULTS

Descriptive statistics for the BSI were consistent with previous studies with other scales. Respondents endorsed more somatic complaints (68% endorsed at least one symptom) than depressive complaints (55% endorsed at least one symptom). Furthermore, of the 8 symptoms endorsed most often, 6 were somatic items (see Table 1). The construct validity of the BSI was tested by performing a factor analysis on the somatic symptoms and the symptoms of depression. In addition to being a preliminary first step to testing our hypothesis, the result of this analysis was interesting in and of itself for the following reason. If the increase of somatic symptoms in the elderly represents a different type of depression, factor analyzing the symptoms from both dimensions should reveal considerable interdependence and a failure to validate the construct validity of the two scales. If, however, the symptoms represent two separate constructs (as has been shown in non-aged samples), the results would indicate that somatic symptoms are representing something other than depression.

Results of the factor analysis yielded high factor loadings for the cluster of items defining each construct. Figure 1 illustrates a factor plot of the symptoms. Somatic symptoms showed high loadings on the somatization factor and depressive symptoms showed high loadings on the depression factor. These results confirmed the independence of somatic and depressive symptoms for older adults and provided evidence for the validity of these BSI subscales.

The distribution of scores for the somatic and depression dimensions were highly skewed, with 29% failing to endorse any items. We reconciled the problem by constructing 2 x 2 contingency tables reflecting the frequency of persons endorsing or not endorsing somatic or depressive

complaints. By performing Chi-square analyses on this data, we were able to test the relationship between the endorsement of somatic complaints and the endorsement of depressive complaints.

In order to examine the influence of health, we divided our sample into two groups according to their self-rated health scores. The mean of the 4 point scale was 2.33 with a higher rating indicating poorer health. These results are consistent with other findings in the literature (Shanas & Maddox, 1968; Tissue, 1972). For the purposes of analysis, persons who rated their health as excellent or good were considered in good health and persons who rated their health as fair or poor were considered in poor health. Fifty-nine percent of the sample fell into the good health group and 41% into the poor health group.

Three separate chi-square analyses were performed: the total sample, the groups in good health, and the group in poor health. The results are illustrated in Figures 2 and 3. A significant relationship between the endorsement of somatic and depressive complaints was found for the total sample. However, looking at the total sample obscured an important difference. When the sample was divided into two groups according to self-rated health, a significant association between depression and somatization was found for the healthy group but not for the unhealthy group. For the group in poor health, somatic complaints appeared to reflect physical health problems, not depression.

DISCUSSION

The results of the factor analysis indicate that the BSI is a valid scale to use with older adults when assessing somatic and depressive symptomatology. By separating out the items that define each construct, a more precise evaluation of the contribution of each dimension can be made. This is especially important when conducting research with older adults since the distinction between symptoms may be less obvious with age (Siegler & Costa, in press).

The results of the chi-square analyses confirmed the influence of health status on the endorsement of somatic symptoms. A significant association between somatic and depressive complaints was found for persons with good health but not for persons with poor health. For the latter group, the lack of association seemed to be due to their greater endorsement of somatic complaints.

These findings support previous explanations for the increase in the reporting of somatic complaints with age, namely, the greater decline in health of older persons. They also indicate that rates of prevalence for depression in the elderly may be overestimated when measures of depressive symptomatology are used that include somatic complaints. Therefore, it is important when using measurements in assessing the elderly to consider the role of health and its influence on patterns of complaints.

REFERENCES


- Berry, J. M. & Storandt, M. (1983, August). Age and sex differences in somatic complaints associated with depression. Paper presented at the meeting of the American Psychological Association, Anaheim, CA.
- Blazer, D. (1983). The epidemiology of depression in late life. In L. D. Breslau & M. R. Haug (Eds.), Depression and aging: Causes, care, and consequences. New York: Springer Publishing Co.
- Blumenthal, M. D. (1975). Measuring depressive symptomatology in a general population. Archives of General Psychiatry, 32, 971-978.
- Derogatis, L. R. & Spencer, P. M. (1982). The Brief Symptom Inventory (BSI): Administration, scoring, and procedures manual - I. Baltimore: Clinical Psychometric Research.
- Fillenbaum, G. G. (1979). Social context and self-assessment of health among the elderly. Journal of Health and Social Behavior, 20, 45-51.
- Gurland, B. J. (1976). The comparative frequency of depression in various adult age groups. Journal of Gerontology, 31, 283-292.
- Kay, D. W. K. & Bergmann, K. (1980). Epidemiology of mental disorders among the aged in the community. In J. E. Birren & R. B. Sloane (Eds.), Handbook of mental health and aging. Englewood Cliffs, N.J.: Prentice-Hall.
- LaRue, A., Bank, L., Jarvik, L., & Hetland, M. (1979). Health in old age: How do physicians' ratings and self-ratings compare? Journal of Gerontology, 34, 687-691.
- Raskin, A. (1979). Signs and symptoms of psychopathology in the elderly. In A. Raskin & L. F. Jarvik (Eds.), Psychiatric symptoms and cognitive loss in the elderly. Washington: John Wiley & Sons.
- Shanas, E. & Maddox, G. L. (1976). Aging, health and the organization of health resources. In R. Binstock & E. Shanas (Eds.), Handbook of aging and the social sciences. New York: Van Nostrand Reinhold.
- Siegler, I. C. & Costa, P. T. (in press). In J. E. Birren & K. W. Schaie (Eds.), Handbook of the psychology of aging. New York: Van Nostrand Reinhold.
- Tissue, T. (1972). Another look at self-rated health among the elderly. Journal of Gerontology, 27, 91-94.
- Waskow, I. E. & Parloff, M. B. (Eds.). (1975). Psychotherapy change measures. Washington, DC: DHEW Publication No. (Adm) 74-120.
- Zemore, R. Eames, N. (1979). Psychic and somatic symptoms of depression among young adults, institutionalized aged and noninstitutionalized aged. Journal of Gerontology, 34, 716-722.
- Zung, W. W. K. (1967). Depression in the normal aged. Psychosomatics, 8, 287-291.

TABLE 1

Descriptive Statistics for Brief Symptom Inventory comparing an elderly sample with norms from Derogatis (1982) for Normal Young, Normal Adolescents, and Psychiatric Outpatient samples

	(Mahurin & Gatz) Older Adults (n=111)	(Derogatis, 1982) Normal Young (n=719)	(Derogatis, 1982) Normal Adolescents (n=2408)	(Derogatis, 1982) Psychiatric Outpatients (n=1002)
BSI SYMPTOMS RANKED BY PERCENTAGE SYMPTOMATIC				
#	%	%	%	%
16: Feeling lonely (d)	47	25	47	82
37: Feeling weak in parts of body (s)	43	22	42	54
17: Feeling blue (d)	39	33	46	88
29: Trouble getting your breath (s)	33	14	19	33
7: Pains in heart or chest (s)	31	16	43	38
23: Nausea or upset stomach (s)	31	23	39	51
33: Numbness or tingling in body (s)	30	24	31	38
2: Faintness or dizziness (s)	28	17	43	42
18: Feeling no interest in things (d)	22	19	38	69
30: Hot or cold spells (s)	22	18	23	40
35: Feeling hopeless about the future (d)	20	18	32	78
50: Feeling of worthlessness (d)	19	11	29	71
9: Thoughts of ending your life (d)	10	3	20	46
Somatization (s)	.48	.29	.63	.83
Depression (d)	.45	.28	.82	1.80
Global Severity Index (GSI)	.41	.30	.83	1.32
Number of Symptoms Endorsed (PST)	13.05	11.45	24.81	30.80

FIGURE 1
 PLOT OF REFERENCE STRUCTURE FOR FACTOR 1 AND FACTOR 2



SOMATIC ITEMS

BSI 23	BSI 2
BSI 33	BSI 7 & BSI 37
BSI 29	BSI 30

DEPRESSIVE ITEMS

BSI 17
BSI 35
BSI 18
BSI 50
BSI 16
BSI 9

FIGURE 2

Frequency data on persons reporting different complaint patterns.

ALL SUBJECTS

Somatic Complaints

	no	yes	
no	32	22	54
yes	9	48	57
	41	70	111=N

Chi-Square = 22.50 **

$p < .05$


* $p < .01$

note: no = endorsed no complaints

yes = endorsed one or more complaints

FIGURE 3

The influence of health on the reporting of different complaint patterns.



* $p < .05$

** $p < .01$