

DOCUMENT RESUME

ED 323 422

CG 022 744

AUTHOR Evans, Shirley King
TITLE Nutrition and the Elderly: January 1987-May 1990.
Quick Bibliography Series.
INSTITUTION National Agricultural Library, Beltsville, MD.
REPORT NO QB-90-78
PUB DATE Aug 90
NOTE 103p.
AVAILABLE FROM U.S. Department of Agriculture, National Agricultural
Library, Public Services Division, Room 111,
Beltsville, MD 20705.
PUB TYPE Reference Materials - Bibliographies (131)
EDRS PRICE MF01/PC05 Plus Postage.
DESCRIPTORS *Adult Day Care; Adult Foster Care; Eating Habits;
Food Standards; *Frail Elderly; Gerontology; Health;
*Nutrition; Nutrition Instruction; *Older Adults
IDENTIFIERS AGRICOLA; *National Agricultural Library DC

ABSTRACT

This bibliography consists of 307 recent citations (January 1987 through May 1990) from AGRICOLA, the National Agricultural Library (NAL) computerized database. The bibliography addresses issues concerning nutrition, adult day care, and the elderly. Each citation includes the NAL call number, the title, the author(s), the city of publication, the name and date of the journal in which it is published (if applicable), page numbers, language, descriptors, and abstract. The citations are listed in alphabetical order. Author and subject indices are included. (TE)

* Reproductions supplied by EDRS are the best that can be made *
* from the original document. *


Nutrition and the Elderly

January 1987 - May 1990

Quick Bibliography Series: QB 90-78

307 Citations from AGRICOLA

**Shirley King Evans
Food and Nutrition Information Center**



**Food &
Nutrition**

National Agricultural Library Cataloging Record:

Evans, Shirley King.

Nutrition and the elderly.

(Quick bibliography series ; 90-78)

1. Aged -- Nutrition -- Bibliography.

aZ5071.N3 no.90-78



AVAILABILITY OF CITED DOCUMENTS

Expanded Service Patrons

The National Agricultural Library, Lending Branch provides direct lending services to persons working for the U.S. Congress, libraries and information centers. Through an interagency agreement with the Food and Nutrition Service (FNS) of USDA, NAL is able to expand the categories of users eligible for direct loan of books and audiovisuals, free photocopies of journal articles, and comprehensive reference/research services. This expanded group includes persons working for federal government agencies and cooperators with FNS programs:

- Elementary and secondary schools and school districts (including teachers, nurses, and food service and media center staff)
- Nutrition Education and Training Program
- Supplemental Food Program for Women, Infants and Children (WIC), and Commodity Supplemental Food Program (CSFP)
- Child Care Food Program (CCFP)
- Food Distribution Program on Indian Reservations (FDPIR)
- State Agencies (State employees only)

Submit lending requests on Individual Request Forms (IRF), one request per form; provide complete address, telephone number, job title and original signature of the requester to:

Interlibrary Loan
Lending Branch, 8th Floor
USDA National Agricultural Library
Beltsville, MD 20705

General Service Patrons

For those who do not fall into the above categories, (including some groups which previously had expanded service), food and nutrition materials must be obtained through interlibrary loan. The librarian in your public, state, university or corporate library can assist you as all requests must comply with the National or International Interlibrary Loan Code. Current charges for photocopies are \$5.00 for the first 10 pages; \$3.00 for each additional 10 pages; \$5.00 for the first fiche and \$.50 for each additional fiche; \$10.00 for duplicate reel of microfilm.

USDA Patrons

The materials listed in this bibliography may be obtained by submitting one Form AD 245 for each item requested to your local Agency or Regional Document Delivery System Library or directly to the National Agricultural Library, Lending Branch.

For general information, please call (301) 344-3755.

For reference service, subject searching and identification of newest editions or titles, please call (301) 344-3719.

For document delivery and booking of audiovisuals, please call (301) 344-2994.

AGRICOLA

**Citations in this bibliography were
entered in the AGRICOLA
database between January 1979
and the present.**

SEARCH STRATEGY

S1	114127	NUTRI?
S2	132522	FOOD?
S3	3731	EATING?
S4	49136	DIET?
S5	237730	(NUTRI? OR FOOD? OR EATING? OR DIET?)
S6	2538	ELDERLY
S7	1274	SENIOR
S8	753	CITIZEN?
S9	133	SENIOR (W) CITIZEN?
S10	7015	ADULT
S11	10065	DAY?
S12	14	ADULT (W) DAY
S13	2205	AGING
S14	2162	S5 AND (ELDERLY OR SENIOR ()CITIZEN? OR ADULT ()DAY/ OR AGING)
S15	36969	SF=FNC (FOOD AND NUTRITION)
S16	1411	S14 AND SF=FNC
S17	203956	PY=1987:PY9999
S18	307	S16 AND PY=1987:9999

Nutrition and the Elderly

1 NAL Call No: TX361.A3J6
The ABCs of nutritional assessment in older adults.

Fanelli, M.T.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (3): p. 33-40. charts; 1987. Includes 11 references.

Language: English

Descriptors: Nutritional assessment; Elderly; Nutritional state; Anthropometric dimensions; Clinical investigations; Nutrient intake; Blood composition; Standards; Guidelines

Abstract: A brief report provides an overview of nutritional assessment techniques and standards appropriate for elderly subjects over the continuum of care, and discusses criteria for detecting malnutrition risk in such subjects. Specific attention is given to separate discussions of anthropometric, biochemical, clinical, and dietary techniques and standards. It is concluded that suitable assessment techniques and standards are limited for accurate assessments of nutritional status of the elderly. A list of nutritional risk factors for elderly subjects is appended.(wz)

2 NAL Call No: TX355.A2 1987
About good nutrition.

Channing L. Bete Co
South Deerfield, Mass. : Channing L. Bete Co., Inc; 1987.
15 p. : col. ill. ; 21 cm. Cover title. New] (First press run) sample copy. A scriptographic booklet.

Language: English

Descriptors: Nutrition--Study and teaching; Popular works; Diet; Exercise; Health

Abstract: This booklet describes how eating nutritiously promotes good health. It recommends eating a variety of foods from the basic food groups; exercising regularly to maintain a desirable weight; and moderating sugar, salt, fat, and alcohol intake. It also discusses the special nutrition needs of children, teenagers, women, and elderly individuals. Illustrations are included.

3 NAL Call No: TX361.A3J63
Acetylsalicylic acid dose fails to affect energy intake of osteoarthritic elderly.

Taylor, J.; Krondl, M.; Rao, V.
Binghamton, N.Y. : Haworth Press; 1989.
Journal of nutrition for the elderly v. 8 (3,4): p. 79-96. charts; 1989. Includes 29 references.

Language: English

Descriptors: Caloric intake; Acetylsalicylic acid; Nutrient-drug interactions; Osteoarthritis; Elderly; Men

Abstract: The objective of the study was to determine the effect on energy intake and appetite of acetylsalicylic acid (ASA), commonly used by the elderly to treat arthritis. In a double blind crossover study, 23 free-living osteoarthritic patients 60 years of age or older were treated for two-week intervals with a mean daily ASA intake of 2.44 and 1.29 grams, respectively. Twenty healthy persons similar in age, taking no medication, and matched in sociocultural characteristics were included as a control group. Appetite was measured directly, using a visual analogue scale, and indirectly by calculating energy intake from three-day food records. Varying the dose of ASA was without effect on appetite and food energy intake; however, as appetite was rated lower by the medicated osteoarthritic than by the healthy group, although the energy intakes were not significantly different, the former should be considered as potentially at nutritional risk.

4 NAL Call No: TX361.A3J63
Actual and perceived nutritional knowledge of elderly individuals.

Mann, N.L.; Hildreth, G.J.; Draughn, P.; Hegsted, M.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 8 (4): p. 25-39. charts; 1988. Includes 10 references.

Language: English

Descriptors: Nutrition knowledge; Surveys; Demography; Socioeconomic status; Ethnicity; Elderly; Men; Women

Abstract: The perceived and actual nutritional knowledge of 150 noninstitutionalized elderly individuals was measured using mean scores on a 20 item test. Over 90% of those surveyed knew about the daily dietary need for vitamin C; about the life-long need for calcium; and about sources of fiber coming from whole grain breads and cereals. The elderly who thought they had fair knowledge of nutrition, in fact, did have higher mean scores on the nutritional knowledge test. Data analysis indicated that mean scores on actual nutritional knowledge were higher for those who were white and for those who were in a higher socioeconomic level. Mean scores of elderly men and women were found to be similar.

Quick Bibliography Series

5 NAL Call No: 389.8 AM34
ADA testimony on the 1987 reauthorization of the Older Americans Act of 1965.
Chicago, Ill. : The Association; 1987 Jul.
Journal of the American Dietetic Association v. 87 (7): p. 943-944; 1987 Jul. Includes 11 references.

Language: English

Descriptors: USA; Elderly nutrition; Federal aid; Federal programs; Congressional reports; Legislation

Abstract: The testimony given by the President of the American Dietetic Association (ADA) before the US Senate Subcommittee on Aging is reproduced, concerning ADA impressions and recommendations for the 1987 reauthorization of the 1965 Older Americans Act. Specific attention is given to: (1) Title III meals programs (home-delivered meals, congregate meals) and the need for improved training in food safety and sanitation; and (2) nutrition education needs. Additional programmatic considerations concerning reauthorization of the Older Americans Act also are described.(wz)

6 NAL Call No: 389.8 AM34
Age-related olfactory and taste changes and interrelationships between taste and nutrition.
Chauhan, J.; Hawrysh, Z.J.; Gee, M.; Donald, E.A.; Basu, T.K.
Chicago, Ill. : The Association; 1987 Nov.
Journal of the American Dietetic Association v. 87 (11): p. 1543-1550. charts; 1987 Nov. Literature review. Includes 91 references.

Language: English

Descriptors: Taste; Smell; Aging; Taste threshold; Age groups; Nutrient intake

Abstract: Extract: This article critically reviews the literature on age-related olfactory and taste changes and the interrelationships between nutrition and taste. Age-related taste alterations are controversial, but there is consensus on an olfactory decline with age. Much of the age-related chemoreceptive research has used inappropriate methodology, thus leaving many findings inconclusive. Changes in the taste system with age may be related to nutrient intake, which in turn can be influenced by taste. Literature on such interrelationships is limited. Vitamins A, thiamin, B-6, B-12, folacin, zinc, and copper are thought to influence taste function. Moreover, those nutrients are of concern in the aged. The use of appropriate methodology is critical not only for the study but

also for the nutritional aspects, particularly for qualitative and quantitative assessments of nutrient intakes. This review considers the experimental deficiencies (small sample size, inappropriate age range, inadequate psychophysical and stimuli presentation procedures, and poor dietary intake assessment) of studies reported in the literature and suggests major needs for research.(author)

7 NAL Call No: 321.8 C162
Aging in Canada: a nutritional perspective.
Kronld, M.; Coleman, P.
Ottawa : Canadian Home Economics Association; 1987.

Canadian home economics journal v. 37 (3): p. 119-123. ill; 1987.

Language: English

Descriptors: Canada--fre; Elderly; Aging; Nutrition physiology; Nutritional disorders

Abstract: Extract: With advancing age, physiological changes and socio-economic considerations become key issues influencing the vulnerability of elderly persons to nutrition-related disorders. Physical impairment may hinder food-related activities and medications may precipitate or aggravate nutritional deficiencies. Isolation can affect motivation to buy, prepare, and eat adequate meals. With persons over 75 years of age expected to be the fastest growing segment of the Canadian population in the near future, it is important that the role of nutrition "successful aging" is emphasized. Appropriate intervention programs using approaches which recognized the heterogeneity of elderly populations and build on reliable guidelines will serve to decrease nutritional risks. These services should offer to Canadian seniors a supportive network to assist them in delaying health deterioration and maintaining life satisfaction. (author)

8 NAL Call No: 389.8 AM34
Aging population presents marketing opportunities for dietitians.
Foltz, M.B.; Ryan, A.S.
Chicago, Ill. : The Association; 1987 May.
Journal of the American Dietetic Association v. 87 (5): p. 633-635. charts; 1987 May. Includes 16 references.

Language: English

Descriptors: USA; Elderly nutrition; Marketing; Dietitians; Socioeconomic status; Nutritional state; National surveys

Abstract: An evaluation of statistical data

Nutrition and the Elderly

developed during the 2nd National Health and Nutrition Examination Survey (NHANES II) was applied to develop marketing strategies for dietitians, focusing on the expanding elderly market. This evaluation showed that socioeconomic status influenced the level of fatness, with poorer men tending to be leaner and poorer women fatter than their richer counterparts. Other findings on factors influencing eating and nutrition (living alone; eating habits; medical history conditions) also are discussed. Anthropometric and nutrient intake data by age, per capita income, and living arrangement are summarized, covering both sexes between ages 55-74. The evaluation specifically identified several factors that could be integrated into nutrition programs for the older US population.(wz)

- 9 NAL Call No: 389.8 N953
The aging process: lessons from observations in man.
Leaf, A.
Washington, D.C. : Nutrition Foundation; 1988 Feb.
Nutrition reviews v. 46 (2): p. 40-44. charts; 1988 Feb. Includes 27 references.

Language: English

Descriptors: Elderly nutrition; Trends; Mortality; Demography
Abstract: A brief report summarizes evidence concerning changes in the major causes of mortality due to the increase in life span. Attention is given to: changes in the leading causes of mortality and in the per capita consumption of tobacco and various foods of potential health risk in the US during the 20th century; the dietary composition of elderly people; and the association between diet and longevity in different countries. The evidence suggests that diseases of affluence disproportionately afflict older people and that overnutrition or improper dietary balance directly affect the prevalence of atherosclerotic heart disease in our current societies.

- 19 NAL Call No: 448.8 B77
Alcohol consumption and dependence in elderly patients in an urban community.
Bridgewater, R.; Leigh, S.; James, O.F.W.; Potter, J.F.
London : British Medical Association; 1987 Oct10. British medical journal v. 295 (6603): p. 884-885. charts; 1987 Oct10. Includes 5 references.

Language: English

Descriptors: United Kingdom; Alcoholic beverages; Consumption patterns; Elderly; Urban areas; Sex differences

Abstract: A cross-sectional survey assessed the prevalence of alcohol consumption and dependence in elderly subjects (53 women, 48 men; mean age, 71) residing in an urban community. Mean alcohol consumption was 16.5 times/week, with men consuming more frequently than women (22.6 vs 10.9 times/week), and subjects 75 years old and younger consuming more frequently than subjects over age 75 (20.5 vs. 8.7 times/week). Sixteen of the subjects (31%; 4 men, 12 women) were abstainers, while 17 subjects (11 men, 6 women) were classified as being alcohol dependent. A survey employed a structured ("CAGE") questionnaire.(wz)

- 11 NAL Call No: 449.9 AM3J
Alcohol intake in a healthy elderly population.
Goodwin, J.S.; Sanchez, C.J., Thomas, P.; Hunt, C.; Garry, P.J.; Goodwin, J.M.
Washington, D.C. : American Public Health Association; 1987 Feb.
American journal of public health v. 77 (2): p. 173-177; 1987 Feb. Includes 29 references.

Language: English

Descriptors: Alcoholic beverages; Consumption patterns; Socioeconomic status; Psychological factors; Elderly; Sex differences; Cognitive development
Abstract: Extract: The relationship between amount of alcohol consumption and social, psychological, and cognitive status was examined in 270 healthy, independently living men and women over age 65. Forty-eight per cent of the sample recorded some alcohol intake during a three-day diet record, with 8 per cent drinking 30 or more grams of alcohol daily. Alcohol intake was positively associated with male gender, income, and amount of education and negatively associated with age. Alcohol intake was not associated with any changes in social or psychological status, but was positively associated with several measurements of cognitive status. These correlations were weak, however, and tended to disappear after controlling for income, education, gender, and age. Past alcohol intake was not associated with any indicators of present social, psychological, or cognitive functioning.(author)

- 12 NAL Call No: 389.8 Z22
Anemia and iron status in the free-living and institutionalized elderly in Kentucky.

Quick Bibliography Series

Chen, L.H.; Cook-Newell, M.E.
Bern : Hans Huber; 1989.
International journal for vitamin and nutrition research v. 59 (2): p. 207-213. charts; 1989. Includes 30 references.

Language: English

Descriptors: Kentucky; Nutritional state; Iron; Nutrient intake; Blood plasma; Iron deficiency anemia; Elderly

Abstract: The prevalence of iron deficiency in the elderly population of Central Kentucky was studied using hemoglobin and other biochemical indices of iron status. The association of iron intake with blood plasma levels of iron was examined for free-living and for institutionalized elderly.

13 NAL Call No: GN51.A58
Anthropometric standardization reference manual.

Lohman, Timothy G.,1940-; Roche, Alex F.,1921-; Martorell, Reynaldo,1947-
Champaign, IL : Human Kinetics Books; 1988.
vi, 177 p. : ill. ; 29 cm. Includes bibliographies and indexes.

Language: English

Descriptors: Anthropometry

Abstract: This reference manual is designed to serve as a comprehensive set of measurement procedures describing over 40 anthropometric dimensions. The purpose of the manual is to provide a standardized set of descriptions that can be used across disciplines (for example: epidemiology, exercise and sport science, human biology, human nutrition, medicine, physical anthropology, and physical education). The manual also includes a section on special issues such as right vs. left side, measurement error, equipment availability, and applications. The papers on applications show the many applications of anthropometry to research involving children, the elderly, handicapped or obese individuals, clinical nutrition, epidemiology, physical anthropology, sports medicine, and coronary heart disease.

14 NAL Call No: RM214.D47
Anthropometrics: an update on techniques.

Lieberman, L.S.; Kurtz, C.S.
Rockville, Md. : Aspen; 1987 Apr.
Directions in applied nutrition v. 1 (6): p. 1-2, 4-6. ill; 1987 Apr. Includes 22 references.

Language: English

Descriptors: Nutritional assessment; Methodology; Anthropometric dimensions; Infants; Children; Elderly; Body composition

Abstract: A report describes standard anthropometric techniques and instruments for assessment of nutritional state with respect to accuracy and precision and the potential limitations and misinterpretations of resultant data, including longitudinal measurements. Attention is focused on measurements of small-for-gestational-age infants, growth rates of boys and girls, nutritional assessment of the elderly, and innovative methods for measuring body composition.(wz)

15 NAL Call No: 389.8 J824
Ascorbate and dehydroascorbate: distribution in mononuclear cells of healthy elderly people.

VanderJagt, D.J.; Garry, P.J.; Bhagavan, H.N.
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Mar.
American journal of clinical nutrition v. 49 (3): p. 511-516; 1989 Mar. Includes 33 references.

Language: English

Descriptors: Ascorbic acid; Dehydroascorbic acid; Lymphocytes; Monocytes; Nutrient intake; Diet; Vitamin supplements; Elderly; Men; Women

Abstract: Concentrations of ascorbic acid and total ascorbate (ascorbic acid + dehydroascorbic acid) were determined in mononuclear cells from eight elderly men and nine elderly women (aged greater than 65 y) on controlled intakes of vitamin C. No increase in mononuclear cell ascorbic acid concentration was observed for intakes between 30 and 280 mg/d. A significant increase in total ascorbate concentrations was observed for the same range of intakes (p less than 0.01), primarily because of an increase in dehydroascorbic acid. Women were found to have significantly higher ascorbic acid concentrations in mononuclear cells than men (p less than 0.02) for similar intakes of vitamin C. No difference was observed for total ascorbate concentrations. The ratio of dehydroascorbate to ascorbate increased with increasing intake of vitamin C for both men and women.

16 NAL Call No: RC620.A1N8
Ascorbic acid and carnitine in man.

Davies, H.E.F.; Gruffudd, S.; Hughes, R.E.; Jones, E.
Stoneham, Mass. : Butterworth Publishers; 1987 Nov.
Nutrition reports international v. 36 (5): p. 941-948. ill., charts; 1987 Nov. Includes 17 references.

Nutrition and the Elderly

Language: English

Descriptors: Men; Women; Elderly; Ascorbic acid; Carnitine; Amino acid metabolism

Abstract: Extract: In apparently healthy elderly men supplementation of diet with 200 mg ascorbic acid daily increased urinary excretion of carnitine; ending of supplementation reduced urinary carnitine. In elderly female hospital patients urinary carnitine/creatinine ratio was positively correlated ($P=0.01$) with leucocyte ascorbic acid. Neither in young healthy women nor in elderly female hospital patients was a correlation found between leucocyte ascorbic acid and plasma carnitine concentration.(author)

17 NAL Call No: RC620.A1J6
Ascorbic acid, HDL cholesterol, and apolipoprotein A-I in an elderly Chinese population in Boston.

Dallal, G.E.; Choi, E.; Jacques, P.; Schaefer, E.J.; Jacob, R.A.
New York, N.Y. : John Wiley & Sons; 1989 Feb.
Journal of the American College of Nutrition v. 8 (1): p. 69-74. charts; 1989 Feb. Includes 24 references.

Language: English

Descriptors: Massachusetts; High density lipoprotein; Apolipoprotein; Ascorbic acid; Ethnicity; Nutritional surveillance; Elderly; Men; Women

Abstract: Over 300 elderly Chinese Americans were studied to determine the role that vitamin C may play in HDL cholesterol metabolism. Variables such as smoking, alcohol intake and health status were taken into consideration. Results were discussed and shown on tables.

18 NAL Call No: 389.8 J824
Ascorbic acid intake and plasma levels in healthy elderly people.

VanderJagt, D.J.; Garry, P.J.; Bhagavan, H.N.
Bethesda, Md. : American Society for Clinical Nutrition; 1987 Aug.
American journal of clinical nutrition v. 46 (2): p 290-294. ill., charts; 1987 Aug. Includes 17 references.

Language: English

Descriptors: Elderly; Nutrient intake; Blood composition; Sex differences; Nutritional state; Vitamin supplements; Ascorbic acid

Abstract: Extract: Plasma ascorbic acid concentra-

tions were determined in eight men and nine women over age 65 on controlled intakes of vitamin C. Plasma ascorbic acid levels were significantly lower in elderly men compared with women for vitamin C intakes ranging from 30 to 280 mg/d. A maximum steady-state level of 1.0 mg/dL (56.78 micromoles/L) ascorbic acid was observed in male subjects at an intake of approximately 150 mg/d; female subjects approached a plasma ascorbate level of 1.0 mg/dL (56.78 micromoles/L) with an intake of approximately 80 mg/d. At a total vitamin C intake of 60 mg/d, plasma ascorbate levels in men were estimated to be 0.4 mg/dL (22.71 micromoles/L) or less. The amount of dehydroascorbic acid in plasma did not vary with intake of vitamin C.(author)

19 NAL Call No: 389.8 AM34
Assessing productivity of foodservice systems in nutrition programs for the elderly.

Lieux, E.M.; Winkler, L.L.
Chicago, Ill. : The Association; 1989 Jun.
Journal of the American Dietetic Association v. 89 (6): p. 826-829. charts; 1989 Jun. Includes 15 references.

Language: English

Descriptors: Nutrition programs; Food service management; Program evaluation; Task analysis; Labor productivity; Nursing homes; Elderly

Abstract: A pilot study was conducted to develop a monitoring instrument for the measurement of productivity in nutrition programs for the elderly. Observations of foodservice activities were collected for 3 days in each of three senior centers utilizing cook/serve production models. Direct work activity categories included processing, service, transportation, clerical, cleaning, and receiving. Delay activities included personal delay and idle time. Information was gathered on the number of meals served. A method to recognize the equivalent work of senior citizen volunteers was developed. Productivity ratios were calculated as average labor minutes in each activity per meal served. Labor time was consistent within centers, and productivity varied with number of meals served. Average productivity ranged from 12.95 to 19.30 labor minutes per meal.

20 NAL Call No: 389.8 J824
Assessment of physical activity and energy needs
Shephard, R.J.
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.

Quick Bibliography Series

American journal of clinical nutrition v. 50 (5): p. 1195-1200; 1989 Nov. Includes 38 references.

Language: English

Descriptors: Energy requirements; Physical activity; Research methodology; Questionnaires; Elderly

Abstract: The assessment of physical activity and energy needs is discussed with particular reference to elderly subjects. The only practical method of determining physical activity in epidemiological research remains a retrospective questionnaire. Attitude measurements may nevertheless help in assessing those individuals prone to exaggerate participation, while personal activity monitors and measurement of individual fitness levels may contribute to the refinement of questionnaires. Particular problems in applying questionnaires to elderly subjects can include poor vision, hearing, and recent memory, coupled with slow understanding and garrulousness. The design of forms should be as simple and unambiguous as possible, and timely. Attempts to distinguish occupational from leisure or household chores and to explore seasonal variations merely seem likely to cause confusion.

21 NAL Call No: 389.8 AM34

Attitudes, knowledge, and problem-solving approach of Michigan dietitians about aging.

Noel, M.M.B.; Ames, B.D.

Chicago, Ill. : The Association; 1989 Dec.

Journal of the American Dietetic Association v. 89 (12): p. 1753-1757. charts; 1989 Dec. Includes 36 references.

Language: English

Descriptors: Michigan; Elderly nutrition; Nutrition knowledge; Dietitians; Surveys; Nutrition education; Problem solving; Professional competence

Abstract: Dietitians' attitudes and knowledge about aging are important because of the potential influence on quality and quantity of nutrition services provided to older adults. Four instruments were used to survey active members of The American Dietetic Association (ADA) in Michigan. The survey was mailed to 1,408 dietitians; 738 responded (52.4%). The responding dietitians were predominantly under 40, well educated, and female, which was consistent with the 1986 national census of ADA members. The results indicated that dietitians hold positive attitudes toward older adults in most areas. An exception was that a majority of the dietitians perceived older adults as resistant to treatment. Although dietitians' knowledge about aging generally was adequate, the ques-

tionnaire revealed that dietitians lacked knowledge in the areas of economic and health status; dietitians answered that older adults have a lower income and are sicker and more institutionalized than is in fact true. These are critical areas of deficiency in knowledge by dietitians, since food and nutrition issues often involve money and health. In the problem-solving section of the study, the dietitians usually focused on one specific environment or area of concern. Neither the full range of all possible environments for older adults nor the diversity of older adults' needs was usually considered. Continuing education as well as comprehensive undergraduate courses on aging would have a positive influence on Michigan dietitians' attitudes and knowledge about aging.

22 NAL Call No: TX361.A3J63

Attitudes of rural and urban elderly concerning supplement use.

Betts, N.M.; Rezek, J.

Binghamton, N.Y. : Haworth Press; 1989.

Journal of nutrition for the elderly v. 8 (3,4): p. 67-77. charts; 1989. Includes 10 references.

Language: English

Descriptors: Nebraska; Supplements; Nutrition information; Consumer attitudes; Calcium; Urban population; Rural population; Elderly; Women; Men

Abstract: To examine supplement purchase practices and attitudes toward supplements of rural and urban elderly, two questionnaires were designed. One questionnaire included demographic questions, an attitude scale, and questions concerning supplement purchases and sources of information about nutrition. The second questionnaire contained an advertisement about calcium supplements followed by questions regarding means to "improve calcium" and health concerns. The rural elderly had a significantly lower mean education level and a significantly higher number of misconceptions. The average expenditures on supplements were \$52.00 per month for the total group and sources of information were media, physicians, family and friends. In response to the advertisement, nearly one-third of the respondents reported that they would be unlikely to take calcium supplements to improve calcium even though 86.7% stated they would take calcium supplements to prevent health problems and 51.7% believed there could be dangers in taking calcium supplements. The most commonly reported means to improve calcium were to increase milk and to eat cheese

Nutrition and the Elderly

(86.7% each). Contradictions in the responses may indicate that these elderly are susceptible to advertising implying supplements are needed to prevent health problems.

23 NAL Call No: TX341.C6
B-12 recommendation lowered in 10th edition.
Washington, D.C. : Community Nutrition Institute; 1987 Jun18.
Nutrition week v. 17 (25): p. 6-7. charts; 1987 Jun18.

Language: English

Descriptors: USA; Vitamin B12; Nutrient requirements; Recommended dietary allowances; Recommended dietary intakes; Food and nutrition controversies; Vitamin supplements

Abstract: A brief technical report provides an excerpt of the recommended dietary intake (RDI) for vitamin B12 proposed for the 10th edition of the Recommended Dietary Allowances report rejected by the US National Academy of Sciences. Information is provided on vitamin B12 requirements for maintaining normal nutrient balance, and on vitamin B12 RDIs for sustaining normal infants, children, adults (including the elderly), and pregnant and lactating women. The coupled lack of apparent toxicity and health benefits from daily high intakes of vitamin B12 supplements also is noted.(wz)

24 NAL Call No: QP141.A1J68
Beta-carotene serum response in young and elderly females.
Maiani, G.; Mobarhan, S.; Ceccanti, M.; Ranaldi, L.; Gettner, S.; Bowen, P.; Friedman, H.; De Lorenzo, A.; Ferro-Luzzi, A.
Basingstoke : The Macmillan Press Ltd; 1989 Nov.
European journal of clinical nutrition v. 43 (11): p. 749-761. charts; 1989 Nov. Includes 41 references.

Language: English

Descriptors: Beta-carotene; Blood serum; Food intake; Aging; Experimental diets; Intestine transit time; Intestinal absorption; Blood lipids; Triglycerides; Young adults; Elderly; Women

Abstract: This study investigates the effects of aging and food intake on the serum beta-carotene response curve in humans. Serum carotenoids, retinoids, triglycerides, and cholesterol were measured after the oral administration of a fixed dose of beta-carotene given with a test meal to elderly and young adult women. Small intestine transit

time was determined for the two groups by breath hydrogen test. The data indicate that beta-carotene absorption may be enhanced in old age.

25 NAL Call No: 389.8 J824
Between- and within-subject variation in nutrient intake from infancy to old age: estimating the number of days required to rank dietary intakes with desired precision.

Nelson, M.; Black, A.E.; Morris, J.A.; Cole, T.J.
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Jul.

American journal of clinical nutrition v. 50 (1): p. 155-167. charts; 1989 Jul. Includes 41 references.

Language: English

Descriptors: Food consumption; Dietary history; Nutrition surveys; Nutrient intake; Data analysis; Research methodology

Abstract: In prospective studies of diet it is often necessary to know for how many days subjects should record food consumption in order to be able to rank subjects correctly according to their nutrient intakes. Data from six studies--of toddlers, families, schoolchildren, dietitians, pregnant women, and elderly subjects--were analyzed to estimate the number of recording days necessary for energy, 28 nutrients, and the ratio of polyunsaturated to saturated fatty acids (P:S). The most striking finding is that 7 d of diet record do not rank subjects with the degree of accuracy commonly assumed. For some nutrients, such as iron, zinc, nicotinic acid, and pyridoxine, it may be desirable to record diet over a number of short, separate periods to achieve the number of days required. For others, such as copper, retinol, carotene, vitamin B-12, polyunsaturated fatty acids, P:S, and alcohol, alternative methods of assessment based on dietary histories or questionnaires may be more appropriate.

26 NAL Call No: RC620.A1N8
Biochemical assessment of vitamin status in "healthy" elderly Japanese: relationships between intakes and biochemical measures of thiamine, riboflavin and ascorbic acid.

Itoh, R.; Yamada, K.; Oka, J.; Echizen, H.; Murakami, K.

Stoneham, Mass. : Butterworth; 1989 Mar.
Nutrition reports international v. 39 (3): p. 509-519. charts; 1989 Mar. Includes 20 references.

Language: English

Descriptors: Japan; Thiamine; Riboflavin; Ascorbic

Quick Bibliography Series

acid; Nutrient intake; Nutritional state; Erythrocytes; Enzyme activity; Glutathione reductase; Transketolase; Elderly

Abstract: Intakes and nutritional status of thiamine, riboflavin and ascorbic acid were assessed from 3-day food records and blood samples collected from 183 free-living healthy, middle-income volunteers over 60 years of age from Nagano City area in Japan. About 50% of the subjects consumed thiamine-fortified rice, which was the only vitamin supplement used in this population. Biochemical measures used for the assessment of the status of these three vitamins were erythrocyte transketolase activity coefficient (ETK-AC), erythrocyte glutathione reductase activity coefficient (EGR-AC) and serum ascorbic acid concentration. Significant correlation coefficients between these biochemical measures and estimated intakes for each of the vitamins examined were obtained. Although only 5.5% of the subjects had estimated thiamine intakes less than the 1984 RDA for Japanese, 37.6% of the subjects not taking the supplement and 13.3% of the subjects taking the supplement had ETK-ACs above 1.15 and assessed to be in poor status of this vitamin. With regard to riboflavin, 9.6% of the subjects had its intakes less than the RDA, and 3.7% had EGR-ACs above 1.20. Only 1.1% of the subjects had ascorbic acid intakes less than the RDA, and 4.9% had its serum concentrations less than 0.4 mg/100 ml. Thiamine nutriture seems to be a serious problem in this population.

27 NAL Call No: 389.8 Z33
Biochemical indicators related to antioxidant status and bone metabolic activity in Finnish elderly men.

Maenpaa, P.H.; Pirhonen, A.; Pirskanen, A.; Pekkanen, J.; Alftan, G.; Kivela, S.L.; Nissinen, A. Bern : Hans Huber; 1989.

International journal for vitamin and nutrition research v. 59 (1): p. 14-19. charts; 1989. Includes 29 references.

Language: English

Descriptors: Finland; Nutritional state; Vitamin A; Vitamin D; Vitamin E; Selenium; Calcium; Blood serum; Blood lipids; Tobacco smoking; Alcoholic beverages; National surveys; Elderly; Men

Abstract: A 25-year follow-up survey of the Finnish men originally examined in the Seven Countries Study was conducted. The follow-up survey examined biochemical indicators related to increased cancer risk and bone fragility. Serum concentrations of vitamin A, vitamin D, vitamin E, osteocal-

cin, calcium, selenium, cholesterol and HDL-cholesterol were measured in 270 elderly Finnish men.

28 NAL Call No: RC620.A1N8
Blood pressure and body mass index in elderly long-term vegetarians and nonvegetarians.

Melby, C.L.; Lyle, R.M.; Poehlman, E.T. Stoneham, Mass. : Butterworth Publishers; 1988 Jan.

Nutrition reports international v. 37 (1): p. 47-55. charts; 1988 Jan. Includes 16 references.

Language: English

Descriptors: USA; Elderly; Diet; Vegetarians; Blood pressure; Seventh day adventists; Weight; Height; Indexes

Abstract: Extract: Several cross-sectional studies have found vegetarians to have lower blood pressure (BP) than nonvegetarians. However, differences in non-dietary health practices exist between the two groups which could confound the apparent relationship between diet and BP. Also, few studies have examined the diet-BP relationship in the elderly. This study addressed these shortcomings by examining elderly (age greater than or equal to 60) vegetarian and nonvegetarian Seventh Day Adventists. Subjects voluntarily completed a questionnaire which included information on demographics, health history, dietary practices, and other health-related habits. Height and weight were measured and multiple blood pressure readings were taken for each individual. Subjects were divided into long term vegetarian (n=61, x age=69.8) and nonvegetarian (n=29, x age=70.0) categories based on self-reported dietary habits. Long term vegetarians were defined as adhering to a meatless dietary regime for at least 3 years prior to the study; non vegetarians were defined as current omnivores. Twenty-one percent of the vegetarians and 24% of the nonvegetarians were on current antihypertensive drug therapy, and were excluded from further analysis. Age and sex-adjusted systolic BP was significantly lower (p less than .01) in the vegetarians (n=48, x=122.2 mm Hg) compared to the nonvegetarians (n=22, x=136.1 mm Hg). Age and sex-adjusted diastolic BP (vegetarians, x=71.2 mm Hg vs. nonvegetarians, 73.7 mm Hg) was not significantly different between the two groups. Body mass index (weight/height²) was significantly greater (p less than .001) in the nonvegetarians (x=28.6) compared to the vegetarians (x=24.3). There were no differences between the two groups in reported

Nutrition and the Elderly

family history of hypertension, frequency of exercise, use of tobacco, and consumption of beverage ethanol. Further statistical analysis revealed that the lower systolic BP among the elderly vegetarians was best explained by their body mass in

29 NAL Call No: 1X361.A3J63
Blood pressure of elderly persons in relation to dietary practices, sodium preference in foods, and urinary sodium.
Stanek, K.L.; Fox, H.M.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 7 (3): p. 3-15. chart. 1988. Includes 20 references.

Language: English

Descriptors: Hypertension; Diet; Nutrient intake; Sodium; Food preferences; Body weight; Elderly
Abstract: Hypertensive and normotensive elderly patients in government-sponsored nutrition centers were compared with regard to dietary practices, urinary sodium excretions, and taste preference levels for sodium chloride in tomato juice and beef broth. Body mass index and body weight of 265 persons were positively and significantly related to blood pressure. Caloric intake of treated hypertensive but not of normotensive persons was related to blood pressure. Urinary sodium excretions did not differ between 42 treated hypertensive and 40 normotensive persons. Hypertensive persons tended to prefer a higher level of sodium in tomato juice than did normotensive persons.

30 NAL Call No: 389.8 N953
Body composition and aging.
Steen, B.
Washington, D.C. : Nutrition Foundation; 1988 Feb.
Nutrition reviews v. 46 (2): p. 45-51. ill; 1988 Feb. Includes 43 references.

Language: English

Descriptors: Body composition; Aging; Methodology
Abstract: Information and data developed from published studies are discussed concerning the assessment of body composition in elderly subjects. Specific attention is given to: the concepts of lean body mass and fat-free body mass; methods for gaining an understanding of the different body compartments; and assessments of body composition in higher age groups using a 4-compartment model technique considering total body potassium

and total body water. Data concerning body weight, body fat, body cell mass, and body water in the 7th and 8th decade of life are graphically presented and discussed.

31 NAL Call No: 389.8 J824
Body composition in elderly subjects: a critical appraisal of clinical methodology.
Heymsfield, S.B.; Wang, J.; Lichtman, S.; Kamen, Y.; Kehayias, J.; Pierson, R.N. Jr
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.
American journal of clinical nutrition v. 50 (5): p. 1167-1175. charts; 1989 Nov. Literature review. Includes 36 references.

Language: English

Descriptors: Body composition; Laboratory methods; Body density; Neutron activation analysis; Elderly

Abstract: Body composition measurements are integral to nutritional assessment of elderly subjects. The validity of methods available for partitioning body weight into its tissue or chemical subfractions in the geriatric population is of primary importance. Until recently verifying the accuracy of body composition techniques in vivo was limited by prevailing technology. New advances such as neutron activation analysis, neutron inelastic scattering, and dual photon absorptiometry now promise to overcome these earlier limitations. With these new tools investigators can now examine critical underlying assumptions related to body composition assessment in elderly subjects. This review highlights the need and opportunities for future research in this area.

32 NAL Call No: 389.8 J824
Bone mineral content in postmenopausal women: comparison of omnivores and vegetarians.
Hunt, I.F.; Murphy, N.J.; Henderson, C.; Clark, V.A.; Jacobs, R.M.; Johnston, P.K.; Coulson, A.H.
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Sep.
American journal of clinical nutrition v. 50 (3): p. 517-523. charts; 1989 Sep. Includes 31 references.

Language: English

Descriptors: Bone density; Mineral content; Nutrient intake; Calcium; Protein intake; Vegetarians; Elderly; Women

Abstract: This cross-sectional study of bone mineral content (BMC)/bone width (BW) (BMC/BW, cortical radius) was conducted in el-

Quick Bibliography Series

derly, free-living Methodist omnivores (n = 146) and Seventh-day Adventist vegetarians (n = 144) in southern California. The purpose was to investigate relationships between BMC/BW and dietary factors (primarily vegetarianism and intakes of protein and calcium) and use of prescribed drugs. BMC was measured by single-photon absorptiometry. Dietary intakes during current and early periods of life were assessed by 24-h dietary recall and frequency methods. Information about drug use was obtained by interview. BMC/BW was not different in omnivores vs vegetarians. There were no significant relationships or trends between current or early dietary intakes and BMC/BW in either group. In multiple-regression analysis, age, weight, and use of estrogen and thiazide diuretics were predictive of 31% of the variability in BMC/BW in the total group.

33 NAL Call No: 448.8 L22

The bran wagon.

Suri, Y.P.

Boston, Mass. : Little, Brown and Company; 1987 Jul04.

The Lancet v. 2 (8549): p. 42-43; 1987 Jul04. Includes 7 references.

Language: English

Descriptors: Diet; Bran; Fiber; Risks; Calcium; Zinc; Elderly nutrition; Patient care; Nutrient-nutrient interactions; Food and nutrition controversies

Abstract: A letter to the editor critiques a recent editorial on bran, arguing that some of the most serious problems resulting from a high-fiber diet were completely overlooked. These include the impaired absorption of certain minerals (especially Ca, which may accelerate osteoporosis in the elderly) and trace elements (e.g. Zn, needed for tissue repair and bone fracture healing), due to insoluble phytate formation. It is emphasized that these effects need to be considered by the nutritionist and dietitian for exercising moderation with fiber in high-risk groups.(wz)

34 NAL Call No: 389.2538 F82

Breaking down barriers.

Peters, J.E.

New York, N.Y. : Restaurant Business Magazine; 1987 Nov01.

Restaurant business v. 86 (16): p. 134-136, 138, 140. ill; 1987 Nov01.

Language: English

Descriptors: Restaurants; Elderly; Physically handicapped persons; Consumers; Eating out; Guidelines

Abstract: A report argues that restaurants should give increased attention to accommodating elderly and physically handicapped customers. Guidelines for accomplishing this are discussed and the benefits that restaurants can derive are emphasized. Specific examples are given throughout the report. Considerations for special facility and management accommodations and for proper training of the restaurant staff are included.

35 NAL Call No. 389.8 J824

Calcium and vitamin D supplements: effects on calcium metabolism in elderly people.

Chapuy, M.C.; Chapuy, P.; Meunier, P.J.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Aug.

American journal of clinical nutrition v. 46 (2): p. 324-328. charts; 1987 Aug. Includes 43 references.

Language: English

Descriptors: Calcium; Elderly nutrition; Vitamin D; Mineral supplements; Vitamin supplements; Parathyroid hormone

Abstract: Extract: Calcium and vitamin D status were studied in 193 healthy elderly French people. Calcium intake was less than 500 mg/d in 62% of the patients and the vitamin D intake was less than 5 micrograms/d in all patients. They also exhibited reduced levels of serum calcium (SCa) and 25-hydroxyvitamin D (25-OHD) and high levels of parathyroid hormone (PTH) and alkaline phosphatase (SAP). The response to calcium (1000 mg/d) and ergocalciferol (20 micrograms/d) supplementation given for 6 mo was evaluated in 65 patients with 69 subjects taken as controls. The treatment induced a significant increase in SCa and in 25-OHD levels, and a subsequent decrease in PTH levels without modification of the mean calcitriol levels. The biochemical changes were more marked in long-stay hospital patients than in outpatients. All these changes were significantly different from those observed in the control group. Increasing the calcium and vitamin D intake reduces the biochemical signs of secondary hyperparathyroidism in elderly people.(author)

36 NAL Call No: 389.8 J82

Calcium intake and skeletal integrity: is there a critical relationship?

Marcus, R.

Bethesda, Md. : American Institute of Nutrition;

Nutrition and the Elderly

1987 Apr.

The Journal of nutrition v. 117 (4): p. 631-635. ill; 1987 Apr. Includes 33 references.

Language: English

Descriptors: Calcium; Bone mineralization; Nutrient requirements; Skeleton development; Vitamin D; Estrogens; Menopause; Elderly; Reviews

Abstract: A technical review (33 references) summarizes and discusses current research evidence concerning the importance of calcium (Ca) intake on maintaining a peak bone mass and preventing bone loss. Attention is given to: the effect of Ca intake on peak bone mass and on bone loss in young adulthood; the relationship between Ca intake and bone loss in young adulthood; the relationship between Ca intake and bone loss at menopause; and the role of dietary Ca intakes in the maintenance of bone health in postmenopausal women and the elderly. It is argued that, while considerable research has been conducted, current evidence does not point to a clear relationship between high Ca intakes to prevent accelerated bone loss at menopause or their therapeutic value for the elderly. Relationships between dietary Ca intake, estrogen status in postmenopausal women, and vitamin D are discussed with respect to bone health status and bone loss prevention and treatment.(wz)

37

NAL Call No: TX341.H82

Can I afford the diet?

Hanes, F.A.; De Looy, A.E.

London, Eng. : J. Libbey; 1987 Feb.

Human nutrition : applied nutrition v. 41A (1): p. 1-12. charts; 1987 Feb. Includes references.

Language: English

Descriptors: United Kingdom; Food costs; Low income groups; Nutrient intake; Food policy; Adults; Children

Abstract: An authoritative technical review addresses the problem of nutritional inadequacy due to restricted income, and provides recommendations for aiding low-income people residing in the United Kingdom (UK) who are at greatest risk of illness from nutritional deficiencies. Specific attention is given to: the inadequacy of RDA's for ill people; average household expenditures for food by income group vs. the cost for maintaining a healthy diet; the influence of income on food consumption patterns; and, especially, special groups at malnutrition risk (covering: children; pregnant women; black and ethnic minorities; mental-

ly/physically handicapped; and the elderly). Attention also is given to the cost of special or therapeutic diets for adults and children having different requirements due to disorders or diseases (e.g., diabetes, renal failure). It is concluded that financial provision for food for many people on the UK Supplementary Benefit program is inadequate and that there should be no restriction in the existing provision of free school meals.

38

NAL Call No: TX1.I55

Caring for the elderly in America.

Stephenson, S.

Newton, Mass. : Cahners Publishing Company; 1989 Mar06.

Restaurants & institutions v. 99 (7): p. 105-106, 110. ill; 1989 Mar06.

Language: English

Descriptors: Nutrition programs; Adult day care; Community programs; Elderly

Abstract: Elderly day care centers are found in all areas of the country. The number of centers and their clientele are steadily increasing. Several centers and their programs are discussed.

39

NAL Call No: 389.8 J824

Changes in food and nutrient intake in middle-aged men from 1960 to 1985 (the Zutphen Study).

Kromhout, D.; Lezenne Coulander, C. de; Obermann-de Boer, G.L.; Kampen-Donker, M. van; Goddijn, E.; Bloemberg, B.P.M.

Baltimore, Md. : American Society for Clinical Nutrition; 1990 Jan.

American journal of clinical nutrition v. 51 (1). p. 123-129. charts; 1990 Jan. Includes 22 references

Language: English

Descriptors: Netherlands; Food consumption; Nutrient intake; Caloric intake; Dietary surveys; Longitudinal studies; Eating habits; Men

Abstract: Within the Zutphen Study dietary surveys using the cross-check dietary history method were carried out in 1960, 1965, 1970, and 1985. Of the 872 men aged 40-59 y examined in 1960, 315 participated in all four surveys. In 1985 a small random sample of 51 men aged 40-59 y was also investigated. Between 1960 and 1985 the consumption of bread, potatoes, and edible fats decreased and the consumption of fruits, pastries, nuts, and alcoholic beverages increased in both the aging cohort and in the two independent samples of middle-aged men examined 25 y apart. These changes were accompanied by a substantial

Quick Bibliography Series

decrease in the intake of monounsaturated fat, polysaccharides, and potassium and a substantial increase in animal protein and alcohol. Small changes were observed in the intake of saturated fatty acids, polyunsaturated fatty acids, dietary cholesterol, and dietary fiber. Since 1960 some changes in a nutritionally desirable direction were observed but the diet of the Zutphen men in 1985 departs substantially from the guidelines for a healthy diet formulated by The Netherlands Nutrition Council in 1986.

40 NAL Call No: TX361.A3J63
Changes in food preferences of the elderly over a ten year period.

Holt, V.; Nordstrom, J.; Kohrs, M.B.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 7 (4): p. 23-34; 1988. Includes 18 references.

Language: English

Descriptors: Food preferences; Food habits; Dietary surveys; Food groups; Longitudinal studies; Elderly

Abstract: The changes of food preferences of persons who were initially 60 years and older were determined by comparing responses to surveys taken in 1975 and 1985. In 1985 the foods chosen as liked were softer foods with general decreases in preferences for protein and fatty foods. There was an increase in the amount of carbohydrate foods preferred. In 1975 the food items liked the most were beef, potatoes, chicken, green beans, pie, pork, dried beans, lettuce salad, fish and cake. In 1985 the foods liked the most often were potatoes, chicken, beef, pie, green beans, fish, fruit, meat, lettuce salad and ice cream. Nine of the top eleven foods disliked in 1985 were vegetables, while eight of the top ten foods disliked in 1975 were vegetables. Items from the basic four food groups were chosen as "good for you" the most frequently in both the 1975 and 1985 surveys.

41 NAL Call No: 389.8 J824
Changes in total body water with age.

Schoeller, D.A.
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.
American journal of clinical nutrition v. 50 (5): p. 1176-1181. charts; 1989 Nov. Includes 15 references.

Language: English

Descriptors: Body water; Body composition; Labo-

ratory methods; Physiology; Men; Women; Elderly

Abstract: Extensive cross-sectional studies demonstrate a diminution of total body water in elderly and very old subjects. These findings are supported by less extensive longitudinal studies. Cross-sectional studies indicate that the decrease in total body water is mainly due to decreased intracellular water, but this is not supported by the findings of longitudinal studies. Despite the observed changes in total body water, both animal and human studies indicate little or no change in the relationship between total body water and fat-free mass with aging.

42 NAL Call No: HD9001.F65
Charting the course of change: what to expect as the food industry nears the 21st century.

Fair Lawn, N.J. : American Institute of Food Distribution, Inc; 1988 Apr09.
The Food distribution report (15): p. 4-6; 1988 Apr09.

Language: English

Descriptors: Consumption patterns; Trends; Food production; Food preferences; Consumer preferences; Prediction

Abstract: Recent trends in food consumption patterns are discussed relevant to projections of anticipated changes in food production and marketing most likely to occur by the end of the century. It is anticipated that over the next few years convenience stores and frozen microwaveables will be among the biggest competitors, that middle-class consumers are more likely to purchase prepared foods at a food store than white-collar workers, that teenagers are beginning to take nutrition seriously, and that elderly preferences will be more strongly felt with the anticipated growth in the US elderly population. A listing of anticipated changes is included.(wz)

43 NAL Call No: 448.8 L22
Cholesterol as risk factor for mortality in elderly women.

Forette, B.; Tortrat, D.; Wolmark, Y.
Boston, Mass. : Little, Brown and Company; 1989 Apr22.
The Lancet v. 1 (8643): p. 868-870. charts; 1989 Apr22. Includes 30 references.

Language: English

Descriptors: Cholesterol; Blood serum; Risks; Mortality; Carcinoma; Cardiovascular diseases; Longi-

Nutrition and the Elderly

tudinal studies; Elderly; Women

Abstract: This article reports the results of a study designed to assess the relation between mortality and total serum cholesterol in a group of elderly women. Mortality increased at both ends of the cholesterol distribution curve. The results suggest that cancer mortality does not account for the excess of deaths in elderly women with low cholesterol.

44 NAL Call No: RC952.A1G4
Cholesterol changes with aging: their nature and significance.

Kasim, S.
Cleveland : Modern Medicine Publications, Inc; 1987 Mar.
Geriatrics v. 42 (3): p. 73-76, 79, 82. ill., charts; 1987 Mar. Includes 27 references.

Language: English

Descriptors: Atherosclerosis; Aging; Cholesterol; High density lipoprotein; Triglycerides

Abstract: Total cholesterol has been directly correlated with the risk of coronary artery disease, while levels of high density lipoproteins have been shown to be inversely related with the disease. These relationships have been found, to a lesser extent, to occur in the elderly population as well. In general, plasma triglycerides, total cholesterol, and high density lipoproteins have been shown to increase progressively with age. Because data among studies vary, prudent management of cholesterol levels through proper diet, and appropriate medications, if necessary, are the suggested guidelines at this time.(lsp)

45 NAL Call No: TX341.N836
Chromium and nicotinic acid in glucose tolerance.
Van Nuys, Calif. : Nutrition & the M.D.; 1988 Mar.
Nutrition & the M.D. v. 14 (3): p. 1; 1988 Mar. Includes 1 references.

Language: English

Descriptors: Chromium; Niacin; Glucose tolerance; Elderly nutrition; Nutrient-nutrient interactions

Abstract: A brief report discusses the findings of a recently reported dietary study that assessed the effect of a possible interaction between chromium (Cr) and nicotinic acid (NA) on glucose tolerance in healthy elderly subjects. The combination of Cr and Na, but not either individually, produced a significant 7% decrease in the fasting glucose level, suggesting that sufficient amounts of both are needed to achieve optimum glucose metabolism.

46 NAL Call No: 389.1 W892
Clinical aspects of osteoporosis.

Duursma, S.A.; Raymakers, J.A.; Beresteyn, E.C.H. van; Schaafsma, G.

Basel : S. Karger; 1987.

World review of nutrition and dietetics v. 50: p. 92-121. ill., charts; 1987. Literature review. Includes 155 references.

Language: English

Descriptors: Osteoporosis; Clinical investigations; Bone mineralization; Bone resorption; Aging; Diagnosis; Therapeutic diets; Disease prevention; Treatment; Reviews

Abstract: A brief technical review surveys and discusses various clinical aspects associated with the prevention, causes, etiology, and treatment of osteoporosis. Specific attention is given to: Ca and bone homeostasis (including the maintenance and regulation of Ca and bone balance and the role of the hormonal system in Ca homeostasis); bone mass and the effects of aging (including physiological vs. pathological bone loss and the varied causes of osteoporosis); clinical approaches to diagnosing osteoporosis in patients; and the various treatments for osteoporotic patients (including dietary, physical activity, and drug therapies, and osteoporosis prevention). Practical measures and case studies are included.(wz)

47 NAL Call No: 389.8 J824
Clinical chemistry reference intervals for healthy elderly subjects.

Garry, P.L., Hunt, W.C.; VanderJagt, D.J.; Rhyne, R.L.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.

American journal of clinical nutrition v. 50 (5): p. 1219-1230. charts; 1989 Nov. Includes 8 references.

Language: English

Descriptors: Nutritional assessment; Blood chemistry; Reference standards; Elderly

Abstract: In 1979, 304 healthy elderly individuals in New Mexico were recruited for a longitudinal study of nutrition and aging. Repeat measurements on a yearly basis of commonly requested clinical chemistry analytes allowed the calculation of reference intervals, between and within-subject variance components, and percentiles for change in concentration between two yearly measurements. The latter was further divided into analytical and

Quick Bibliography Series

biological variance components. The upper 95th percentile for the difference between two yearly measurements, expressed as a percent of the population mean, ranged from 4% for Na⁺ to approximately 20% for total cholesterol and to greater than 90% for ferritin. Year-to-year differences attributable to the biological component ranged from a low of 2% of the population mean for Na⁺ to 70% for gamma-glutamyltransferase.

48 NAL Call No: TX341.C6
A closer look at the mature market.

Papa, A.
Washington, D.C. : Community Nutrition Institute;
1987 Mar05.
Nutrition week v. 17 (10): p. 4-5. charts; 1987
Mar05.

Language: English

Descriptors: Eating habits; Eating out; Elderly; Restaurants; Consumption patterns; Consumer behavior; Food preferences

Abstract: A brief report discusses factors that influence the food consumption patterns of the US "mature" population, defined as age 55 and above. It is pointed out that: mature consumers have money to spend; have fewer serious health problems than previously because of advances in medical technology and increased awareness of diet and exercise; spending for restaurant meals by mature consumers has increased; and that mature consumers prefer family-style restaurants. It also is noted that most (90%) mature consumers feel that they are informed about health and nutrition. Data are shown that indicate that mature consumers frequent fast food and moderately priced restaurants about as often, with less attention (10-15%) to attending fine restaurants.(wz)

49 NAL Call No: TX341.J6

A college-based nutrition counseling service for homebound elderly persons.

Fishman, P.; Van Marrelo, C.; Arthur, E.
Philadelphia, Pa. : George F. Stickley Company;
1987 Dec.

Journal of nutrition education v. 19 (6): p. 273-276.
ill., charts; 1987 Dec. Includes 6 references.

Language: English

Descriptors: New York; Nutrition education; Patient education; Home delivered meals; Diet counseling; Program evaluation

Abstract: A technical report discusses the design and implementation of a unique college-based

nutrition counseling service for educating homebound elderly subjects. Attention is given to the special opportunities and challenges associated with this service, operated as an adjunct to New York City's Meals on Wheels program. The procedures and assessment tools for this service are described and illustrated, and selected findings of a pilot evaluation project are discussed.

50 NAL Call No: TX341.C6
Community-based care should stress nutrition.

Bartlett, B.J.
Washington, D.C. : Community Nutrition Institute;
1989 Apr13.
Nutrition week v. 19 (15): p. 4-5; 1989 Apr13.

Language: English

Descriptors: Ohio; Nutrition programs; Community nutrition; Elderly nutrition; Home care; Home delivered meals; Elderly

Abstract: This article is an excerpt from the testimony by the Ohio Dietetic Association during a hearing before the Governor's Home and Community Care Council. The testimony emphasized that any program providing home care or community services to the elderly should include nutritional assessment and monitoring of nutritional status to ensure that nutritional needs are being met.

51 NAL Call No: QP141.A1N88

Comparative effects of linoleic acid and gamma-linolenic acid intake on plasma lipids and platelet phospholipids in elderly people.

Vericel, E.; Lagarde, M.; Mendy, F.; Courpron, P.; Dechavanne, M.

Elmsford, N.Y. : Pergamon Press; 1987 Jun.
Nutrition research v. 7 (6): p. 569-580. charts; 1987
Jun. Includes 24 references.

Language: English

Descriptors: Diet; Nutrient intake; Linoleic acid; Linolenic acid; Blood plasma; Lipids; Platelets; Phospholipids; Elderly

Abstract: Extract: It has been speculated that linoleic acid might be a precursor for platelet potentially anti-thrombotic polyunsaturated fatty acids like dihomo-gamma-linolenic acid. In elderly people, who are considered as a pre-thrombotic state, desaturation of linoleic acid may be diminished. To by-pass this rate limiting step, gamma-linolenic acid was given to elderly people and compared to linoleic acid with regard to the composition of plasma and platelet lipids. Sixteen old subjects took a daily supplement of 1 g of

Nutrition and the Elderly

linoleic acid plus 1 g of gamma-linolenic acid (primrose oil) or 2 g of linoleic acid (sunflower oil) for periods of two months in a cross over study. Increases in both dihomogamma-linolenic acid and arachidonic acid were noted in plasma lipids and platelet phospholipids after primrose oil supplementation. In contrast, sunflower oil did not induce such increases. Since dihomogamma-linolenic and arachidonic acid have opposite effects upon platelet functions, these findings could explain why primrose oil intake did not lead to any modification of platelet functions in elderly people.(author)

52 NAL Call No: 389.8 AM34
Comparing participants' and managers' perception of services in a congregate meals program. Harris, L.J.; Hodges, P.A.M.; Johnson, J.M.; Shifflett, P.A.
Chicago, Ill. : The Association; 1987 Feb.
Journal of the American Dietetic Association v. 87 (2): p. 190-195. charts; 1987 Feb. Includes 38 references.

Language: English

Descriptors: Elderly nutrition; Supplemental feeding programs; Consumers; Food quality; Nutrition education; Health services

Abstract: Extract: Managers and participants in the Title III C meals program were surveyed to determine differences in perception of meal and service quality. Manager vs. participant response significantly differed for food temperature, special diet needs, nutrition education, transportation, and accuracy of posted menus. Examination of discrepancies uncovered in this way in conjunction with foodservice monitoring will eliminate biased views and allow administrators to plan and evaluate the program more effectively.(author)

53 NAL Call No: 389.8 AM34
A comprehensive nutrition case management system. Clay, G.; Bouchard, C.; Hemphill, K.
Chicago, Ill. : The Association; 1988 Feb.
Journal of the American Dietetic Association v. 88 (2): p. 196-199. ill., forms; 1988 Feb. Includes 7 references.

Language: English

Descriptors: Nutritional assessment; Patient care; Nutritional intervention; Program development
Abstract: Extract: An innovative, state-of-the-art, comprehensive nutrition assessment and case man-

agement system has been developed by nutritionists (R.D.s) and public health nurses with the San Bernardino County Public Health Department in California. The system consists of three major components: flowsheets for the infant, the child, the perinatal client, family planning, and the adolescent/adult/aging client; a manual that defines and describes nutrition assessment guidelines and interventions; and an appendix that includes resource/reference information. Distinct advantages and unique features of this system include the following: (a) decrease in charting time; (b) precise documentation; (c) a workable diagnostic tool for implementing nutrition interventions; (d) a standard for consistent, quality nutrition care; (e) computer adaptability; and (f) easy accessibility for quality assurance audits. (author)

54 NAL Call No: 389.8 AM34
Computer use in Massachusetts elderly nutrition programs. Messina, M.S.; Balsam, A.L.
Chicago, Ill. : The Association; 1989 Oct.
Journal of the American Dietetic Association v. 89 (10): p. 1502-1504. charts; 1989 Oct. Includes 6 references.

Language: English

Descriptors: Massachusetts; Elderly nutrition; Nutrition programs; Computer applications; Questionnaires; Food service management; Health care; Home care; Hospitals; Dietitians

Abstract: The federally funded elderly nutrition program (ENP) began in 1972 under Title 7 of the Older Americans Act of that year. The program has as its primary mission the provision of food and nutrition services to persons 60 years old and older, targeting those in greatest social and economic need. This article reports the results of a questionnaire designed to describe to what extent ENPs make use of computer applications

55 NAL Call No: TX341.C6
Consumer may be poised on new era of activism. Washington, D.C. : Community Nutrition Institute; 1987 Jun11.
Nutrition week v. 17 (24): p. 4-5. charts; 1987 Jun11.

Language: English

Descriptors: Consumer preferences; Consumer behavior; Food preferences; Consumption patterns; Trends; Consumer protection
Abstract: A 1987 report of the Food Marketing In-

Quick Bibliography Series

stitute identifies new patterns of consumer practices that have significant implications for food policy and nutrition programs, and for food marketing. Data are presented that indicate that: more elderly people than younger people are very concerned about the nutritional content of the food they eat; consumers rely mostly on themselves that the food they purchase are safe; consumers frequently select foods to balance their family's diet and serve nutritional snacks; and that consumers are becoming increasingly concerned about the salt, mineral, and vitamin content of foods, and less concerned about sugar content. However, consumers are more concerned about each of these and other factors (fat and cholesterol content and the presence of preservatives and chemical additives) than they are about freshness, junk food, fiber content, and protein value. Issues that could lead to increased policy activism include: biotechnology, detection technology, preservation and packaging techniques, and new discoveries concerning relationships between diet and health and disease.(wz)

56 NAL Call No: 389.8 F7398

Consumer reaction to flavor innovations.

Horton, H.W.

Chicago, Ill. : Institute of Food Technologists; 1987 Jun.

Food technology v. 41 (6): p. 80-82, 84; 1987 Jun. Based on a paper presented during the symposium, "Flavor trends and technologies," sponsored by the Chicago Section of the Institute of Food Technologists, March 9, 1987, Chicago, IL.

Language: English

Descriptors: Flavorings; Consumer preferences; Food acceptability; Consumer behavior; Food industry; Food technology

Abstract: A brief commentary discusses factors influencing consumer satisfaction with food flavors and how rapid responses can be made by the food industry to consumer comments and complaints. Attention is focused on: the results of a consumer flavor survey affirming that blandness is out and flavor variety is in; the growth of consumer sophistication and experience, and how it affects food taste preferences; consumer health and fitness concerns; how manufacturers can keep abreast of changing consumer demands; guidelines for marketing the elderly and younger populations; and the influence of consumer nutrition education. The importance of the consumer's role in defining future trends for developing and marketing of new

flavors also is discussed.(wz)

57 NAL Call No: 449.9 AM3J
Continuance of elderly on home-delivered meals programs.

Frongillo, E.A. Jr; Williamson, D.F.; Roc, D.A.; Scholes, J.E.

Washington, D.C. : American Public Health Association; 1987 Sep.

American journal of public health v. 77 (9): p. 1176-1179. ill., charts; 1987 Sep. Includes 10 references.

Language: English

Descriptors: New York; Elderly nutrition; Home delivered meals; Supplemental feeding programs; Participation; Patient compliance; Surveys

Abstract: Extract: Data from 2,002 elderly receiving home-delivered meals in 23 counties of New York State in 1984-85 were analyzed using a baseline survey and a six-month follow-up survey. Survival analysis estimated that 37.4 per cent of the clients left the program by the follow-up. The initial drop-out rate was estimated to be 14.2 per cent per month; by six months the rate fell to 3.3 per cent. Whites left the program at a rate 2.6 times (95% CI = 1.9, 3.8) that of minorities. Recent discharge from hospital was associated with the highest relative probability of program discontinuance (2.7, 95% CI = 2.2, 3.4), followed by having cancer (2.1, 95% CI = 1.6, 2.8), and having a non-hip fracture (2.0, 95% CI = 1.3, 2.9). Many of those with cancer died, while those recently discharged and those with non-hip fractures were more likely to leave the program because they were again able to cook for themselves.(author)

58 NAL Call No: RM222.5.N87

The cost of home total parenteral nutrition.

Baptista, R.J.; Blackburn, G.L.; Champagne, C.D.; Miller, D.G.; Bistran, B.R.

Baltimore, Md. : Williams & Wilkins ; 1987 Feb. Nutrition in clinical practice : official publication of the American Society for Parenteral and Enteral Nutrition v. 2 (1): p. 14-22. ill., charts; 1987 Feb. Includes 9 references.

Language: English

Descriptors: Parenteral feeding; Elderly; Costs

Abstract: A survey assessed the differences in charges among 6 multiple home care vendors for hypothetical home total parenteral feeding and intravenous hydration patients. Comparisons of charges for self-mixed vs. pre-mixed solutions for

Nutrition and the Elderly

each of these therapies also was assessed. The cost estimates for self- and pre-mixed solutions for 2 different formulas are presented in detailed tables. Considerable variation was observed in the annual cost estimates among the vendors.(wz)

59

NAL Call No: 389.8 J823

Determination of zinc and copper absorption at three dietary Zn-Cu ratios by using stable isotope methods in young adult and elderly subjects.

August, D.; Janghorbani, M.; Young, V.R.

Mysore : Association of Food Scientists and Technologists; 1989 Dec.

Journal of food science and technology v. 50 (6): p. 1457-1463. charts; 1989 Dec. Includes 32 references.

Language: English

Descriptors: Zinc; Copper; Intestinal absorption; Nutrient intake; Ratios; Stable isotopes; Young adults; Elderly; Men; Women

Abstract: Nine young men and six elderly men and women ingested low-zinc, low-copper, and adequate diets for 2-wk periods. The Zn-Cu ratios averaged 2:1, 15:1, and 5:1. On day 8 of each period, subjects ingested doses of 70Zn and 65Cu so that net absorption could be determined. Zn absorption on the adequate diet was $39 \pm 3\%$ (average \pm SEM) in young subjects and $21 \pm 1\%$ (p less than 0.05) in the elderly subjects. During the low-Zn period, Zn absorption was higher as compared with that on the adequate diet in both young ($64 \pm 5\%$, p less than 0.05) and elderly subjects ($43 \pm 7\%$, p less than 0.05). Cu absorption on the adequate diet was $60 \pm 4\%$ and $53 \pm 2\%$ in young and elderly subjects, respectively. During the low-Cu period, Cu absorption was higher as compared with that on the adequate diet in the elderly subjects ($67 \pm 4\%$, p less than 0.05). Cu absorption was higher in six of seven young subjects on the restricted Cu diet (p greater than 0.05). The effect upon absorption of the change in the Zn-Cu ratio was less than the effect of dietary restriction.

60

NAL Call No: TX361.A3J6

Developing a continuum of community nutrition services: Massachusetts elderly nutrition programs.

Balsam, A.; Osteraas, G.

Binghamton, N.Y. : Haworth Press; 1987.

Journal of nutrition for the elderly v. 6 (4): p. 51-67. ill; 1987. Includes 18 references.

Language: English

Descriptors: Massachusetts; Elderly nutrition; Supplemental feeding programs; Federal programs; Program effectiveness; Nutrient requirements; Reports; Nutritional support; Community feeding centers

Abstract: A detailed report discusses the costs, characteristics, and effectiveness of elderly nutrition programs in Massachusetts. Specific attention is given to: the functioning and effectiveness of these programs for providing nutritional support for ambulatory elderly subjects having the greatest economic and social needs; serving the frail elderly; the benefits and effectiveness of luncheon clubs; requirements in addition to provision of "meals on wheels"; and the importance of nutrition and health education for elderly subjects. Special attention also is given to the cost of such programs, including meal costs and a breakdown of state nutrition program expenditures according to the diversification of income from federal and state programs and from donations.

61

NAL Call No: 389.8 AM34

Development and testing of a statistical and graphics-enhanced nutrient analysis program.

Manore, M.M.; Vaughan, L.A.; Lehman, W.

Chicago, Ill. : The Association; 1989 Feb.

Journal of the American Dietetic Association v. 89 (2): p. 246-250. charts; 1989 Feb. Includes 48 references.

Language: English

Descriptors: Nutrient contents; Computer software; Computer graphics; Nutrient data banks; Diet study techniques

Abstract: The authors have developed and tested a nutrient analysis program that will compute and present graphically summary statistics of population and population subgroup nutrient intakes. The program analyzes for 44 nutrients from 5,800 separate food items. Capabilities of the program include: storage of large numbers of diet records and evaluations of their nutrients; calculation of nutrient means and standard deviations; data sorting based on subject characteristics, such as age, sex, and supplement use; and generation of bar graphs and line plots for individual and/or group data. To test this computerized nutrient analysis program, two sets of 3-day diet records from 200 elderly individuals were analyzed. The program was then used to generate means, differences between means, and distribution frequencies of designated nutrients for various population subgroups (e.g., men greater than or equal to 65 years vs. men

Quick Bibliography Series

greater than or equal to 80 years) as well as comparisons with individual files (e.g., Mr. Smith vs. all men greater or equal to 65 years). The statistical and graphics capabilities also function within the context of recipe analysis and menu planning, which enhances the application of this program in institutional and community nutrition settings.

62 NAL Call No: 389.8 J824
Development of a diet history questionnaire for studies of older persons.

Hankin, J.H.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.

American journal of clinical nutrition v. 50 (5): p. 1121-1127; 1989 Nov. Includes 37 references.

Language: English

Descriptors: Elderly nutrition; Dietary history; Questionnaires; Research methodology; Data collection

Abstract: This paper describes the criteria of a simplified diet history method and the procedures for developing a dietary questionnaire for studies of older persons. Suggestions are included on selecting food items representative of the population's dietary patterns, designing objective methods to estimate quantitative and frequency intakes of food items, and collecting information on other dietary practices. Potential problems in dietary surveys among older persons are discussed, along with their prevention or alleviation.

63 NAL Call No: 389.8 N953
Diet and maintenance of mental health in the elderly.

Hodkinson, H.M.

Washington, D.C. : Nutrition Foundation; 1988 Feb.

Nutrition reviews v. 46 (2): p. 79-82; 1988 Feb. Includes 7 references.

Language: English

Descriptors: Mental health; Diet; Elderly; Neurotransmitters; Metabolism; Depression

Abstract: A brief report discusses the role of diet in maintaining mental health in old age. Topics include: dietary nutrients that serve as neurotransmitter precursors; nutrients that affect neurotransmitter metabolism and general brain metabolism; and the effect of changes in the intake of nutrients on the development of confusional states of mind. Information gaps requiring further research study also are discussed.

64 NAL Call No: S22.C6
Diet and senile dementia.

Ames, Iowa : The Council; 1987 Mar.

Council for Agricultural Science and Technology report v. 111: p. 55-56. charts; 1987 Mar. Includes 22 references.

Language: English

Descriptors: Senility; Alzheimer's disease; Nutrient deficiencies; Choline; Nutrient excesses; Aluminum; Elderly

Abstract: A brief task force report discusses 2 dietary factors that may be related to the etiology and prevention of the type of senile dementia known as Alzheimer's disease, viz.: inadequate intakes of choline; and excessive intakes of aluminum (Al). Data on excessive Al intakes from antacids and analgesics relative to typically safe dietary intakes of Al are included. It is emphasized that evidence relating diet to the incidence or treatment of Alzheimer's disease is very limited.(wz)

65 NAL Call No: 389.8 N953
Diet, exercise, and chronic disease patterns in older adults.

Smith, E.L.; Smith, P.E.; Gilligan, C.

Washington, D.C. : Nutrition Foundation; 1988 Feb.

Nutrition reviews v. 46 (2): p. 52-61. charts; 1988 Feb. Literature review. Includes 96 references.

Language: English

Descriptors: Cardiovascular diseases; Diabetes; Osteoporosis; Aging; Diet; Exercise; Disease prevention

Abstract: A technical review considers the role of physical activity and diet in elderly subjects in maintaining the cardiovascular system, in lipid metabolism, in adult-onset diabetes, and in the skeletal system. Particular attention is given to changes in cardiovascular function during aging; plasma lipoproteins; glucose tolerance; osteoporosis; dietary considerations; and the effects of inactivity and physical activity. It is concluded that diet and exercise are being related increasingly to the prevention/delay of diabetes, atherosclerosis, and osteoporosis onset during aging.

66 NAL Call No: QP751.D53
Dietary fat requirements in health and development.

Beare-Rogers, Joyce

Champaign, Ill. : American Oil Chemists Society;

Nutrition and the Elderly

1988.

v, 206 p. : ill. ; 24 cm. includes bibliographies.

Language: English

Descriptors: Lipids in human nutrition; Fat; Fat metabolism; Dietary fat; Nutrition; Fatty acids

Abstract: This publication is a collection of works on fat requirements in development and health. The role of fats during human development and throughout a lifespan are discussed. Topics include: suitable fat foundations for infant feeding; lipid digestion in the developing infant; fat effects on fatty acids and cholesterol metabolism in animal experiments; the biochemistry and role of (n-3) fatty acids in the brain and retina; dietary factors in immune responsiveness; aging and nutritional requirements of essential fatty acids; and challenges for lipid nutritionists.

67

NAL Call No: RC682.A37

Dietary fats and the aging adult.

Willcox, K.K.; Lei, K.Y.

Boca Raton, Fla. : CRC Press; 1987.

Nutrition and heart disease / editor, Ronald R. Watson. v. 1 p. 21-28; 1987. Literature review. Includes 56 references.

Language: English

Descriptors: Source fat; Cardiovascular diseases, Elderly nutrition; Food and nutrition controversies; Risks; Lipid metabolism; Aging; Reviews

Abstract: A technical review focuses on dietary fat consumption by Americans, with particular emphasis on elderly intakes. The review addresses the association between dietary fat and cardiovascular disease (CVD), as well as age-related changes in dietary fat metabolism. It is emphasized that the type of fat ingested appears to be equally as if not more important than the quantity of fat consumed with respect to hyperlipidemia, atherosclerosis, and heart disease risk. It is concluded that conflicting opinions on the role of fats in CVD make the diet-heart hypothesis a controversial issue.

68

NAL Call No: 389.8 AM34

Dietary habits, weight history, and vitamin supplement use in elderly osteoarthritis patients.

White-O'Connor, B.; Sobal, J.; Muncie, C.L. Jr
Chicago, Ill. : The Association; 1989 Mar.

Journal of the American Dietetic Association v. 89 (3): p. 378-382. charts; 1989 Mar. includes 33 references.

Language: English

Descriptors: Arthritis; Cartilage; Food consumption; Food groups; Physical activity; Weight gain; Weight losses

Abstract: Osteoarthritis is a chronic disabling disease in the elderly, but few studies have examined nutritional parameters of osteoarthritis patients. For 82 ambulatory elderly osteoarthritis patients, a registered dietitian assessed the following: consumption of 72 food items, using a food-frequency questionnaire; weight history, by measuring current weight and asking the weight at age 20, maximum adult weight, and minimum adult weight; dietary habits; and vitamin supplement consumption. Joint pain and activities of daily living (ADL) were assessed by a physician. On the basis of the Four Food Group guidelines, dietary intakes were suboptimal in the dairy and grain groups, which are important sources of calcium, vitamin D, thiamin, iron, and riboflavin. Eighty percent of the sample were obese (BMI greater than or equal to 27). The average weight change since early adulthood was a gain of 59 lb. Current joint pain and ADL restrictions were not related to obesity or weight gain. Vitamin/mineral supplements were consumed by 37% of the sample.

69

NAL Call No: TX341.C6

Dietary info. lacking for elderly Americans.

McGandy, R.B.

Washington, D.C. : Community Nutrition Institute; 1987 Mar19.

Nutrition week v. 17 (12): p. 4-5; 1987 Mar19.

Language: English

Descriptors: USA; Elderly nutrition; Food and nutrition controversies; Nutritional state; Nutrient requirements; Supplemental feeding programs; Income

Abstract: Two papers presented at a 1986 USDA conference argue the need for further studies to assess and quantify dietary information concerning the US elderly. The first paper addressed various factors influencing the quality of elderly diets; the second paper compared income trends between the young and the elderly and discussed the effect of changes in the economy and future income prospects.(mjs)

70

NAL Call No: 389.8 J824

Dietary intakes and biochemical indicators of nutritional status in an elderly, institutionalized population.

Sahyoun, N.R.; Otradovec, C.L.; Hartz, S.C.; Jacob, R.A.; Peters, H.; Russell, R.M.; McGandy,

Quick Bibliography Series

R.B.

Bethesda, Md. : American Society for Clinical Nutrition; 1988 Mar.

American journal of clinical nutrition v. 47 (3): p. 524-533. charts; 1988 Mar. Includes 39 references.

Language: English

Descriptors: Nutritional state; Elderly; Nutrient intake; Anthropometric dimensions; Nutritional assessment

Abstract: Extract: A dietary and biochemical assessment of the nutritional status of 260 elderly men and women, 60-101 y (average 80.5 y), was conducted in 15 long-term-care facilities in the Boston area. Subjects were free of clinically apparent terminal or wasting illness. Nutrient intakes were comparable to those in a simultaneously studied free-living population as were most biochemical markers of nutrient status. Although no specific nutrient deficiencies were identified, blood levels of vitamin A and retinol-binding protein in males and of zinc in both sexes were lower in this institutionalized group than in the free-living subjects. Hematologic indices, albumin, prealbumin, and transferrin levels were also lower than in non-institutionalized elderly populations. These differences may reflect the greater prevalence of chronic diseases and medication use in a long-term-care population. However, there is no evidence that institutionalization in itself leads to impairment of nutritional status.(author)

71

NAL Call No: 389.8 F7332

Dietary patterns and nutritional status of low income, free-living elderly.

Vaughan, L.A.; Manore, M.M.

Chicago, Ill. : National Live Stock and Meat Board; 1988 Nov.

Food and nutrition news v. 60 (5): p. 27-30. charts; 1988 Nov. Includes 28 references.

Language: English

Descriptors: Nutritional state; Nutrient intake; Vitamin And mineral supplements; Poultry meat; Fish; Nutritional assessment; Low income groups; Elderly

Abstract: The U.S. population is rapidly aging, and the proportion of elderly and very elderly Americans continues to increase. Inadequate dietary intakes and low nutritional status have been identified within elderly populations. Research suggests the elderly are at risk for inadequate consumption of dietary vitamin B-6, folacin and zinc;

vitamin B-12 and iron are rarely a problem. Biochemical data indicate vitamin B-6 status is frequently low in the elderly, while folacin and vitamin B-12 status is generally adequate. Iron status indices in the elderly are more difficult to interpret, but suggest a marginal adequacy. The nutritional status of the elderly can be influenced by their dietary patterns, supplementation practices, medication use and state of health.

72

NAL Call No: 448.8 N442

Dietary potassium and stroke-associated mortality.

Khaw, K.T.; Barrett-Connor, E.

Boston, Mass. : Massachusetts Medical Society; 1987 Jan 29.

New England journal of medicine v. 316 (5): p. 235-240. ill., charts; 1987 Jan 29. Includes 54 references.

Language: English

Descriptors: Diet; Stroke; Potassium; Mortality; Hypertension; Longitudinal studies

Abstract: A study using data from a 12-year longitudinal, regional study of 859 middle-aged to elderly subjects (ages 50-79) tested the hypothesis that a high dietary potassium (K) intake reduces stroke incidence. Statistical analysis of K intake data indicated that risk of stroke-related mortality was reduced by 40 per cent by an increase in K intake of 10 mmol/day. This effect appeared to be unaffected by accepted cardiovascular disease risk factors (e.g., age, sex, blood pressure level, obesity, cigarette smoking, blood glucose level) and other dietary intakes (total energy, fat, protein, fiber, alcohol, Ca, Mg). The results supported the hypothesis that increased K intakes protect against stroke-related mortality.(wz)

73

NAL Call No: 389.8 J82

Dietary restriction and aging: historical phases, mechanisms and current directions.

Walford, R.L.; Harris, S.B.; Weindruch, R.

Bethesda, Md. : American Institute of Nutrition; 1987 Oct.

The Journal of nutrition v. 117 (10): p. 1650-1654. ill., charts; 1987 Oct. Includes 51 references.

Language: English

Descriptors: Calorie-restricted diets; Aging; Health promotion; Longevity; Models; Animal experiments

Abstract: A technical overview discusses the historical justification of energy intake restriction (EIR),

Nutrition and the Elderly

in the absence of essential nutrient deficiencies, for retarding aging and disease susceptibility and extending life span, the potential mechanisms involved, and current research directions aimed at improving the understanding of this effect. Potential mechanisms involved in this effect may include effects on the immune defense system, cell proliferation potential, metabolic rate, DNA repair, protein biosynthesis/turnover, and a control of free radical scavengers. The utility of the EIR model for examining single vs. multiple factorial theories of aging also is discussed.

74 NAL Call No: R31.B55
Dietary salt affects biochemical markers of resorption and formation of bone in elderly women.

McParland, B.E.; Goulding, A.; Campbell, A.J.
London : British Medical Association; 1989 Sep30.
BMJ : British medical journal v. 299 (6703): p. 834-835; 1989 Sep30. Includes 5 references.

Language: English

Descriptors: Sodium chloride; Intake; Diet; C-amp; Hydroxyproline; Vitamin D; Bone resorption; Elderly; Women

Abstract: High dietary intakes of sodium chloride increase calciuria and are considered a risk factor for osteoporosis. We investigated the short term effects of salt intakes of 70 and 170 mmol/day on variables of resorption of bone (urinary hydroxyproline excretion) and formation of bone (serum osteocalcin concentration) in elderly women.

75 NAL Call No: 389.8 AM34
Dietary status of Seventh-Day Adventist vegetarian and non-vegetarian elderly women.

Nieman, D.C.; Underwood, B.C.; Sherman, K.M.; Arabatzis, K.; Barbosa, J.C.; Johnson, M.; Shultz, T.D.

Chicago, Ill. : The Association; 1989 Dec.
Journal of the American Dietetic Association v. 89 (12): p. 1763-1769. charts; 1989 Dec. Includes 46 references.

Language: English

Descriptors: Nutrient intake; Vegetarianism; Nutrition surveys; Food preferences; Elderly; Women

Abstract: The purpose of this study was to investigate nutrient intakes of Seventh-Day Adventist elderly women who were similar in many demographic and life-style factors except for choice of diet. Twenty-three vegetarian and 14

non-vegetarian elderly women (mean +/- standard error ages 72.2 +/- 1.3 and 71.1 +/- 1.4 years, respectively) were recruited on the basis of several selection criteria, including race, religion, education, geographic area, Quetelet index, self-reported absence of major chronic disease and use of medications, and physical activity. Average years +/- SE of adherence to dietary regimens were 47.0 +/- 2.9 and 71.2 +/- 1.4 in the vegetarian and non-vegetarian groups, respectively. Results from analysis of 7-day food records showed that vegetarians consumed significantly less cholesterol, saturated fatty acids, and caffeine but more carbohydrate, dietary fiber, magnesium, vitamins E and A, thiamin, pantothenic acid, copper, and manganese than non-vegetarians (p less than .05). On the basis of group means, 67% of the Recommended Dietary Allowance was met for all nutrients except zinc and vitamin D in both groups, and vitamins E-6, folacin, and vitamin E in the non-vegetarians. Compared with non-vegetarians, vegetarians had significantly lower serum glucose (5.18 +/- 0.11 vs. 4.65 +/- 0.09 mmol/L), low-density-lipoprotein cholesterol (4.08 +/- 0.25 vs. 3.34 +/- 0.19 mmol/L), and total cholesterol levels (6.46 +/- 0.27 vs. 5.62 +/- 0.21 mmol/L) (p less than .05). In summary, when healthy elderly vegetarian women were compared with closely matched non-vegetarian peers, the vegetarian diet was associated with improved nutrient intake and associated reductions in blood glucose and lipid levels.

76 NAL Call No: QP141.A1N88

Dietary survey of a large random sample of elderly people: energy and nutrient intakes.

Horwath, C.C.

Elmsford, N.Y. : Pergamon Press; 1989 May.
Nutrition research v. 9 (5): p. 479-492. charts; 1989 May. Includes 62 references.

Language: English

Descriptors: South australia; Nutrient intake; Caloric intake; Dietary surveys; Food consumption; Protein intake; Dietary carbohydrates; Fat consumption; Vitamins; Minerals; Food fiber; Elderly; Men; Women

Abstract: The average daily intakes of energy and nutrients were estimated in a randomly selected group of 2195 people aged 65 years and over residing in Adelaide, South Australia, by means of a semi-quantitative food frequency questionnaire. Mean intakes for most nutrients were equal to or above the recommended intakes. The proportion of people with intakes below two-thirds of the

Quick Bibliography Series

RDA, however, was highest for folate, calcium, magnesium, copper, and zinc, and varied from approximately 10 to 35 per cent of the total group. Comparisons with the results of other large surveys of noninstitutionalized elderly people in Boston and Gothenburg, Sweden, showed some variation in total intakes, but many similarities in patterns of nutrient density and those nutrients least adequately supplied.

77 **NAL Call No: 389.8 F7332**
The dining environment and food intake of the elderly in long-term care.

Doyle, B.

Chicago, Ill. : National Live Stock and Meat Board; 1989 Jan.

Food and nutrition news v. 61 (1): p. 5-6. ill; 1989 Jan. Includes 8 references.

Language: English

Descriptors: Food intake; Long term care; Malnutrition; Dining facilities; Food serving methods; Interior design; Physical environment; Sensory disorders; Elderly

Abstract: Appropriate environmental adaptations in long-term care dining settings can improve the ongoing nutrient intake of the older resident. Improvement in intake not only improves overall nutritional status but also makes mealtime an anticipated and positive "dining experience," rather than just a time to consume food.

78 **NAL Call No: 389.8 J824**
Discriminant biochemical markers for evaluating the nutritional status of elderly patients in long-term care.

Kergoat, M.J.; Lerclerc, B.S.; PetitClerc, C.; Imbach, A.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Nov.

American journal of clinical nutrition v. 46 (5): p. 849-861. ill., charts; 1987 Nov. Literature review. Includes 65 references.

Language: English

Descriptors: Elderly nutrition; Nutritional state; Nutritional assessment; Patient care; Carotenes; Creatinine; Malnutrition; Anthropometric dimensions; Diagnosis

Abstract: Extract: To determine the most discriminant serum markers of protein-energy status in elderly patients, we performed a discriminant analysis of 85 subjects grouped according to triceps skinfold and midarm circumference values as com-

pared with reference percentiles. Results indicated that neither the classic serum indices of nutritional assessment nor retinol-binding protein can predict undernutrition. However, creatinine, urea, carotene, complement C3, and prealbumin included in a function enabled high discrimination between groups: 68% of subjects in 0-5th percentile for triceps skinfold and 75% of subjects in 0-5th percentile for midarm circumference are correctly predicted. Lower serum concentration was found in the lower anthropometric percentiles except for serum carotene, which showed an inverse relation not explained by diet. We found that nutritional alterations exist in hospitalized elderly patients. We emphasize the importance of considering several biochemical markers for detection of malnutrition and the pertinency of further exploration of serum carotene profiles in undernourished elderly patients.(author)

79 **NAL Call No: HQ1060.J68**
Does nutrition affect surgical outcome?

Cooper, J.K.

New York : Elsevier Science Publishing Company; 1987 Mar.

Journal of the American Geriatrics Society v. 35 (3): p. 229-232; 1987 Mar. Includes 33 references.

Language: English

Descriptors: Supplements; Malnutrition; Surgery; Nutritional state

Abstract: A review article addresses the question of whether a higher nutritional status is associated with an improved surgical outcome in elderly patients, concluding that it is difficult to diagnose protein malnutrition in the elderly due to lack of well-defined norms and anthropometric characteristics change with age in a non-predictable variable way. It also is pointed out that while low serum albumin have been correlated with poor surgical outcome and nutritional supplements can reverse some nutritional deficiencies, the value of supplementation to elderly surgery candidates who are not severely malnourished is unproven and the benefits of nutritional intervention must be weighed against costs of delayed surgery and hospital stay.(wz)

80 **NAL Call No: aHD9001.N275**
Domestic food programs: an update.

Matsumoto, M.

Washington, D.C. : The Service; 1987.

National food review NFR - United States Department of Agriculture, Economic Research Ser-

Nutrition and the Elderly

vice (NFR-38): p. 24-25. ill., charts; 1987.

Language: English

Descriptors: USA; Supplemental feeding programs; Federal programs; Food stamp program; School lunches; Elderly nutrition; Program participants; Costs; Trends

Abstract: A technical report compares food program participation and costs for January-March 1987 with the same period for 1986. Summary data and findings are presented and discussed concerning the US Food Stamp Program, child nutrition programs (National School Lunch Program; School Breakfast Program; Child Care Food Program; Special Milk Program), supplemental food programs (WIC; the Commodity Supplemental Food Program; Elderly Feeding Pilot Project), and food distribution programs (Food Distribution Program on Indian Reservations; Nutrition Program for the Elderly; Temporary Emergency Food Assistance Program). Data for 1985 also are included.

81 NAL Call No: 389.8 J824
Dose-response relationships regarding vitamin B-6 in elderly people: a nationwide nutritional survey (Dutch Nutritional Surveillance System).

Lowik, M.R.H.; Berg, H. van den; Westenbrink, S.; Wedel, M.; Schrijver, J.; Ockhuizen, T. Baltimore, Md. : American Society for Clinical Nutrition; 1989 Aug.

American journal of clinical nutrition v. 50 (2): p. 391-399. charts; 1989 Aug. Includes 40 references.

Language: English

Descriptors: Netherlands; Pyridoxine; Nutrient intake; National surveys; Nutritional state; Vitamin supplements; Food intake; Elderly; Men; Women

Abstract: The dietary intake and biochemical status of vitamin B-6 in 476 apparently healthy Dutch elderly people (aged 65-79 y), who were not using drugs known to affect vitamin B-6 metabolism, were evaluated. Intake of vitamin B-6 per gram protein was related to biochemical data, namely plasma pyridoxal 5'-phosphate (PLP) and cofactor stimulation of aspartate aminotransferase in erythrocytes (AST-AC). Based on a cutoff point of 2.02 for AST-AC, approximately 9% of the elderly people not using vitamin B-6 supplements had a marginal vitamin B-6 status. About 7% were using vitamin B-6 supplements. Dietary intake of vitamin B-6 per gram protein was negatively related to AST-AC. Vitamin B-6 intakes per gram protein

higher than 0.020 mg were necessary to ensure an AST-AC value less than 2.02. At high PLP values AST-AC hardly varied. The results seem to indicate a higher requirement of vitamin B-6 in elderly people than in younger adults.

82 NAL Call No: RM302.D76
Drug and nutrient interactions in elderly cardiac patients.

Roe, D.A.

New York, N.Y. : Alan R. Liss; 1988.

Drug-nutrient interactions v. 5 (4): p. 205-212. charts; 1988. Includes 33 references.

Language: English

Descriptors: Cardiovascular diseases; Drug therapy; Nutrient-drug interactions; Diuretics; Hypertension; Wheat bran; Potassium deficiency; Digoxin; Purgatives; Elderly

Abstract: In the elderly, drug treatment is used for the prevention and control of cardiovascular disease. Prevention of cardiovascular disease includes management of hypercholesterolemia to reduce risk of myocardial infarction, treatment of acute myocardial infarction to reduce the risk of recurrence, and treatment of hypertension to reduce the risk of hypertensive heart disease and stroke. Management of chronic cardiovascular disease with permanent disability is the major therapeutic goal. The most commonly treated disorders of the heart and peripheral vascular system include congestive heart failure, arrhythmias, angina, and thromboses. Reduction of plasma lipid levels may be undertaken by drug therapy as a means of preventing myocardial infarction.

83 NAL Call No: RM302.D76
Drug and nutrient interactions in the elderly diabetic.

Roe, D.A.

New York, N.Y. : Alan R. Liss; 1988.

Drug-nutrient interactions v. 5 (4): p. 195-203. charts; 1988. Literature review. Includes 29 references.

Language: English

Descriptors: Diabetes; Nutrient-drug interactions; Drug combinations; Nutrient deficiencies; Hypoglycemia; Hyperglycemia; Appetite; Weight gain; Elderly

Abstract: Elderly diabetics take more drugs than other groups of elderly patients. Their multiple drug use is largely explained by the drugs that they take for complications of their primary disease;

Quick Bibliography Series

these include cardiovascular drugs for macrovascular disease and antibiotics for secondary infections. They also take more drugs for control of other conditions that are etiologically associated with the development and progression of their diabetes, including antihypertensive agents, antilipemic agents and steroids, and nonsteroidal antiinflammatory drugs (NSAIDs), which are taken for relief of joint pain that is intensified by arthritic joints bearing excess weight. Drugs taken by elderly diabetics that contribute to the high prevalence of drug-nutrient interactions include those taken as antidiabetic agents, including both insulin and sulfonylureas as well as calcium channel blockers; they also include thiazides, loop diuretics, sulfa drugs, cephalosporin antibiotics, tetracyclines, antifungal agents, cholestyramine and colestipol, niacin, prednisone and other corticosteroids, and NSAIDs. These drugs and drug combinations contribute to the risk of hyperglycemia, which can cause nonketotic hyperglycemia in the elderly; to the risk of hypoglycemia, which in the elderly carries the risk of inducing pseudo-stroke; to the risk of drug-induced nutritional deficiencies from antilipemics and cephalosporins, which can induce vitamin K deficiency; to the risk of acute incompatibility reactions, including flush reactions from chlorpropamide, niacin, and calcium channel blockers; and to the risk of edema, anemia, and hyperkalemia from NSAIDs. The risk of these adverse outcomes can be minimized by carefully pruning drug lists, by discontinuing medications when they are no longer needed, by monitoring drug times relative to meal times, and by monitoring the electrolyte levels, blood count, and nutrient levels in elderly diabetics. Promoting abstinence also serves to reduce risk.

84 NAL Call No: Audiocassette no.78
Drug and nutrient interactions information and instructions for dietitians and their patients.
Roe, Daphne A.,1923-
American Dietetic Association
Chicago, Ill. : American Dietetic Association, Publications Dept; 1987.
1 sound cassette (41 min.) : 1 7/8 ips, mono. + 1 study guide. (Audio cassette series ; 1987 ACS 6).

Language: English

Descriptors: Drug-nutrient interactions; Drug interactions; Audiovisual aids; Nutrient-drug interactions; Elderly; Children; Health hazards; Patient education; Dietitians; Guidelines

Abstract: An audio cassette tape and an accompanying study guide provides basic information and guidelines to aid dietitians in understanding and minimizing drug-nutrient interactions in clinical settings in particular patient groups and in individual elderly patients and other patients taking medication, and to provide dietitians with advice on appropriate intervention. The cassette tape covers: definitions of drug-nutrient interactions; the classification of such interactions by adverse effects and by risk level; specific examples of drug-nutrient interactions; guidelines for controlling common drug-nutrient interactions in elderly patients, diabetics, and elderly patients with neurological or mental health problems (as well as for situations involving development disorders and anorexia in children and problems elderly patients face from over-the counter drugs and the mass media); information on foods, formulas, and diets that reduce the effectiveness of prescribed drugs (including dietary changes, dietary factors, and the use of nasogastric tubes); and advice concerning the monitoring of drug-nutrient interactions for their control. The accompanying study guide provides tabular summaries of specific food-drug incompatibilities, the chemical and brand names of various drugs, factors that influence drug absorption, and information on the type of nutrient interaction and its risk for drugs used for infants through the elderly.

85 NAL Call No: TX341.N836
Drug-induced malnutrition in the geriatric patient.
Blumberg, J.B.
Van Nuys, Calif. : Nutrition & the M.D.; 1987 Aug.
Nutrition & the M.D. v. 13 (8): p. 1-3; 1987 Aug.
Includes 5 references.

Language: English

Descriptors: Elderly nutrition; Nutrient-drug interactions; Patient care; Guidelines
Abstract: A brief technical review highlights the effects of prescribed drugs on nutritional status and the effect of nutrient intake on the effectiveness of such medications in elderly patients. Specific attention is given to: alterations of food intake by drugs that influence appetite, taste, and smell, including their effect on nutrient absorption, and on nutrient metabolism and excretion. Practical guidelines for reducing the risk of drug-induced malnutrition risk in the elderly are included.(wz)

86 NAL Call No: TX341.N98

Nutrition and the Elderly

Eating for your 80s.

Liebman, B.; Davis, R.

Washington, D.C. : Center for Science in the Public Interest; 1987 Nov.

Nutrition action health letter v. 14 (9): p. 1, 4-7. ill.; 1987 Nov. Includes 6 references.

Language: English

Descriptors: Elderly nutrition; Exercise; Health promotion; Aging; Guidelines

Abstract: A report for aging people discusses how dietary modifications (including the use of the Pritikin diet) and exercise can be used to promote and extend a sound health status into their 80's. A number of examples and guidelines are given. Specific attention is given to: methods for unclogging arteries; the positive effects of exercise; increasing fiber intakes; reduction of diabetes; the use of sensible vitamin-mineral supplements; the dual need for both calcium and vitamin D; and unrealistic RDA's. Practical suggestions for elderly people are included.(wz)

87

NAL Call No: TX361.A3E3

Eating the low fat way a nutrition program for older adults, with home and quantity-size recipes. Philadelphia Health Management Corporation, Thomas Jefferson University, Dept. of Geropsychiatry

Philadelphia, PA : Philadelphia Health Management Corporation : Thomas Jefferson University; 1987.

139 p. ; 28 cm. "The nutrition program described ... is part of a larger project, "Exercise and Nutrition Program for the Elderly", which was developed by the Philadelphia Health Management Corporation and Thomas Jefferson University, Department of Geropsychiatry."--Pref. Bibliography: p. 137-139.

Language: English

Descriptors: Low-fat diet--Recipes; Aged--Nutrition; Popular works; Elderly; Nutrition programs

Abstract: This manual describes a nutrition program for the elderly. This program is part of a larger project: "Exercise and Nutrition Program for the Elderly" developed by Thomas Jefferson University's Dept. of Geropsychiatry and the Philadelphia Health Management Corporation. It is intended to alleviate or improve some of the health problems associated with aging and to improve the functional ability and independence of residents in nursing homes. This nutrition program recom-

mends changes in eating habits and includes: nutritional guidelines for changing eating habits; and how to reduce sodium in the diet. Recipes, references, appendices, illustrations, and a glossary are included.

88

NAL Call No: TX715.W5

Eating well when you just can't eat the way you used to.

Wilson, Jane Weston; Karas, G. Brian
New York : Workman Pub; 1987.

x, 390 p. : ill. (some col.) ; 26 cm. Includes index. Bibliography: p. 376-378.

Language: English

Descriptors: Cookery; Aged--Nutrition; Recipes; Dietary guidelines; Elderly nutrition

Abstract: This text is a guide to eating and living well for those who are fifty and over. The aim of this book is to show how to feast every day (but lightly), how to entertain often, how to splurge on occasion; and how to do it all in a healthy way. Food shopping, a philosophy of eating, suggested menus, food preparation and a large section of recipes are included.

89

NAL Call No: TX945.S344

Education, training, and development of the older worker.

DeMicco, F.J.

Englewood, Colo. : American School Food Service Association; 1988.

School foodservice research review v. 12 (1): p. 6-12. charts; 1988. Includes 34 references.

Language: English

Descriptors: School food service; Personnel; Training; Demography; Aging; Labor force; Employer employee relationships; Attitudes; Elderly

Abstract: Demographic trends will have a significant impact on the food service labor force. With fewer teenagers available, the commercial sector (that is, fast food restaurants) is experiencing acute labor shortages. This trend will impact the institutional sector (that is, school food services) as commercial and institutional employers compete for a similar labor pool. The older worker will be a valuable resource in light of a shrinking labor pool and increased competition. Stereotypes regarding the training, education, and development of older workers are pervasive. To achieve financial success in their food service operations, employee training, education, and development programs are necessary. Therefore, managers will have to overcome

Quick Bibliography Series

stereotype barriers and develop effective training and educational programs for older employees. This paper examines the age-related stereotypes regarding education, training, and development of the older worker.

90 NAL Call No: 389.8 AM34
Effect of household size on nutritional patterns among the Dutch elderly.

Westenbrink, S.; Lowik, M.R.H.; Hulshof, K.F.A.M.; Kistemaker, C.

Chicago, Ill. : The Association; 1989 Jun.

Journal of the American Dietetic Association v. 89 (6): p. 793-799. charts; 1989 Jun. Includes 25 references.

Language: English

Descriptors: Netherlands; Nutritional state; Food consumption; Nutrition surveys; Eating patterns; Risks; Family size; Households

Abstract: Elderly persons are vulnerable to nutritional risks. Factors such as living alone make the elderly prone to poor nutritional status as a result of inadequate diets. The results of a Dutch national nutrition survey of food consumption and nutritional status among 539 independently living, apparently healthy elderly persons, aged 65 to 79 years, show that in general the intake of macronutrients deviates from the Dutch recommendations. In particular, saturated fat intake and P:S ratio were unfavorable, especially for the solitary elderly persons (saturated fat intake 19.0 +/- 3.2% and 17.7 +/- 3.3% total energy for men and women, respectively). Except for vitamin B-6 (about 92% of the recommendation on a group level for men and 78% for women), the intake of micronutrients studied is considered to be adequate. Elderly women living alone had a higher calcium:phosphorus ratio (0.76 +/- 0.12), lower meat consumption (90 +/- 31 gm), and a lower mean HDL-cholesterol concentration (1.22 +/- 0.28 mmol/L) in serum than non-solitary elderly women (Ca:P ratio 0.73 +/- 0.12, meat consumption 100 +/- 36 gm, HDL-cholesterol concentration 1.45 +/- 0.38 mmol/L). Potatoes, vegetables, and meat were more often prepared for more than one day at a time by elderly persons living alone than by the non-solitary elderly.

91 NAL Call No: RC620.A1N8
Effect of vitamin E supplement intake on HDL and total cholesterol in the elderly.

Jacques, P.F.; Hartz, S.C.; McGandy, R.B.; Russell, R.M.; Jacob, R.A.

Stonham, Mass. : Butterworth Publishers; 1988 Feb.

Nutrition reports international v. 37 (2): p. 363-370. charts; 1988 Feb. Includes 25 references.

Language: English

Descriptors: Vitamin E; Elderly nutrition; Vitamin supplements; High density lipoprotein; Cholesterol

Abstract: Extract: The relationships of vitamin E supplement use with plasma HDL cholesterol (HDL-C) and total cholesterol (T-C) were examined in 252 males and 465 females, age 60 to 98 years. The relative difference between those subjects consuming greater than or equal to 10 times the recommended dietary allowances (RDA) for vitamin E from supplements and those consuming less than the RDA from supplements was 3% for HDL-C and 2% for T-C. Neither of these differences was statistically significant (p greater than 0.05). The results suggest that vitamin E intake does not influence HDL-C or T-C. The present study, because of the large sample size, overcomes the problem of low statistical power to detect small, but clinically important, differences in cholesterol levels between those with high and low vitamin E intakes.(author)

92 NAL Call No: TX361.A3J63
Effect of zinc supplementation on the development of cardiovascular disease in the elderly.

Hale, W.E.; May, F.E.; Thomas, R.G.; Moore, M.T.; Stewart, R.B.

Binghamton, N.Y. : Haworth Press; 1988.

Journal of nutrition for the elderly v. 8 (4): p. 49-57; 1988. Includes 8 references.

Language: English

Descriptors: Zinc; Mineral supplements; Surveys; Risks; Cardiovascular diseases; Elderly

Abstract: Supplemental zinc intake by an ambulatory elderly population was examined. Health questionnaire, electrocardiogram, laboratory, and medication use data from a geriatric health screening program were compared for 69 participants who were taking zinc supplements and 1,832 participants who were not taking zinc supplements during 1985. A major purpose of this study was to evaluate whether ingestion of zinc supplements by elderly persons is a risk factor for development of cardiovascular disease. No association between zinc supplementation and increased risk of developing cardiovascular disease was found for the indices examined. More information is needed to

Nutrition and the Elderly

resolve the question of whether or not zinc supplementation is a risk factor for the development of cardiovascular disease. It is important that elderly persons be provided information to evaluate the possible risk versus benefit of taking zinc supplements.

93 NAL Call No: RM214.T66
Effective nutrition education for older adults.

Fanelli, M.T.
Gaithersburg, Md. : Aspen Publishers; 1988 Oct.
Topics in clinical nutrition v. 3 (4): p. 65-71; 1988 Oct. Includes 17 references.

Language: English

Descriptors: Nutrition education; Elderly; Test construction; Teaching methods

Abstract: The special needs of the older adult learner are discussed. A test to examine the educational needs of the adult is described and instructional methods to reach the older learner are recommended.

94 NAL Call No: TX341.J6

The effectiveness of a television quiz show in providing nutrition information to the elderly.

Goldberg, J.P.; Gershoff, S.N.; McGandy, R.B.; Hartz, S.C.
Baltimore, Md. : Williams & Wilkins; 1989 Apr.
Journal of nutrition education v. 21 (2): p. 86-89; 1989 Apr. Includes 18 references.

Language: English

Descriptors: Nutrition education; Nutrition information; Television; Games; Program evaluation; Elderly

Abstract: This paper describes a television quiz show designed to convey information to the elderly, and provides data on the show's effectiveness.

95 NAL Call No: TX341.C6

Elderly face special nutrition, food problems.

Washington, D.C. : Community Nutrition Institute; 1987 Aug06.

Nutrition week v. 17 (31): p. 4-5; 1987 Aug06.

Language: English

Descriptors: USA; Elderly nutrition; Malnutrition; Socioeconomic status; Nutrient requirements; Health services

Abstract: Excerpts are provided from a position paper and technical support paper issued by the American Dietetic Association on nutrition, aging, and the continuum of health care, noting and em-

phasizing the special food and nutrition problems faced by the US elderly. Special attention is given to nutritional problems of the elderly caused by disease complications, loss of teeth, socioeconomic factors, changes in nutrient requirements for health promotion, and the inadequacy of health care.(wz)

96 NAL Call No: 389.8 B773

Energy, protein, zinc and copper status of twenty-one elderly inpatients: analysed dietary intake and biochemical indices.

Thomas, A.J.; Bunker, V.W.; Hinks, L.J.; Sodha, N.; Mullee, M.A.; Clayton, B.E.

Cambridge : Cambridge University Press; 1988 Mar.

The British journal of nutrition v. 59 (2): p. 181-191. ill., charts; 1988 Mar. Includes 53 references.

Language: English

Descriptors: United Kingdom; Elderly; Protein intake; Energy intake; Zinc; Copper; Nutritional state; Recommended dietary intakes

Abstract: Extract: Duplicate diet analysis for energy, protein, zinc and copper with estimates of biochemical status for Zn and Cu were undertaken in twenty-one elderly long-stay inpatients (mean age 82 (range 63-89) years) consuming their customary hospital diet and in a stable medical condition. Fourteen patients had a long standing and significant healing problem, either a leg ulcer or pressure sore. Mean daily intakes of energy (5.2 MJ), protein (45 g), Zn (85 micromoles) and Cu (14 micromoles) were low in comparison with both official recommendations and levels of intake at which metabolic equilibrium was observed in healthy elderly people studied by the same methods (Bunker et al. 1984a). Mean leucocyte Zn (9 pmol/10(6) cells) and Cu (7.5 pmol/10(6) cells) were low in comparison with results from healthy elderly people (Bunker et al. 1984a), implying sub-optimal status for these elements. Those patients with healing problems tended to have the lower values within the range. Recommendations are made with respect to improving nutritional status in this disadvantaged group of people.(author)

97 NAL Call No: RC620.A1N47

Essential fatty acid metabolism in the elderly: effects of dietary manipulation.

Dillon, J.C.

New York : Raven Press; 1987.

Nesle nutrition workshop series v. 13: p. 93-106. ill., charts; 1987. Papers presented at the Second

Quick Bibliography Series

Nestle Nutrition Research Symposium, Sept 26-27, 1985, Vevey, Switzerland. Includes 47 references.

Language: English

Descriptors: Essential fatty acids; Polyunsaturated fatty acids; Elderly nutrition; Nutrient requirements; Supplements; Nutritional intervention; Enzyme deficiency; Lipid metabolism

Abstract: A technical review summarizes and discusses evidence linking alterations in polyunsaturated fatty acid (PUFA) metabolism, platelet dysfunction, and thrombotic events as they occur in the elderly, and presents clinical and biochemical results on dietary interventions conducted in elderly subjects. Attention is given to: the nature of delta-6-desaturase deficiency in the elderly; age-related disorders in eicosanoid synthesis; and the effect of dietary supplementation with PUFA's in the elderly. The results of a beneficial clinical trial with gamma-linolenate supplementation in a group of 60 elderly patients are included.(wz)

98 **NAL Call No: RC620.A1N8**
Estimation of available dietary iron in meals of "healthy" elderly Japanese.

Itoh, R.; Oka, J.; Echizen, H.

Stoneham, Mass. : Butterworth Publishers; 1988 Jan.

Nutrition reports international v. 37 (1): p. 127-135. ill., charts; 1988 Jan. Includes 13 references.

Language: English

Descriptors: Japan; Elderly; Diet; Nutrient availability; Iron

Abstract: Extract: To establish the amount of absorbable iron necessary to maintain optimum iron nutriture in free living "healthy" elderly Japanese, we tried to estimate the availability of dietary iron in 67 subjects 60 years of age and older, whose iron stores were assessed to be normal. From the value of basal iron loss of 14 micrograms/kg body weight/day, the averages of the iron requirement of the subjects were calculated to be about 0.8 mg/day for males and 0.7 mg/day for females. From geometric means and 95% confidence intervals of serum ferritin concentration, iron stores of most of the male and female subjects were estimated to be in the range of 400-2500 mg and 300-1000 mg, respectively. Using dietary records of each meal in three successive days and a food composition table, we estimated availability of iron, according to the model proposed by Monsen et al. (Am. J. Clin. Nutr., 31, 134, 1987). Averages of dietary iron intake differed in sexes and residential

areas, and ranged from 10 to 12 mg. When the iron stores of the subjects were assumed to be 500 mg, estimated absorbability of dietary iron was about 8% and its amount was 0.8-1.0 mg/day. When the iron stores were assumed to be about 1000 mg, estimated absorbability of dietary iron was 4-4.5% and its amount was 0.5-0.6 mg/day. (author)

99 **NAL Call No: 389.8 J823**
Ethnic and genetic differences in bone mass: a review with a hereditary vs environmental perspective.

Pollitzer, W.S.; Anderson, J.J.B.

Mysore : Association of Food Scientists and Technologists; 1989 Dec.

Journal of food science and technology v. 50 (6): p. 1244-1259. charts; 1989 Dec. Literature review. Includes 90 references.

Language: English

Descriptors: Bone density; Bone weight; Body composition; Calcium; Ethnicity; Heredity; Dietary factors; Nutrient intake; Physical activity; Ethnic groups; Men; Women

Abstract: Based on a review of the literature, ethnic and genetic factors are significant determinants of bone mass, along with such environmental factors as diet and exercise. Differences in bone density between blacks and whites remain even after adjustment for body mass. Black-white differences in bone mass appear to be related to ethnicity because blacks have not only greater skeletal calcium content, but also greater total body potassium and muscle mass. Genetic studies of twins and parent-offspring pairs reflect strong constitutional associations of both bone mineral content and bone density at commonly measured skeletal sites. At least for females, bone mass accumulation by age 20 y is highly associated with maternal bone mass; up to menopause it is enhanced by child-bearing and lactation; beyond menopause environmental factors seem to dominate. Dietary calcium and physical activity are significant in the control of bone mass. These findings are important for osteoporosis and fractures, especially in elderly people.

100 **NAL Call No: 389.9 C1632**
Evaluation of the dietary adequacy of elderly osteoarthritic individuals treated with acetylsalicylic acid.

Taylor Kennedy, J.; Krondl, M.

Toronto, Ont. : The Association; 1987.

Journal of the Canadian Dietetic Association v. 48

Nutrition and the Elderly

(4): p. 232-238. charts; 1987. Literature review. Includes 56 references.

Language: English (Summary in French)

Descriptors: Acetylsalicylic acid; Osteoporosis; Nutrient-drug interactions; Elderly nutrition; Vitamin A; Ascorbic acid

Abstract: A study of 23 elderly osteoarthritic patients taking acetylsalicylic acid (ASA) assessed the effects of ASA on nutrient intakes relative to 20 healthy controls not taking ASA. The patients had higher intakes of niacin and other nutrients (except Ca) than controls, but vitamin A, Ca, and ascorbic acid intakes were below their RDA's. The clinical implications of these findings are discussed.(wz)

101 NAL Call No: TX341.H82

An evaluation of the nutrient intake of a group of elderly people attending a luncheon club.

Potts, M.C.

London : J. Libbey; 1987 Oct.

Human nutrition : applied nutrition v. 41A (5): p. 352-356. charts; 1987 Oct. Includes 13 references.

Language: English

Descriptors: England; Lunch; Groups; Elderly; Nutrient intake; Energy intake; Consumption patterns

Abstract: A study of the nutrient intakes of 18 elderly people (ages 67-85) who frequent a London luncheon club revealed generally adequate intakes (except for Zn), but with wide variations between individuals, including sub-optimal intakes. The luncheon did not markedly alter mean intakes of energy and other nutrients (except for protein). However, it is possible that the luncheon had a greater contribution to daily intake for individuals having low intakes of certain nutrients.(wz)

102 NAL Call No: 389.8 F7398

Factors affecting folate bioavailability.

Bailey, L.B.

Chicago, Ill. : Institute of Food Technologists; 1988 Oct.

Food technology v. 42 (10): p. 206-210, 212, 238. charts; 1988 Oct. Literature review. Includes references.

Language: English

Descriptors: Folic acid; Bioavailability; Intestinal absorption; Ph; Food fiber; Wheat bran; Milk protein; Thermal processes; Nutrient-drug interactions; Aging

103

NAL Call No: 389.8 AM34

Factors affecting food intake of women with Alzheimer's type dementia in long-term care.

Suski, N.S.; Nielsen, C.C.

Chicago, Ill. : The Association; 1989 Dec.

Journal of the American Dietetic Association v. 89 (12): p. 1770-1773. charts; 1989 Dec. Includes 20 references.

Language: English

Descriptors: Nutrient intake; Alzheimer's disease; Eating habits; Supplements; Food preferences; Elderly

Abstract: This study assessed the nutrient intake and problems associated with the feeding of individuals who have advanced senile dementia of the Alzheimer's type (SDAT). The sample consisted of 19 non-ambulatory women, ranging in age from 67 to 105 years, with documented SDAT, who reside in a long-term-care facility. Dietary intake was recorded, and feeding practices were observed for a 3-day period. The mean daily energy intake was 1,558 kcal, with a mean protein intake of 65 gm during the study. Nutritional supplements, consisting of fortified beverages and/or puddings, provided 29% of the calories and 41.5% of the protein intake while contributing considerable amounts of vitamins and minerals. Maintenance of optimal caloric and protein intake is difficult for advanced SDAT patients who are at high risk for infection and skin breakdown related to poor swallowing ability, incontinence, and immobility. Four major factors emerged that promoted optimal intake: using skillful feeding techniques, selecting appropriate food consistency, providing adequate time in which to feed, and capitalizing on the midday meal when cognitive abilities were at their peak.

104

NAL Call No: 389.8 AM34

Factors that influence the elderly to use traditional or nontraditional nutrition information sources.

Probart, C.K.; Davis, L.G.; Hibbard, J.H.; Kime, R.E.

Chicago, Ill. : The Association; 1989 Dec.

Journal of the American Dietetic Association v. 89 (12): p. 1758-1762. charts; 1989 Dec. Includes 30 references.

Language: English

Descriptors: Oregon; Nutrition education; Nutritional surveys; Health services; Utilization; Information sources; Food misinformation; Sex

Quick Bibliography Series

differences; Socioeconomic status; Supplements; Elderly

Abstract: This study analyzed sociodemographic variables, access to medical care, health status-related variables, and nutrition status-related variables to determine their ability to predict an orientation toward traditional or nontraditional nutrition information sources among aged subjects in Eugene, OR. Data were collected from 165 randomly selected subjects by trained interviewers using face-to-face interviews. Nutritionist orientation was measured by a Traditional/Nontraditional Index (TNT Index) developed for the study. Nutrition knowledge and misinformation, supplement use, and sources of nutrition information were also investigated. Predictors of nontraditional nutritionist orientation were acceptance of nutrition misinformation and male sex (adjusted $R^2 = .09$). For women, predictors of a similar orientation were acceptance of nutrition misinformation, low knowledge of general nutrition information, and low functional health status (adjusted $R^2 = .19$). Mean nutrition knowledge scores were high, with a mean of 4.7 (standard deviation [SD] = 0.8) on a scale ranging from 4 (high) to 8 (low). However, many of the subjects did not list dairy foods (28%) and grains (27%) as necessary in the diet. In addition, belief in certain nutrition myths was widespread. More than 58% were taking one or more supplements (mean = 1.4 supplements per day, SD = 1.98). Physicians were mentioned as most likely sources of nutrition information, with dietitians and health food store personnel mentioned second, indicating the importance of marketing efforts for registered dietitians.

105 . NAL Call No: QP141.A1N88
Failure of dietary zinc supplementation to improve the antibody response to influenza vaccine. Bracker, M.D.; Hollingsworth, J.W.; Saltman, P.D.; Strause, L.G.; Klauber, M.R.; Lugo, N.J. Elmsford, N.Y. : Pergamon Press; 1988 Jan. Nutrition research v. 8 (1): p. 99-104. charts; 1988 Jan. Includes 15 references.

Language: English

Descriptors: Influenzavirus; Vaccination; Zinc; Mineral supplements; Antibodies

Abstract: Extract: The effect of administering zinc gluconate on the immune response of elderly subjects to influenza vaccine was studied. Sixty health men and women aged 64 to 90 years (mean = 74) were randomly assigned to two groups and asked

to discontinue all mineral and vitamin supplements for a period of one month. Members of each group were then started on either 50 mg of elemental zinc as zinc gluconate or an identical appearing and tasting placebo in a double blinded manner for one month. Blood samples were then obtained to determine serum zinc and copper levels, and antibody titers to A/Philippines, A/Chile, B/USSR, B/Ann Arbor strains of influenza virus. All individuals were then immunized using trivalent influenza vaccine (Fluogen, Park-Davis), and continued on either zinc or placebo for one month. A second blood sample was then obtained and similarly analyzed. Serum zinc levels increased in both groups, however in the group receiving zinc gluconate supplementation there was a 5.8-fold greater increase than in the control group (p less than 0.001). There was no significant difference in the immune response in the zinc treated group as compared to the placebo group. This study suggests that elevating serum zinc levels in healthy elderly subjects does not enhance the immune response to influenza vaccine.(author)

106 NAL Call No: 389.8 J824
Fasting plasma amino acids in elderly men. Rudman, D.; Mattson, D.E.; Feller, A.G.; Cotter, R.; Johnson, F.C. Baltimore, Md. : American Society for Clinical Nutrition; 1989 Mar. American journal of clinical nutrition v. 49 (3): p. 559-566. charts; 1989 Mar. Includes 56 references.

Language: English

Descriptors: Amino acids; Blood plasma; Fasting; Nutritional state; Elderly; Men

Abstract: The fasting amino acid profile in 22 healthy young men aged 25-35 y (group A) was compared with the fasting profile in 21 healthy independent men aged 65-85 y (group B), in 23 orally-fed nursing home men with dementia aged 65-92 y (group C), and in 17 tube-fed nursing home men with dementia aged 65-88 y (group D). Groups B, C, and D had significantly (p less than 0.05) lower levels of methionine and branched-chain amino acids than group A. Methionine was significantly lower in groups C and D than in group B. The ratio of essential to nonessential amino acids was significantly lower in groups B, C, and D than in group A. The data suggest that the intake of essential amino acids may often be suboptimal in both independent and institutionalized elderly men.

Nutrition and the Elderly

107 NAL Call No: 389.8 J824
Fatty acid composition of serum lipids and its relation to diet in an elderly institutionalized population.

Asciutti-Moura, L.S.; Guillard, J.C.; Fuchs, F.; Richard, D.; Klepping, J.
Baltimore, Md. : American Society for Clinical Nutrition; 1988 Oct.
American journal of clinical nutrition v. 48 (4): p. 980-991. charts; 1988 Oct. Includes 14 references.

Language: English

Descriptors: Fatty acids; Blood serum; Nutrient intake; Source fat; Acyl-coa desaturase; Elderly; Nursing homes; Nutritional state

Abstract: This study was conducted to determine the nutritional status with respect to fatty acids of 53 institutionalized elderly subjects (group A). Seven-day food records and biochemical determinations were used. The same protocol was applied to 25 healthy young adults (group B). The 1981 French Recommended Dietary Allowances were used to assess adequacy of intake. Total fat intake was found to be too high in both groups. Higher intakes of animal fats and saturated fatty acids and lower intakes of polyunsaturated fatty acids and linoleic acid were documented in group A than in group B. Elderly subjects exhibited decreased 18:2-omega-6 and 20:4-omega-6 in serum phospholipids and triglycerides. Other indicators of essential fatty acid status in group A differed from group B in the direction of deficiency (double-bond index [DBI], 0.93 +/- 0.01 vs 1.13 +/- 0.04, p less than 0.001; total omega-6, 25.07 +/- 0.46% vs 32.49 +/- 0.89%, p less than 0.001). These findings indicate that institutionalized elderly people are deficient in circulating essential fatty acids, which appears to be caused by both diet and metabolism.

108 NAL Call No: TX361.A3J63
Feeding the dysphagic elderly resident: a case study.
Matthews, L.E.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 8 (4): p. 59-64; 1988. Includes 1 references.

Language: English

Descriptors: Individual feeding; Long term care; Clinical nutrition; Menus; Case studies; Elderly

Abstract: A brief case study of a nursing home resident with chewing and swallowing problems is presented. The nutritional assessment, the

nutritional plan and the daily menu plan for the patient is included.

109 NAL Call No: TX341.F69
Focusing elderly nutritional programs.
Leonard, R.
New York, N.Y. : Harcourt Brace Jovanovich Publishers; 1987 Mar.
Food management v. 22 (3): p. 34. ill; 1987 Mar.

Language: English

Descriptors: Elderly nutrition; Supplemental feeding programs; Federal programs; Food and nutrition controversies

Abstract: A brief commentary critiques perceived shortcomings of elderly nutrition programs, arguing that federal and state "aging" program managers give little real attention to elderly nutrition and that a managerial system is needed that can accomplish the goals of the Older American Act and its authorized service programs at a more reasonable cost than currently is being paid. The view is expressed that in the current system, individuals become recipients rather than participants whose needs are determined by their interviewers as they are processed through an intake system. It is suggested that the current system may encourage Congress to ask whether the current program is achieving the goal of aiding older people to maintain independent lifestyles in their communities and how program management may be improved.(wz)

110 NAL Call No: RC455.4.N8N8
Food and fluid intake as criteria of disturbed adaptivity in old age.
Janicke, B.; Cooper, H.
Basel : Karger; 1987.
Nutrients and brain function / editor, W.B. Esman. p. 103-115. ill; 1987. Includes 17 references.

Language: English

Descriptors: Adaptation; Nutrient requirements; Aging; Food intake; Fluid intake

Abstract: A technical report discusses and provides experimental animal data concerning the association between food and fluid intake and disturbed adaptivity in old age. The data, addressing various physiological mechanisms of adaptive behavior, show that latent functional losses of adaptability in the 3rd phase of life can be detected. The study approach is based on subjecting aging animals to physical and drug challenges.

Quick Bibliography Series

111 NAL Call No: TX361.A3J6
Food and nutrient intake of institutionalized and noninstitutionalized elderly in the Philippines.
Florencio, C.A.; Macatangay, L.A.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (4): p. 17-30. charts; 1987. Includes references.

Language: English

Descriptors: Philippines; Elderly nutrition; Nutrient intake; Energy intake; Food habits; Hospitals; Nutritional adequacy; Dietary surveys; Recommended dietary allowances; Institutional food service

Abstract: Extract: The study was conducted to compare the food and nutrient intake of institutionalized and noninstitutionalized elderly from the same geographical area. Data were obtained using food weighing-recall method and interview. Sixty elderly in the same section of Quezon City, Philippines participated in the study: thirty from a government institution for the elderly and thirty from among those living with their families. Both groups failed to meet the Philippine Recommended Daily Allowances for energy and the eight nutrients considered. However, the average diet of the noninstitutionalized elderly was more nutritionally adequate than that of the institutionalized. (author)

112 NAL Call No: TX341 C6
Food contamination: widespread, increasing.
Washington, D.C. : Community Nutrition Institute; 1987 Apr02.
Nutrition week v. 17 (14): p. 4-5. ill; 1987 Apr02.

Language: English

Descriptors: USA; Foodborne diseases; Salmonella; Campylobacter; Trends; Costs; National surveys

Abstract: A special report provides data illustrating the increase in foodborne illnesses in the US, up to half of which resulted from contaminated meat and poultry, particularly from Salmonella and Campylobacter. These 2 bacteria are separately discussed with respect to their contamination levels and health impacts. Infants, elderly, malnourished, and people with an immune deficiency are most susceptible. Human and economic costs associated with such illnesses are discussed.(mjs)

113 NAL Call No: TX361.A3J6
Food preferences and perceptions of institution-

alized elderly: a case report.

Pierce, R.P. Jr; Hodges, P.A.M.; Merz, B.; Olivey, A.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 7 (1): p. 35-42. charts; 1987. Includes 6 references.

Language: English

Descriptors: Elderly; Institutions; Food preferences; Environmental factors; Case studies
Abstract: Extract: Providing nutritious, appetizing meals to institutionalized elderly is a continuing challenge. Menu planning must reflect sensitivity to regional, cultural, and ethnic food preferences, in addition to nutritional needs of the population to be served. In order to raise the level of awareness of dietetic technician students--future menu planners--regarding the eating habits and food preferences of the elderly, this project was undertaken in Tidewater, Virginia. Two facilities were used to interview selected institutionalized elderly patients. Results are reported, with considerations for future menu planning, as well as implications of the impact of early eating habits upon preferences in later years.(author)

114 NAL Call No: TX361.A3J6
Food preferences of older adults.
Holt, V.; Nordstrom, J.; Kohrs, M.B.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (3): p. 47-55. ill; 1987. Includes 20 references.

Language: English

Descriptors: Missouri; Food preferences; Elderly; Food habits; Regional surveys

Abstract: Extract: Food preferences of 488 elderly persons over 59 years old were determined. Beef, potatoes, chicken, green beans, pie, pork, dried beans, lettuce salad, cake and fish were given most often as liked. Nine of the top ten food items given as disliked were vegetables. Items from the basic four food groups were given as "good for you" the most frequently.(author)

115 NAL Call No: 389.8 AM34
Food stamps and the energy status of the U.S. elderly poor.
Lopez, L.M.; Habicht, J.P.
Chicago, Ill. : The Association; 1987 Aug.
Journal of the American Dietetic Association v. 87 (8): p. 1020-1024. charts; 1987 Aug. Includes 34 references.

Nutrition and the Elderly

Language: English

Descriptors: USA; Elderly; Low income groups; Food stamp program; Participation; Energy consumption; National surveys

Abstract: Extract: The energy status of poor elderly individuals in relation to the nonpoor elderly was examined using data from NHANES I and II. Within the poor population, the status of Food Stamp Program participants was compared with that of nonparticipants. The indicators of energy status used were weight (adjusted for height), triceps skinfold, subscapular skinfold, and energy intake. Energy status was examined by analyses of variance; the effects associated with sex, race, age, geographic region, and other potentially confounding variables were included in the model. In both surveys, there were interactions of income and food stamp use with sex. That is, poor women had greater means for weight and triceps skinfold than the nonpoor. Among the poor women, the Food Stamp Program participants weighed more than the nonparticipants, and their triceps skinfold mean was greater in NHANES II than in NHANES I. The poor men had a lower mean for triceps skinfold than the nonpoor, but there was no difference between the program participants and nonparticipants. Hence, for the poor elderly in NHANES I and II, food stamp use was not consistently associated with differences in energy status, as indicated by weight, skinfold, and energy intake.(author)

116 NAL Call No: 389.8 AM34
Food stamps and the iron status of the U.S. elderly poor.
Lopez, L.M.; Habicht, J.P.
Chicago, Ill. : The Association; 1987 May.
Journal of the American Dietetic Association v. 87 (5): p. 598-603. charts; 1987 May. Includes 38 references.

Language: English

Descriptors: USA; Iron; Nutritional state; Elderly; Food stamp program; Low income groups; Elderly nutrition; National surveys

Abstract: Extract: With the use of data from NHANES I and II, the iron status of poor elderly persons was examined by comparing it with that of nonpoor elderly persons. Within the poor group, the iron status of Food Stamp Program participants was compared with that of nonparticipants. The poor were those below the poverty line; the nonpoor were those with incomes at least twice the poverty line. The indicators of iron status were:

hemoglobin, hematocrit, free erythrocyte protoporphyrin, transferrin saturation, total iron-binding capacity, serum iron, and iron intake. Iron status was examined by analyses of variance. The effects associated with sex, race, age, geographic region, and other potentially confounding factors were included in the model. In both surveys, there were interactions with race. That is, the poor whites had lower transferrin saturation and iron intake means than the nonpoor whites, while the Food Stamp Program participants were no different from the nonparticipants in transferrin saturation. Within the nonwhites, the poor had a lower hemoglobin mean than the nonpoor, and the program participants had a lower hemoglobin mean than the nonparticipants. Therefore, the Food Stamp Program was not consistently associated with better iron nutrition of the elderly poor in NHANES I and II.(author)

117 NAL Call No: RC620.A1J6
Fractures in the men of a Veterans Administration Nursing Home: Relation of 1,25-dihydroxyvitamin D.
Rudman, D.; Rudman, I.W.; Mattson, D.E.; Nagraj, H.S.; Caindec, N.; Jackson, D.L.
New York, N.Y. : John Wiley & Sons; 1989 Aug.
Journal of the American College of Nutrition v. 8 (4): p. 324-334. charts; 1989 Aug. Includes 42 references.

Language: English

Descriptors: 1,25-dihydroxycholecalciferol; Bone fractures; Blood chemistry; Diseases; Drug therapy; Nursing homes; Elderly; Men
Abstract: The authors studied the incidence of bone fracture in middle-aged and elderly men living in nursing homes. Health status, types of medication, age, blood chemistries and vitamin D levels were measured. The emphasis of the study was to correlate vitamin D deficiency with increased bone fractures.

118 NAL Call No: 389.8 J82
The future direction of nutrition research: unraveling the diet-disease connection.
Feldman, E.B.
Bethesda, Md. : American Institute of Nutrition; 1989 Feb.
The Journal of nutrition v. 119 (2): p. 323-324; 1989 Feb. Includes 3 references.

Language: English

Descriptors: Nutrition research; Trends, Health

Quick Bibliography Series

promotion; Disease prevention; Maternal nutrition; Fetus; Cardiovascular diseases; Aging; Behavior; Nutrition policy

Abstract: This opinion statement proposes that the future of nutrition research is in the direction of diet and disease relationships. Specific areas mentioned are fetal nutrition, heart disease, aging and longevity, diet and behavior, and nutrition and public policy.

119 NAL Call No: QP141.A1N88
Genetic influences on the selection of brassica vegetables by elderly individuals.
Niewind, A.; Kronld, M.; Shrott, M.
Elmsford, N.Y. : Pergamon Press; 1988 Jan.
Nutrition research v. 8 (1): p. 13-20. charts; 1988 Jan. Includes 33 references.

Language: English

Descriptors: Elderly nutrition; Brassica; Vegetables; Consumption patterns, Genetic factors; Taste

Abstract: Extract: The relative influence of a genetic trait, the ability to taste phenylthiocarbamide (PTC), on rating bitterness of cabbage and liking it, and on the use of Brassica vegetables was examined among 32 subjectively healthy women, 53 to 76 years of age. Sensitivity to PTC was assessed with a forced choice staircase procedure. Flavour intensity and hedonic ratings of cabbage in raw and cooked form were obtained by quantitative descriptive analysis techniques. Frequency of use of Brassica vegetables were recorded. Clear sensory difference in PTC sensitivity categorized the sample into 17 PTC tasters and 15 non-tasters, but it did not map on the suprathreshold bitter sensitivity, preference or frequency of use of Brassica vegetables, although the PTC tasters rated the overall flavour intensity of cooked cabbage higher than did the non-tasters. The results indicate that other than genetic factors may impede the use of a variety of Brassica vegetables by old women.(author)

120 NAL Call No: TX341.C6
Genetic variation and nutrition: highlights.
Washington, D.C. : Community Nutrition Institute; 1989 Jul13.
Nutrition week v. 19 (27): p. 4-5; 1989 Jul13.

Language: English

Descriptors: Nutrition; Genetic variation; Genotype nutrition interaction; Food preferences; Taste threshold, Food allergies

Abstract: An individual phenotype consists of interactions of genotype, environment and institutional influences. Food preferences, nutritional thresholds adverse reactions to foods, and the aging process may be determined by genetic factors. The effect of genetic factors on the intake of essential or nonessential nutrients is discussed. Knowledge of genetic individuality may help in disease prevention when the environment in which the particular gene is likely to act is taken into account.

121 NAL Call No: TX361.A3R63 1987
Geriatric nutrition.
Roe, Daphne A.,1923-
Englewood Cliffs, N.J. : Prentice-Hall; 1987.
xi, 267 p. : ill. ; 24 cm. Includes bibliographies and index.

Language: English

Descriptors: Aged--Nutrition; Geriatric Reference works, Elderly nutrition; Patient care Nutritional requirements; Therapeutic diets; Nutritional assessment

Abstract: A basic text for senior undergraduate or graduate nutrition majors and a reference text for health professionals treating elderly patients provides scientific and practical information and guidelines on the nutritional needs and care of the elderly. The 11 text chapters cover: life style characteristics of the elderly, the physiology and pathology of aging (including the aging processes of tissues and organs and the characteristics of a variety of diseases); nutrition surveys on the nutritional status of the elderly; specific nutritional requirements; factors influencing food intake; procedures for assessing nutritional status; nutritional deficiencies; diseases that respond to dietary modifications; the effect of drugs and drug-nutrient interactions in the elderly; nutrition services for the elderly; and elderly nutrition in the third world with respect to common elderly diseases (hypertension, coronary artery disease, diabetes, musculoskeletal diseases). A set of appendices includes class projects, essay questions, a listings of resource materials, and a glossary of medical terms.

122 NAL Call No: TX361.A3J63
Geriatric nutrition in Canada--a review.
Yeung, D.L.; Imbach, A.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 7 (3): p. 27-45. charts; 1988. Literature review. Includes 30 references.

Nutrition and the Elderly

Language: English

Descriptors: Canada; Elderly nutrition; Nutrient requirements; Recommended dietary intakes; Nutrient intake; Caloric intake; Food preferences; Elderly

Abstract: Since 1965, a small number of studies on geriatric nutrition have been conducted in various parts of Canada. This paper summarizes the major observations of these studies and recommends areas for future research.

123 NAL Call No: HD9000.9.U5A1
The 'grazing' of America: a guide to healthy snacking.
Weinstock, C.P.
Rockville, Md. : Food and Drug Administration, Department of Health & Human Services; 1989 Mar.
F.D.A. consumer v. 23 (2): p. 8-13. charts; 1989 Mar.

Language: English

Descriptors: USA; Snacks; Eating patterns; Feeding habits; Nutrient intake; Low calorie foods; Children; Adolescents; Elderly

Abstract: Since many people do not eat three regular meals a day, snacking has become a part of the nation's eating pattern. Healthy snacks for different stages of life are discussed. Nutrient and caloric intake charts are included.

124 NAL Call No: TX361.A3J6
Health factors associated with food intolerances among meals-on-wheels clientele.
Zimmerman, S.; Kronld, M.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 7 (1): p. 23-33. charts; 1987. Includes 16 references.

Language: English

Descriptors: Canada; Elderly; Home delivered meals; Vegetables; Food intolerance; Incidence; Health; Drugs; Supplements

Abstract: Extract: A study was carried out with Meals-on-Wheels clientele to determine differences between tolerators and intolerators of vegetables in regard to health status, eating problems, medication and supplement use. Sixty-six subjects were divided into two groups on the basis of perceived tolerance to twenty different vegetables. Significantly more intolerators had digestive disorders and took more medications. Supplement use was more uniform in the two

groups. There were no significant differences in subjective health ratings or in chewing and swallowing problems. Identification of meal recipients on the basis of food intolerances would alert the meal providers to persons who need special care.(author)

125 NAL Call No: 389.9 C1632
Health-related characteristics and dietary intake of a selected sample of elderly Canadians.

Martinez, O.B.
Toronto, Ont. : The Association; 1989.
Journal of the Canadian Dietetic Association v. 50 (1): p. 36-40. charts; 1989. Includes 27 references.

Language: English

Descriptors: Ontario--fre; Food intake; Interviews; Questionnaires; Computer analysis; Nutrient intake; Caloric intake; Vitamin And mineral supplements; Physical activity; Health; Drug therapy; Sex differences; Elderly

Abstract: The following study describes selected health characteristics and the dietary intakes of 78 free-living volunteers, 59-88 years of age from Southern Ontario. Data on health, food supplement and medication use and other information were obtained during home interviews using a questionnaire. Dietary data were obtained through three 24-hour recalls. The information was coded and analyzed by computer. Seventy-one percent of the women (n=56) and 32% of the men (n=22) reported at least one chronic health problem, especially arthritis/rheumatism, hypertension, gastrointestinal problems and heart disease. Most subjects (79%) reported habitual use of at least one drug but 27% of the women used four or more drugs simultaneously. Reported compliance with medical treatment was high in both sexes. Twenty percent of the women and 4.5% of the men had lower than recommended energy intakes. In women, 21 and 16%, respectively had lower than recommended intakes of calcium and vitamin D. Men's diets were generally adequate. Five to 23% of the men and 29-45% of the women reported the use of individual nutrient supplements. In conclusion these subjects showed generally adequate nutrient intakes in spite of low energy intakes; calcium and vitamin D were the nutrients more likely to be present in lower than recommended levels in the diet.

126 NAL Call No: aTX341.F615
Helping young and old.
Rudman, L.

Quick Bibliography Series

Washington, D.C. : U.S. Department of Agriculture; 1987 Apr.
Food & nutrition v. 17 (2): p. 22-23. ill; 1987 Apr.

Language: English

Descriptors: Michigan; Nutrition programs; Supplemental feeding programs; Community action; Mothers; Federal programs

Abstract: An illustrated report describes the activities and effectiveness of a Detroit agency food center, Focus HOPE, for providing USDA-donated foods to 38,000 low-income mothers, infants, and children and 26,000 elderly people. The program includes nutrition education as part of its operation, including a cookbook for mothers with recipes using USDA commodity foods. The operational structure of the food center also is briefly described.(wz)

127 NAL Call No: RM216.M342
Home nutrition support in elderly patients.
Chernoff, R.
St. Louis, Mo. : C. V. Mosby Company; 1987 Jan.
Clinical nutrition v. 6 (1): p. 36-39. charts; 1987 Jan. Includes 9 references.

Language: English

Descriptors: Parenteral feeding; Elderly; Home care; Therapeutic diets

Abstract: Extract: The greatest users of acute health care resources are persons aged 65 and older. Because of the impact of hospital reimbursement systems, elderly patients are going home from the hospital earlier with nutrition support therapies. Elderly persons require special considerations because of the physiologic and functional changes that occur as part of the normal aging process. They require careful monitoring because they are more vulnerable to complications and less likely to respond rapidly to change. They are difficult to monitor by usual parameters because of age-related changes that occur in many of these measures. In addition, there are many other factors that are unique in elderly patients, such as polypharmacy and potential rapid shifts in fluid compartments. Nutritional rehabilitation in frail, older patients can be accomplished by encouraging oral feeding or providing supplemental meals, or by instituting enteral or parenteral nutrition support. There are many factors to consider when assessing elderly patients for these therapies. Social, financial, physical, psychological, and ethical aspects of nutrition support should be explored during

decision-making. With the increasing number of persons living beyond age 65, home nutrition support and the factors affecting its successful use in elderly persons will become more important.(author)

128 NAL Call No: 389.8 AM34
Home-delivered meals: food quality, nutrient content, and characteristics of recipients.

Asp, E.H.; Darling, M.E.
Chicago, Ill. : The Association; 1988 Jan.
Journal of the American Dietetic Association v. 88 (1): p. 55-59. charts; 1988 Jan. Includes 25 references.

Language: English

Descriptors: Home delivered meals; Food quality; Nutrient intake; Elderly nutrition; Nutritional adequacy; Program evaluation

Abstract: Extract: Meals provided by a home-delivered meal program with the same supplier were evaluated for 5 consecutive days in each of 6 years. In the evaluations, meals were compared with federal guidelines for meal pattern, serving size, and temperature of hot food, and the quality was evaluated. Meal pattern and serving size guidelines were not always met by the meals. Problems included no delivery of milk with the meals and variations in serving sizes for meat or alternate, fruit, vegetable, and dessert. Temperatures of hot foods were often much lower than the 140 degrees or 150 degrees F specified in food safety guidelines. Food quality varied. Protein, iron, phosphorus, vitamin A, and niacin were consistently greater than 33% of the RDA; energy, thiamin, riboflavin, and vitamin C were less than 33% of the RDA some years, and calcium was consistently less than 33% of the RDA. Estimated total daily nutrient intakes of 27 recipients followed the same pattern as the nutrient content of the meals. For many recipients, estimated nutrient intakes from food other than the home-delivered meal was less than 33% of the RDA. Food preferences were fairly well satisfied. Recall of nutrition-related vocabulary was poor. In this study, home-delivered meals were found to make important contributions to the nutrition needs of the recipients.(author)

129 NAL Call No: 389.8 F7398
How the aging explosion will create new food trends.
Gerber, J.
Chicago, Ill. : Institute of Food Technologists; 1989 Apr.

Nutrition and the Elderly

Food technology v. 43 (4) p. 134-135. 150; 1989 Apr. Paper presented at the "Symposium on Changing Trends in Diet, Nutrition, and Consumer Taste: Implications for Food Technologists," June 19-22, 1988, New Orleans, Louisiana.

Language: English

Descriptors: Food industry; Trends; Nutrition labeling; Ethnic foods; Life style; Elderly

Abstract: Food industry innovations which satisfy the needs of the older consumer may also motivate the buying trends of the younger market. Examples given in this article include improved labeling and packaging of food and foods designed for health promotion.

130 NAL Call No: 389.8 AM34
Hypophagia among hospitalized elderly.

Rammohan, M.; Juan, D.; Jung, D.
Chicago, Ill. : The Association; 1989 Dec.
Journal of the American Dietetic Association v. 89 (12): p. 1774-1779. charts; 1989 Dec. Includes 38 references.

Language: English

Descriptors: Nutrient intake; Age differences; Sex differences; Body weight; Eating habits; Appetite; Elderly

Abstract: A prospective study was done in 21 hospitalized medical patients to determine the effects of age and gender on food intake. From precise weighing of food served and uneaten, macronutrients and micronutrients ingested during the hospital stay were determined. Even though the hospital diet served was adequate in energy and protein, 38% of the patients ingested less than 65% of their nutritional requirements, with a higher percentage of those with inadequate intake among the elderly than among the young (28% vs. 10%). Patients over 65 years of age ingested less energy and protein than those under 65 year of age (1,167 vs. 1,967 kcal, p less than .01; 47 vs. 78 gm, p less than .05). The gender of the patients had no influence on food intake. More than 60% of the medical patients, especially the elderly, had an inadequate intake of micronutrients: folate, 100% of the elderly vs. 93% of the young; zinc, 90% vs. 64%; magnesium, 90% vs. 36%; and vitamin B-6, 90% vs. 64%. Multiple stepwise regression analysis showed that age and body weight on admission were of predictive value in terms of subsequent nutrient intakes during the hospitalization. Patients who were over 65 years old and weighed less than

80% of their ideal body weight consumed significantly less energy and fewer macronutrients and micronutrients.

131 NAL Call No: QP141.A1N88
Immunodeficiency and lymphocyte ecto-5'-nucleotidase activity in the elderly: a comparison of the effect of a trace mineral supplement (1 USRDA) with high zinc (6.7 X USRDA).
Hollingsworth, J.W.; Otte, R.G.; Boss, G.R.; Fryberger, M.F.; Strause, L.G.; Saltman, P.
Elmsford, N.Y. : Pergamon Press; 1987 Aug.
Nutrition research v. 7 (8): p. 801-811. ill., charts; 1987 Aug. Includes 26 references.

Language: English

Descriptors: Zinc; Immunity; Elderly nutrition; Nucleotidase; Enzyme activity; Mineral supplements

Abstract: Extract: The effect of administering a total mineral supplement (TMS) or 100 mg of elemental zinc alone on immune function was studied in 17 normal subjects, aged 66-85. Subjects were given either TMS containing 1 USRDA of zinc, copper, iron, and manganese or 100 mg of elemental zinc alone as zinc sulfate (ZnSO₄). Administration of either supplement led to a slight increase in peripheral blood lymphocytes and improved maturation of B-lymphocytes to antibody-producing cells in culture of peripheral blood mononuclear cells. Ecto-5'-nucleotidase, a zinc obligate enzyme of lymphocyte cell membranes, increased when TMS was administered daily but failed to increase when ZnSO₄ was administered. Serum zinc levels increased in all subjects receiving the ZnSO₄. Only those subjects with low initial serum zinc levels had higher serum zinc levels after 3 months on the TMS. These studies indicate that dietary supplementation with a TMS (1 USRDA) can improve certain in vitro parameters of immune competence in the aged, avoiding the deleterious effects possible with excessive intake.(author)

132 NAL Call No: TX361.A3J6
The impact of food stamps and other variables on nutrient intake in the elderly.

Posner, B.M.; Ohls, J.C.; Morgan, J.C.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (3): p. 3-16. charts; 1987. Includes 29 references.

Language: English

Descriptors: USA; Elderly nutrition, Food stamp program; Nutrient intake; Nutritional intervention;

Quick Bibliography Series

Supplemental feeding programs; Federal programs; Program effectiveness; National surveys

Abstract: Extract: The effect of the national Food Stamp program and other variables on nutrient intake and dietary adequacy in low-income elderly persons was determined by comparing the dietary behavior of 1400 program participants and 500 eligible non-participants. Data were gathered using a validated telephone-administered 24-hour dietary recall and household interview. Food Stamps did not produce statistically significant improvements in nutrient intake in these households. Functional health status, gender, race, nutrition knowledge, social contact and perceived problems with food preparation were variables found to predict differences in nutrient intake among the low-income elderly. An indirect income transfer program like Food Stamps appears at best to have limited effectiveness as a method of improving the nutrient intake of low-income households headed by older persons. (author)

133 NAL Call No: 389.8 AM34

The impact of the diagnosis-related groups/prospective payment system on nutrition needs in home health and extended-care facilities. Sandall, M.J.; Massey, L.K.

Chicago, Ill. : The Association; 1989 Oct.
Journal of the American Dietetic Association v. 89 (10): p. 1444-1447. charts; 1989 Oct. Includes 17 references.

Language: English

Descriptors: Washington; Health services; Health care costs; Diagnosis; Elderly nutrition; Health care; Home care; Hospitals; Nutritional assessment; Nursing homes; Elderly

Abstract: The effect of the diagnosis-related groups (DRG)/prospective payment system (PPS) on the health status complexity and nutrition needs of patients was studied in home health and five extended-care facilities in Spokane, WA. Three periods were studied: pre-DRG (1982), early DRG implementation (1984), and full DRG implementation (1986). The complexity of patient health status and nutrition needs was measured in three areas: number of total diagnoses, nutrition-related diagnoses, and diet modifications per patient. In home health care, those variables increased significantly from a mean of 4.35 diagnoses per patient in 1982 to 6.12 in 1986, from 1.31 nutrition-related diagnoses per patient in 1982 to 1.99 in 1986, and from 6% of patients with four or more diet modifications in 1982 to 16% in 1986. The complexity

of patient health status in extended-care facilities remained stable, with a mean of 6.06 diagnoses per patient in 1982 vs. 5.87 in 1986, 1.84 nutrition-related diagnoses per patient vs. 1.73, and 14% of patients having two or more diet modifications in 1982 vs. 11% in 1986. A consistent odds ratio of 2.06 was observed between number of diagnoses and diet modifications per patient, indicating a direct relationship.

134 NAL Call No: RM214.D47

Impairment of taste and smell in the elderly.

Schiffman, S.S.; Warwick, Z.S.

Rockville, Md. : Aspen; 1988 Jan.

Directions in applied nutrition v. 2 (3): p. 1, 4-6. charts; 1988 Jan. Includes 4 references.

Language: English

Descriptors: Smell; Taste; Food intake; Flavor enhancers

Abstract: The effects of taste and smell impairment on decreased food intake of the elderly are discussed. Also discussed are the physiology and treatment by flavor enhancement of chemosensory loss. Conclusions are that addition of palatable flavors to nutrient-dense foods for elderly persons with reduced taste and smell perception may increase consumption.

135 NAL Call No: 389.8 AM34

Implications of malnutrition and diagnosis-related groups (DRGs).

Delhey, D.M.; Anderson, E.J.; Laramée, S.H.

Chicago, Ill. : The Association; 1989 Oct.

Journal of the American Dietetic Association v. 89 (10): p. 1448-1451. charts; 1989 Oct. Includes 20 references.

Language: English

Descriptors: Health care costs; Malnutrition; Elderly nutrition; Diagnosis; Hospitals; Nutritional assessment; Elderly

Abstract: The implications of malnutrition on the Medicare Prospective Payment System of diagnosis-related groups (DRGs) were examined in 185 Medicare patients, aged 65 to 69 years, admitted to an acute-care tertiary hospital. Upon admission, patients were classified as malnourished if they were below the acceptable level in at least two of four parameters: serum albumin concentration, total lymphocyte count, percent ideal body weight, and percent weight loss. On the basis of that criterion, 8.6% (no.=16) of the 185 patients were classified as malnourished. Although patients

Nutrition and the Elderly

were classified as malnourished, malnutrition was among several comorbidity and complication (CC) factors, and, therefore, coding for malnutrition in any of these cases would not have increased hospital reimbursement. If malnutrition had been the only CC factor, hospital reimbursement would have been enhanced.

136 NAL Call No: 389.8 AM34
Inclusion of geriatric nutrition in ADA-approved undergraduate programs.

Shoaf, L.R.; Jensen, H.M.

Chicago, Ill. : The Association; 1989 Sep.

Journal of the American Dietetic Association v. 89 (9): p. 1282-1285. charts; 1989 Sep. Includes 22 references.

Language: English

Descriptors: Dietetic education; Curriculum, Elderly nutrition; Educational programs

Abstract: All ADA Plan IV programs were surveyed to determine whether geriatric nutrition was included in their curriculums. Of the 268 Plan IV programs, 66% responded. Less than one-fifth of the programs offered or planned to offer a specific geriatric nutrition course. An overview of geriatric nutrition occurred most frequently in a human nutrition course. A practicum/clinical experience or a course other than nutrition most frequently provided in-depth study, if such was available. Nursing homes and congregate meal sites were the primary locations for experiences with the geriatric population. Major activities with that age group included (a) taking diet histories, (b) making nutrition assessments, and (c) providing diet instruction. In some programs, didactic and experimental training with the geriatric population may not be adequate to prepare dietetic undergraduate students to meet the health care needs of that growing segment of society.

137 NAL Call No: QP141.A1N88
Indices of vitamin, iron and hematological status of a selected sample of elderly Canadians.

Martinez, O.B.

Elmsford, N.Y. : Pergamon Press; 1988 Dec.

Nutrition research v. 8 (12): p. 1345-1351. charts; 1988 Dec. Includes 23 references.

Language: English

Descriptors: Ontario; Ascorbic acid; Folic acid; Riboflavin; Vitamin B12; Ferritin; Blood plasma; Elderly; Nutritional state

Abstract: This study investigated vitamin, iron and

hematological indices of 52 women and 22 men volunteers, 59-88 years of age from Southern Ontario. Background information was obtained in home interviews. Fasting blood samples were obtained for measurements of plasma ascorbate, ferritin and vitamin B12 levels, and for red cell folate and red cell glutathione reductase activity (GRA) coefficients. The information was coded and analyzed by computer. Using hemoglobin values as criteria, there was no evidence of anemia in this sample. Ascorbate, folate and iron levels were adequate. The results suggest that subclinical nutritional deficiencies of riboflavin and vitamin B12 may occur in some elderly and that supplementation with the respective vitamins may be beneficial

138 NAL Call No: 389.8 B773

The influence of guar-gum bread on the regulation of diabetes mellitus type II in elderly patients.

Sels, J.P.; Flendrig, J.A.; Postmes, T.J.

Cambridge : Cambridge University Press; 1987 Mar.

The British journal of nutrition v. 57 (2): p. 177-183. ill., charts; 1987 Mar. Includes 33 references.

Language: English

Descriptors: Netherlands; Man; Elderly; Diet; Diabetes; Bread; Fiber; Guar gum; Hemoglobin a1; Blood serum; Glucose; High density lipoprotein; Cholesterol; C-peptide; Triglycerides

Abstract: Extract: The effect of a high-fibre bread mixed with guar-gum (75 g/kg flour) on serum glucose, connecting peptide (C-peptide), haemoglobin A1 (HbA1), high-density-lipoprotein-cholesterol (HDL-cholesterol) and triglycerides was examined in fourteen elderly patients with diabetes mellitus type II. The mean daily consumption of guar gum was 8.1 g. Gastrointestinal disorders were not observed. Consumption of guar-gum bread resulted in a significant decrease in C-peptide values on the 1st day and blood glucose values after 3 weeks (both measured 90 min after breakfast). C-peptide values remained low, while an unaccountable 'rebound' phenomenon was seen in the blood glucose values 90 min after breakfast at the end of the study.(author)

139 NAL Call No: QP141.A1A64

Interindividual and intraindividual variation in repeated measures of 24-hour dietary recall in the elderly.

McAvay, G.; Rodin, J.

London : Academic Press; 1988 Dec.

Appetite v. 2 (2): p. 97-110. charts; 1988 Dec. In-

Quick Bibliography Series

cludes 21 references.

Language: English

Descriptors: Dietary history; Nutrient intake; Elderly; Longitudinal studies; Human nutrition research; Variations

Abstract: This study examines the validity and variability of the 24-hour dietary recall method for determination of the nutrient intake of older adults.

140 NAL Call No: QP141.A1N88
Intraindividual variation in zinc intake among elderly women: the effect of diet and supplements. Ervin, R.B.; Smiciklas-Wright, H.A.; Fosmire, G.J. Elmsford, N.Y. : Pergamon Press; 1989 Jun. Nutrition research v. 9 (6): p. 613-624. charts; 1989 Jun. Includes 30 references.

Language: English

Descriptors: Zinc; Nutrient intake; Dietary history; Supplements; Data analysis; Variations; Elderly; Women

Abstract: Thirty-six women (65-84 y) kept 3-day dietary records and recorded supplement use at three periods in time, each 28 days apart. Sources of intraindividual variation in zinc intake were examined. Mean dietary zinc intake was 7.3 +/- 2.9 mg/d; mean total zinc intake (diet plus supplements) was 11.7 +/- 9.3 mg/d. For both types of zinc intake, there was significant intraindividual variation among the time periods (p less than 0.0001), but not among the three days each subject kept her records. The intraindividual variation for independent records was not significantly different from that for consecutive records. Primary sources of intraindividual variation in zinc intake appeared to be dietary modifications related to acute illnesses, changes in calorie, protein and/or zinc intakes, and variable use of supplements. Intraindividual variance for irregular supplement users was significantly larger (p less than 0.0005) than the intraindividual variances for both non-supplement users and regular supplement users.

141 NAL Call No: 389.1 W892
Iron and folate nutriture during life cycle. Bailey, L.B.; Cerda, J.J. Basel : S. Karger; 1988. World review of nutrition and dietetics v. 56: p. 56-92. charts; 1988. In the series analytic: Aspects of Nutritional Physiology / edited by G.H. Bourne. Literature review. Includes 116 references.

Language: English

Descriptors: Iron; Folates; Life cycles; Pregnancy; Lactation; Adolescence; Elderly; Infants; Nutritional state; Iron deficiency anemia; Recommended dietary allowances

Abstract: Iron and folate are two essential nutrients that are often deficient in diets of population groups. Both are required for the normal functioning of the hematopoietic system as well as numerous critical metabolic processes. A deficiency of either will result in anemia with the associated physiological consequences. Periods of rapid growth and development as during pregnancy, early childhood, and adolescence increase the requirement for both iron and folate. An increase in the prevalence of deficiency is associated with these periods of the life cycle. Changes in iron and folate status in the aged may occur due to a combination of physiological factors such as reduction in nutrient intake coupled with drug and alcohol use. The purpose of this paper is to review the basics of iron and folate nutriture in conjunction with a discussion of how the physiological changes during the life cycle may affect the requirement for these two nutrients.

142 NAL Call No: RC620.A1N8
Iron status in free-living, low income very elderly. Manore, M.M.; Vaughan, L.A.; Carroll, S S. Stoneham, Mass. : Butterworth; 1989 Jan. Nutrition reports international v. 39 (1): p. 1-12. charts; 1989 Jan. Includes 22 references.

Language: English

Descriptors: Nutritional state; Iron; Nutrient intake; Ferritin; Blood serum; Anthropometric dimensions; Supplements; Low income groups; Elderly; Sex differences; Men; Women

Abstract: In order to determine the effect of age, sex, dietary and supplemental iron intakes and chronic disease factors on iron status and serum ferritin concentrations, 69 free-living very elderly (greater than or equal to 75 yr) were recruited. Subjects completed demographic and health history questionnaires, 3-day diet records and provided a fasting blood sample for determination of iron status parameters. Subjects were predominately white (mean age 80 yrs) and low income (52% less than \$10,000; 23% \$10,000-\$15,000, annually). Women had significantly lower hemoglobin, serum iron and percent transferrin saturation as compared with the men. Only one subject had a total iron intake less than 2/3 the RDA, but 22% supplemented with iron. Iron

Nutrition and the Elderly

deficiency anemia was prevalent in only 3% of the population; iron status in general, however, was lower than that reported in other free-living elderly. Neither serum ferritin, total iron intake, age, nor income were good predictors of iron status in this population. Serum ferritin was greater than 200 ng/inL in 38 of the subjects, but half of these high values could be explained by pre-existing disease factors.

143 NAL Call No: TX353.B76 1987

Jane Brody's nutrition book a lifetime guide to good eating for better health and weight control. Brody, Jane E.

Toronto ; New York: Bantam Books; 1987.
xx, 552 p. : ill. ; 23 cm. Includes bibliographies and index.

Language: English

Descriptors: Nutrition; Diet; Nutrition; Nutrition; Preventive nutrition; Health hazards; Diet planning; Nutrient sources; Recipes

Abstract: A sensible approach to good nutrition throughout life is based on making constructive changes in life style, practicing moderation, and understanding basic nutrition principles and needs. Consumers concerned about additives, alcohol, caffeine, cholesterol, salt, or sugar will find facts and discussions clarifying cloudy issues. Diet components--protein, identified and their nutritional aspects examined. Health hazards associated with everything from alcohol to water are discussed. Special dietary needs are identified for pregnant and lactating women, children, adolescents, athletes, vegetarians, and the elderly. The problems of obesity, the failures of diet fads, and tips for successful weight control are discussed. The text is interspersed with charts, tables, recipes, and menus.

144 NAL Call No: 389.8 F7398

Lifestyle-driven foods and ingredients required for their development.

Miller, G.A.; Frier, H.I.

Chicago, Ill. : Institute of Food Technologists; 1989 Apr.

Food technology v. 43 (4): p. 136, 138, 140, 143; 1989 Apr. Paper presented at the "Symposium on Changing Trends in Diet, Nutrition, and Consumer Taste: Implications for Food Technologists," June 19-22, 1988, New Orleans, Louisiana. Includes 16 references.

Language: English

Descriptors: Food industry; Product development; Nutritional value; Disease prevention; Aging; Life style; Consumer attitudes

Abstract: Changes in American demographics and lifestyles will provide opportunities for the food industry to produce new products and ingredients. The aging population, changing family structure and increased demand for ethnic foods will influence the makeup of our food supply. Concern for health and fitness will spur development of nutrient-dense foods and nutrient-free foods.

145 NAL Call No: 389.8 J824

Local action of oral 1,25-dihydroxycholecalciferol on calcium absorption in osteoporosis.

Francis, R.M.; Peacock, M.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Aug.

American journal of clinical nutrition v. 46 (2): p. 315-318. ill; 1987 Aug. Includes 21 references.

Language: English

Descriptors: Vitamin D; Osteoporosis; Absorption; Calcium; Nutrient-nutrient interactions; Bone mineralization; Women

Abstract: Extract: To investigate whether low calcium absorption in osteoporosis improves by increasing 1,25-dihydroxyvitamin D both systemically in plasma and locally in gut, the effects of oral 25-hydroxycholecalciferol and oral 1,25-dihydroxycholecalciferol on plasma 1,25-dihydroxy-vitamin D (1,25-(OH)2D) and calcium absorption were studied in 20 postmenopausal patients with vertebral osteoporosis. In 10 patients taking oral 0.25 micrograms 1,25-dihydroxycholecalciferol twice daily for 7 d, calcium absorption increased more than in 10 patients taking oral 40 micrograms 25-hydroxycholecalciferol once daily for 7 d (p less than 0.02) despite both groups having a similar increase in plasma 1,25-(OH)2D. These results support the view that the major effects of oral 1,25-dihydroxycholecalciferol on absorption is due to a local action on the gut and that it is possible to increase calcium absorption in osteoporosis with oral 1,25-dihydroxycholecalciferol without increasing its undesirable action on bone resorption.(author)

146 NAL Call No: TX341.C6

The MD Food Committee: hunger knows no season.

Washington, D.C. : Community Nutrition Institute; 1989 Feb09.

Quick Bibliography Series

Nutrition week v. 19 (6): p. 4-5; 1989 Feb09.

Language: English

Descriptors: Maryland; Hunger; Food distribution programs; Nutrition programs; State government; Low income groups; Children; Elderly

Abstract: This article is an excerpt from a report by the Maryland Food Committee. The report was based on a survey of 133 food pantries in Maryland. Food pantries distribute food to needy households. The demographics of the people receiving food aid are discussed.

147

NAL Call No: TX341.F69

Meeting a need efficiently.

King, P.

Cleveland, Ohio : Edgell Communications, Inc; 1989 Mar.

Food management v. 24 (3): p. 52, 55. ill; 1989 Mar.

Language: English

Descriptors: California; Food service; Cook-chill system; On-site food preparation; Nutrition programs; Elderly

Abstract: Feedback Foundation, a non-profit group, gives assistance to the elderly in Orange County, California. The need for its services, especially foodservice, has grown, necessitating a more efficient means of preparing and serving meals. They found the cook-chill method of preparation to be cost-effective and nutritious. Details of their foodservice operation are discussed.

148

NAL Call No: TX341.F69

Meeting elderly patients' needs with cook-chill.

King, P.

Cleveland, Ohio : Edgell Communications, Inc; 1989 Jun.

Food management v. 24 (6): p. 80, 82-83. ill; 1989 Jun.

Language: English

Descriptors: Missouri; Hospital food service; Cook-chill system, On-site food preparation; Long term care; Elderly

Abstract: A food and meal production system for a continuing care center is described. The center, which is located in John Knox Village, a retirement community outside Kansas City, Missouri, uses a cook-chill production system that involves the blast-chilling of bulk menu items, cold-plating of those foods well in advance of service and reheating of trayed foods immediately before meals are

served.

149

NAL Call No: TX361.A3J6

Meeting nutritional needs of Alzheimer patients who pace constantly.

Rheume, Y.; Riley, M.E.; Volicer, L.

Binghamton, N.Y. : Haworth Press; 1987.

Journal of nutrition for the elderly v. 7 (1): p. 43-52. charts; 1987. Includes 11 references.

Language: English

Descriptors: Elderly; Patient care; Alzheimer's disease; Energy intake; Energy expenditure; Nutritional requirements

Abstract: Extract: Energy intake and expenditure were observed in six patients suffering from dementia of the Alzheimer type. Three of these patients walked constantly and three walked only occasionally. Our clinical observation indicates that constant walking (pacing) increases energy demand by 1600 kilocalories/day. The implication of this observation and the nutritional strategies for prevention of metabolic imbalance in these patients are described.(author)

150

NAL Call No: 389.8 J824

Metabolic balance studies for zinc and copper in housebound elderly people and the relationship between zinc balance and leukocyte zinc concentrations.

Bunker, V.W.; Hinks, L.J.; Stansfield, M.F.; Lawson, M.S.; Clayton, B.E.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Aug.

American journal of clinical nutrition v. 46 (2): p. 353-359. ill., charts; 1987 Aug. Includes 38 references.

Language: English

Descriptors: Nutritional state; Zinc; Copper; Leukocytes; Balance studies; Elderly

Abstract: Extract: Five-day metabolic balance studies for zinc and copper were carried out in 20 housebound elderly people with stable chronic diseases. Subjects were age 70.0-85.1 y, lived in their own homes, and ate self-selected diets. Mean daily intakes for Zn and Cu were 90 micromoles and 13.4 micromoles with mean negative balances of -16 and -1.1 micromoles, respectively. Both balance values significantly differed from equilibrium (p less than 0.05). For Zn and Cu, respectively, mean concentrations were 11.3 and 22.1 micromoles/L plasma, 103 and 19.4 micromoles/L whole blood, and 98 and 8.7 pmol/10(6) leukocytes. The results

Nutrition and the Elderly

for Zn and Cu balance and leukocyte concentrations were different from those obtained by us for healthy elderly people (Am J Clin Nutr 1984;40:1096-120). A highly significant ($r = 0.75$, p less than 0.001) correlation was observed between zinc balance and leukocyte zinc concentrations.(author)

151 NAL Call No: 389.8 J824
Methodological and nutritional issues in assessing the oral health of aged subjects.

Carlos, J.P.; Wolfe, M.D.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.

American journal of clinical nutrition v. 50 (5): p. 1210-1218. charts; 1989 Nov. Includes 43 references.

Language: English

Descriptors: Dental health; Nutrition; Research methodology; Gingivitis; Paradontal disease; Dental caries; Epidemiology; Elderly

Abstract: Despite several serious methodological difficulties in epidemiological research on the oral health of elderly subjects, sufficient data are available to conclude that the prevalence of dental diseases is high in persons aged greater than 65 y in the US. An estimated 41% are completely edentulous, and dental caries, including root-surface caries, and destructive periodontal disease are widespread. Whether nutrition is related to the oral health problems of older persons is unclear; this question has received little research attention, and the results have generally been equivocal. There is some evidence that loss of teeth may be associated with suboptimal intake of nutrients and with changes in food preference.

152 NAL Call No: RA784.A1C3
A methodology for nutrition, health and lifestyle survey of elderly population living in communities.

Sinha, D.P.

Kingston : Caribbean Food and Nutrition Institute; 1987.

Cajanus v. 20 (1): p. 50-64. forms; 1987.

Language: English

Descriptors: Caribbean; Nutritional assessment; Elderly; Health; Dietary surveys; Life style; Methodology

Abstract: Practical steps are identified and discussed concerning the development and implementation of appropriate methodology for conducting

baseline surveys to assess the nutritional, health, and lifestyle status of elderly populations residing in the Caribbean. Attention is given to the survey characteristics, requisite preliminary investigations, the development of the survey instruments and their pre-testing, sampling strategies, essential characteristics concerning the design and conduct of the survey, data analysis, and the presentation of the report and the dissemination and utilization of the survey results. A 50-item questionnaire for a baseline survey is appended.(wz)

153 NAL Call No: 389.8 J824
Moderate alcohol intake and nutritional status in nonalcoholic elderly subjects.

Jacques, P.F.; Sulsky, S.; Hartz, S.C.; Russell, R.M. Baltimore, Md. : American Society for Clinical Nutrition; 1989 Oct.

American journal of clinical nutrition v. 50 (4): p. 875-883. charts; 1989 Oct. Includes 53 references.

Language: English

Descriptors: Nutritional state; Alcoholic beverages; Nutrient intake; Nutritional surveys; Caloric intake; Vitamins; Minerals; Elderly; Men; Women

Abstract: Three hundred and seventy-three female and 213 male nonalcoholic subjects, aged 60-100 y, who had participated in a nutritional status survey of elderly people in the Boston area were grouped according to usual alcohol intake: 0-4, 5-14, or 15+ g/d. The age- and sex-adjusted mean intake of calories, fat, protein, carbohydrate, and 10 micronutrients and the mean levels of 14 nutrient and 22 nonnutrient biochemical indices were compared for the three categories of alcohol intake. The mean micronutrient intakes were also adjusted for total caloric intake and the mean nutrient biochemical concentrations were also adjusted for the corresponding nutrient intakes. The results suggest that caloric intake and blood concentrations of retinol, iron, ferritin, HDL cholesterol, AST, and ALT increased with increasing alcohol intake whereas folate and phosphorus intakes and blood measures of riboflavin, copper, zinc, urea nitrogen, and creatinine decreased with increasing alcohol intake.

154 NAL Call No: TX361.A3N34
A National survey of nutritional risk among the elderly.

Food Research and Action Center

Washington, D.C. : Food Research and Action Center; 1987.

1 v. (various pagings) ; 28 cm. March 1987. In-

Quick Bibliography Series

cludes bibliographical references.

Language: English

Descriptors: Nutrition surveys; Aged--Nutrition; Elderly nutrition; Nutritional surveys

Abstract: This survey of the nutritional status of the elderly is based on 3602 responses by senior citizens to a questionnaire distributed during December, 1984 and January, 1985. In this self-administrated survey, affirmative answers to the "risk questions" indicated a greater chance that the respondent lacked a component of a healthy diet. Results: 35 percent responded that they eat less than three meals per day, 18.1 percent indicated "I do not have enough money to buy the food I need," 20.2 percent had lost weight without trying over the last month, and 21.7 percent gave responses which indicated nutritional risk to five or more of the survey questions. Recommendations are included.

155 NAL Call No: 389.8 J824
Need for body composition information in elderly subjects.

Kuczumarski, R.J.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.

American journal of clinical nutrition v. 50 (5): p. 1150-1157. charts; 1989 Nov. Includes 59 references.

Language: English

Descriptors: Body composition; Aging; Nutritional assessment; Body fat; Anthropometric dimensions; Elderly; Men; Women

Abstract: The need for accurate body composition information in elderly subjects is becoming more important as greater numbers of people reach older ages. An overview is given to describe some of the practical reasons why body composition data are necessary in various applied settings. Major changes in the body mass components and body dimensions are described to provide an enhanced awareness of the utility and increased needs for body composition information in applied and research settings. These needs for additional body composition data in elderly subjects include the following uses: to prevent malnutrition in institutionalized persons; to screen for health risks; to plan intervention and evaluate therapy; to study mechanisms of fat pattern change and correlates of stature loss; to study associations among fat patterning and mortality; as a prognostic indicator for conditions receiving treatment; and to develop im-

proved reference standards for ambulatory and nonambulatory elderly persons.

156 NAL Call No: 389.8 J824
NHANES III for older persons: nutrition content and methodological considerations.

Harris, T.; Woteki, C.; Briefel, R.R.; Kleinman, J.C.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.

American journal of clinical nutrition v. 50 (5): p. 1145-1149; 1989 Nov. Includes 13 references.

Language: English

Descriptors: Nutritional surveys; National surveys; Nutritional assessment; Elderly nutrition; Research methodology

Abstract: The National Health and Nutrition Examination Surveys (NHANES) are important in the assessment of nutritional status of the population of the United States. The utility of these surveys for assessment of the nutritional status of older Americans has been limited because prior NHANES have not included persons aged greater than or equal to 75 y. This paper reviews the role of the NHANES for nutritional epidemiology, highlighting the unique opportunity NHANES III offers to expand the nutrition database for older persons. Data are presented on consequences of nonresponse in analytic work and mechanisms that have been devised to approach the potential problem of nonresponse within NHANES III.

157 NAL Call No: 389.8 B773
Nitrogen balance studies in apparently healthy elderly people and those who are housebound.

Bunker, V.W.; Lawson, M.S.; Stansfield, M.F.; Clayton, B.E.

Cambridge : Cambridge University Press; 1987 Mar.

The British journal of nutrition v. 57 (2): p. 211-221. charts; 1987 Mar. Includes references.

Language: English

Descriptors: Man; Elderly; Nitrogen; Balance studies; Food intake; Protein requirement

Abstract: Extract: Metabolic balance studies (5 d) for nitrogen were carried out in twenty-four apparently healthy elderly people (age 69.7-85.6 years) and a heterogeneous group of twenty housebound elderly people (age 69.9-85.1 years) with chronic diseases. During the study all subjects ate self-selected diets, lived in their own homes and continued their normal daily activities. Seven of the

Nutrition and the Elderly

housebound received meals-on-wheels 5 d/week. Healthy men and women had mean metabolizable energy intakes of 8.7 and 6.6 MJ/d respectively compared with 6.3 and 4.8 MJ/d in the housebound. The average energy content of the meals-on-wheels as delivered was 2.4 MJ per meal, of which 2.1 MJ were consumed. The healthy men and women had average daily protein intakes of 69.4 and 59.7 g respectively compared with 46.3 and 39.1 g in the housebound. Meals-on-wheels as supplied provided 19.4 g protein per meal, of which 16.2 g were consumed. Healthy subjects were in equilibrium for N balance (0 mmol/d) with a daily intake of 733 mmol, which was equivalent to 11.04 mmol N (0.97 g mixed protein)/kg body-weight per d. Housebound individuals were in negative N balance (-95 mmol/d) with an intake of 475 mmol/d, corresponding to 7.59 mmol N (0.67 g mixed protein)/kg body-weight per d. We were unable to determine in the present study whether the negative N balance observed in the housebound people was due to the relatively low N intake or the underlying disease condition. (author)

158 NAL Call No: RM214.N8
No effect on serum lipids by moderate and high doses of vitamin C in elderly subjects with low plasma ascorbic acid levels.
Aro, A.; Kyllastinen, M.; Kostiainen, E.; Gref, C.G.; Elfving, S.; Uusitalo, U.
Basel : S. Karger; 1988 May.
Annals of nutrition and metabolism v. 32 (3): p. 133-137. charts; 1988 May. Includes 27 references.

Language: English

Descriptors: Diet; Ascorbic acid; Vitamin supplements; Blood lipids; Cholesterol; High density lipoprotein; Elderly

Abstract: The influence of dietary supplementation with moderate (200 mg/day) and high (2,000 mg/day) doses of vitamin C on serum lipid levels was studied in 27 female long-stay hospital patients characterized by low plasma ascorbic acid levels during the preceding year. The two doses of vitamin C were compared with placebo in a double-blind, cross-over design during randomly determined 6-week periods followed by 2-week washout intervals. No effect was observed on serum cholesterol, HDL cholesterol, triglyceride levels. Plasma ascorbic acid levels were highly significantly increased (p less than 0.001) by both doses of vitamin C. It is concluded that dietary supplementation with moderate or high doses of vita-

min C does not affect serum lipids of persons who have low plasma ascorbic acid levels suggestive of possible marginal deficiency of vitamin C.

159 NAL Call No: 389.8 AM34
The nutrient intake of a group of older South African hospitalized persons.
Gouws, E.; Rossouw, J.E.; Labadarios, D.; Van Eck, M.
Chicago, Ill. : The Association; 1989 Feb.
Journal of the American Dietetic Association v. 89 (2): p. 255-257. charts; 1989 Feb. Includes 15 references.

Language: English

Descriptors: South africa, Nutrient intake, Hospital diets; Nutritional state; Elderly

Abstract: As little is known about the nutritional status of older patients hospitalized for long-term rehabilitation, the present study was undertaken to assess the dietary intake of a group of such patients. Inadequate nutrition can influence both the chance of recovery and the duration of hospitalization.

160 NAL Call No: 389.8 AM34
Nutrient intake recommendations needed for the older American.
Andres, R.; Hallfrisch, J.
Chicago, Ill. : The Association; 1989 Dec.
Journal of the American Dietetic Association v. 89 (12): p. 1739-1741; 1989 Dec. Includes 4 references.

Language: English

Descriptors: Elderly nutrition; Recommended dietary allowances; Nutrient requirements; Minerals; Vitamins; Nutrition knowledge

Abstract: Physiological, social, economic, and psychological factors all affect nutritional intake in older Americans. Diseases, physical impairments, and mental deterioration can make maintenance of nutritional status complicated. Until recently, little notice was paid to the dietary needs and habits of older Americans except in relation to restrictions imposed by chronic diseases they acquired on their way to old age. Recommended dietary allowances published in 1980 made few distinctions between requirements for young and old adults. The major change recommended as age increased was a reduction in caloric intake at ages 51 and 75. This reduction is in response to the lower metabolic rates and physical activity of many older people.

Quick Bibliography Series

161 NAL Call No: TX361.A3J63
Nutrient intake status, knowledge, source of information and self-perceived health status among older adults in South Carolina.
Ryan, V.C.; Gates, A.D.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 8 (4): p. 41-48; 1988. Includes 7 references.

Language: English

Descriptors: South Carolina; Nutrition knowledge; Nutritional state; Nutrient intake; Information sources; Elderly

Abstract: Data on 339 subjects who were over 51 years of age was analyzed and Pearson product variant correlation coefficients were estimated between the variables of nutrient intake, nutrition knowledge, source of information and self perceived health status. Nutrient status was based on the consumption of eight nutrients (protein, calcium, iron, vitamin A, vitamin C, niacin, riboflavin and thiamin). A significant positive relationship was found between nutrition status and nutrition knowledge, but no significant relationships were found between self-perceived health status and nutrition status or between self-perceived health status and nutrition knowledge. A significant difference was found between sources of nutrition knowledge (physicians, dietitians, nurses, etc.) and nutrition status using the chi-square statistic. There was no significant difference on a chi-square test of sources of nutrition information and self-perceived health status or on self-perceived health status and nutrition status. A stepwise multivariate approach was used to estimate the amount of variance contributed by the independent variables of nutrition knowledge and sources of nutrition information on the dependent variable of nutrition status.

162 NAL Call No: 389.8 AM34
Nutrient intakes and energy expenditures of residents with senile dementia of the Alzheimer's type.
Litchford, M.D.; Wakefield, L.M.
Chicago, Ill. : The Association; 1987 Feb.
Journal of the American Dietetic Association v. 87 (2): p. 211-213. charts; 1987 Feb. Includes 15 references.

Language: English

Descriptors: Elderly; Alzheimer's disease; Nutrient intake; Energy expenditure; Sex differences
Abstract: A 3-day study compared the nutrient in-

take and energy expenditure of 12 women and 3 men (ages 75-95) with senile dementia of the Alzheimer's type (SDAT) with 10 women and 3 men with similar medical profiles but lacking SDAT. The results showed that the SDAT subjects consistently exhibited lower nutrient intakes than the non-SDAT group. Significant intake differences were noted for total calories, fat, carbohydrates, vitamin A, thiamin, niacin, riboflavin, and calcium. The intake results are statistically summarized and discussed.(wz)

163 NAL Call No: TX361.A3J63
Nutrient intakes of a group of independently-living elderly individuals in Toronto.
Scythes, C.A.; Zimmerman, S.A.; Pennell, M.D.; Yeung, D.L.
Binghamton, N.Y. : Haworth Press; 1989.
Journal of nutrition for the elderly v. 8 (3,4): p. 47-66. charts; 1989. Includes 32 references.

Language: English

Descriptors: Canada; Nutrient intake; Nutritional state; Nutritional surveys; Elderly; Women; Men
Abstract: A cross-sectional survey of an age-stratified random sample of 619 independently-living adults (307 males, 312 females) aged 60 years and over, living in Toronto was conducted. Detailed dietary data was collected from three sets of 3-day food records. The mean energy and nutrient intakes appeared to be adequate. However the energy intake of many participants was low, while their fat intake was high. The nutrients of concern for inadequate intakes were calcium, vitamin D and vitamin A. Intakes of zinc, vitamin C and protein were low for a smaller group. Nutrition education is recommended to provide guidance for choosing healthy diets.

164 NAL Call No: 389.8 AM34
Nutrition, aging, and the continuum of health care: technical support paper.
Chicago, Ill. : The Association; 1987 Mar.
Journal of the American Dietetic Association v. 87 (3): p. 345-347; 1987 Mar. Includes 50 references.

Language: English

Descriptors: Elderly nutrition; Nutritional intervention; Disease prevention; Aging; Health care; Public services; Policy
Abstract: A technical review justifies the importance of a continuum of health care to help the aging maintain independent living for as long as possible, including the integration of professional

Nutrition and the Elderly

nutrition services, education and counseling, and research. Specific attention is given to the etiology of nutrition problems of the elderly, the causes of these problems, and effective intervention and preventive strategies. The limitations of existing methods and standards for assessing the nutritional status of the aging also is addressed.(wz)

165 NAL Call No: RC620.A1J6
Nutrition and aging: Dietary intake of "apparently healthy" elderly (Dutch Nutrition Surveillance System).

Lowik, M.R.H.; Westenbrink, S.; Hulshof, K.F.A.M.; Kistemaker, C.; Hermus, R.J.J.
New York, N.Y. : John Wiley & Sons; 1989 Aug.
Journal of the American College of Nutrition v. 8 (4): p. 347-356. charts; 1989 Aug. Includes 43 references.

Language: English

Descriptors: Netherlands, Nutrient intake, Dietary surveys; Dietary history; Nutritional adequacy, Food consumption; Elderly; Men; Women

Abstract: This article details dietary surveys of healthy elderly men and women. Nutrient intake, nutritional supplements, food choices and eating habits are discussed and shown on tables. Some results were compared with statistics for younger men.

166 NAL Call No: RA784.F6
Nutrition and bone.

Clifton-Bligh, P.

London : Libbey; 1987.

Food and health . issues and directions / edited by Mark L. Wahlqvist ... [et al.]. p. 74-77; 1987. (International perspectives in food, diet and health, 0951-869x ; 1). Includes 17 references.

Language: English

Descriptors: Bone mineralization; Vitamin D, Phosphorus; Osteoporosis; Disease prevention

Abstract: A brief technical report discusses the influence of nutritional factors on bone health. Particular attention is given to calcium, vitamin D, and phosphorus and their interrelationships in promoting/maintaining bone health. Nutrient requirements for infants, growing adolescents, and the elderly are addressed, as well as nutritional approaches for managing osteoporosis in the elderly.(wz)

167 NAL Call No: 389.8 AM34
Nutrition and dementia.

Gray, G.E.

Chicago, Ill. : The Association; 1989 Dec.

Journal of the American Dietetic Association v. 89 (12): p. 1795-1802. charts; 1989 Dec. Literature review. Includes 91 references.

Language: English

Descriptors: Nutrient intake; Alzheimer's disease; Nutritional assessment; Therapeutic diets; Supplements; Aluminum; Alcoholism; Vitamin Deficiency; Elderly

Abstract: The nutritional consequences of dementia and the role of diet in the etiology, treatment, and prevention of dementia are the subjects of this review. The major cause of dementia is Alzheimer's disease. Although it has been suggested that aluminum intake may cause this disease, the bulk of scientific evidence suggests that this is unlikely. Dietary supplements of choline and lecithin have been used to treat Alzheimer's disease but are ineffective. Alzheimer's disease patients are at risk of developing protein-energy malnutrition because of poor food intake and increased energy requirements. The second most common cause of dementia is multi-infarct dementia, caused by multiple strokes. Diet may play a role in the prevention of this form of dementia through effects on blood pressure and other risk factors. Control of risk factors may also prevent further progression of the dementia. Patients with multi-infarct dementia often have dysphagia. The third most common cause of dementia appears to be excessive alcohol intake, due both to the direct neurotoxic effects of alcohol and to the effects of alcohol on nutritional status. Alcoholic dementia may be at least partially reversible with abstinence and good nutrition. Other causes are vitamin B-12 and folate deficiencies; these are reversible dementias. In all types of dementia, adequate nutrition may improve physical well-being, help maximize the patient's functioning, and improve the quality of life.

18

NAL Call No: S22.C6

Nutrition and health.

Ames, Iowa : The Council; 1987 Mar.

Council for Agricultural Science and Technology report v. 111: p. 4-7. charts; 1987 Mar. Includes 15 references.

Language: English

Descriptors: Dietary guidelines; Nutrient requirements; Obesity; Aging; Health promotion

Abstract: An expert task force report defines nutri-

Quick Bibliography Series

tion as the interaction between food components and living organisms, and discusses child development, the aging process, and problems of over-nutrition and undernutrition. Attention also is given to Recommended Dietary Allowances for recommended nutrient intakes and Dietary Guidelines for Americans for recommended food habits. Specific attention is given to the effects of special dietary problems (obesity) and chronic diseases of aging on nutritional status, with longevity and life quality given as products of healthful dietary habits.(wz)

169 NAL Call No: RA784.N8
Nutrition and health symposium: The University of Texas at Austin, April 1988--summary report. Lochniskar, M.
Baltimore, Md. : Williams & Wilkins; 1988 Oct.
Nutrition today v. 23 (5): p. 31-35; 1988 Oct.

Language: English

Descriptors: Food allergies; Food fiber; Diabetes. Elderly; Nutrition, Cardiovascular diseases, Food preferences

Abstract: The Jean Andrews Visiting Professorship Lecture Series at The University of Texas at Austin, is designed to present current information in the field of nutrition. This year the symposium was attended by over 300 participants and addressed many areas of nutrition including: food allergies; the effect of fiber on diseases such as cancer, diabetes and heart disease; and nutrition in aging ranging from the prevention and therapy of osteoporosis by nutrition, defining how the elderly make their food choices and changing the RDAs for the elderly.

170 NAL Call No: RC682.N87
Nutrition and heart disease.
Watson, Ronald R.
Boca Raton, Fla. : CRC Press; 1987.
2 v. : ill., 27 cm Includes bibliographies and indexes.

Language: English

Descriptors: Heart--Diseases--Nutritional aspects, Heart--Diseases--Prevention; Heart--Diseases--Diet therapy; Reference works; Cardiovascular diseases; Dietary factors; Disease prevention; Aging; Patient care, Nutrient requirements; Risks; Reviews

Abstract: A 2-volume reference set for nutrition researchers, clinical dietitians, cardiologists, and other health professionals presents 19 authoritative

reviews prepared by experts in their respective fields concerning various key aspects of the role of nutrition in the prevention, cause, and treatment of heart and vascular diseases, covering both experimental animal and human clinical studies. Topics presented in Volume I (9 reviews) include: the effects of dietary fiber, dietary fats, oxidized lipids, and other dietary factors on heart and vascular diseases in adults and the elderly, including the effects of diet on cholesterol metabolism, lipoproteins, hypertension, stroke, and atherosclerosis. The 10 remaining reviews (Volume II) address: the effects of dietary macronutrients on blood pressure; the cardiovascular effects of caffeine and alcohol; the effects of fat-soluble vitamins, herbs and herbal formulas, and trace elements in preventing and treating heart disease; cardiac function in malnutrition and the causes, prevention, and treatment of "hospital malnutrition"; the association between diet, immune function, and cardiovascular disease; and nutrition and heart disease in the aging. Representative data tabulations and illustrations are included, and copious literature citations are appended to each of the reviews.

171 NAL Call No: RC682.N87
Nutrition and heart disease in the aging.
Debry, G.
Boca Raton, Fla. : CRC Press; 1987.
Nutrition and heart disease / editor, Ronald R. Watson. v. 2 p. 119-140; 1987. Literature review. Includes 193 references.

Language: English

Descriptors: Cardiovascular diseases; Elderly nutrition; Energy requirements; Nutrient requirements; Aging

Abstract: A detailed technical review deals specifically with a general and comprehensive view of the relationship between diet and heart disease in aging. Following a discussion of the influence of aging on cardiac function, key nutritional and biochemical changes that occur in aging, and energy and nutrient requirements for the elderly, attention is directed to heart diseases commonly found in the elderly. A structured examination of the short-term and long-term effects of the intakes of energy and other nutrients (protein, carbohydrate, fat, minerals, alcohol) on heart disease is included.

172 NAL Call No: 389.8 D14
Nutrition and modern lifestyles.
McBean, L.P.

Nutrition and the Elderly

Rosemont, Ill. : National Dairy Council; 1988 Sep. Dairy Council digest v. 59 (5): 6 p.; 1988 Sep. Literature review. Includes 13 references.

Language: English

Descriptors: Food habits; Eating patterns; Consumer attitudes; Household income; Family size; Employed women; Nutritional adequacy; Convenience foods; Consumer behavior; Nutrition surveys; Low calorie foods

Abstract: Modern lifestyles are influenced by recent demographic, economic, and social changes. Among these are an aging but more affluent population, a growing number and proportion of smaller (one or two person) households, and more women in the work force. Also, the trend in the 1980s toward self improvement has stimulated interest in health, fitness, and nutrition. These factors, in turn, have influenced the way people purchase, prepare, and consume food. The greater percentage of women in the work force and today's fast-paced lifestyle have led to a growing demand for convenience, especially as related to food. Modern lifestyles also are characterized by more fragmented meals--"eating on the run," eating away from home, buying take-out foods, or ordering foods for home delivery. The nutritional implications of these new eating patterns are being questioned. However, it is the actual foods consumed throughout the entire day, not necessarily where or when people eat, that determine nutritional adequacy. The increased health consciousness of Americans is responsible for their heightened concern about the nutritional content of food, in particular fat, cholesterol, sodium/salt, sugar, vitamins, and minerals. Improving nutritional intake/status is being viewed by some consumers not just as a means to avoid specific chronic diseases but to improve the quality of life. Consumers' purchasing behavior in grocery stores and when eating out reflects their health perceptions and lifestyle. In response to restaurant patrons' growing awareness of nutrition and health, many restaurants and foodservice establishments are offering healthy menu options and preparation methods. Similarly, food manufacturers, cognizant of this health trend, have developed new food products. In recent years, growth of sales of "healthy" and "diet" foods has been greater than that for their regular counterparts. Although consumers are interested in a healthy lifestyle, they behave in seemingly inc

173

Nutrition and the aging brain.

NAL Call No: RM214.N87

DeBruyne, L.K.

Philadelphia, Pa. : George F. Stickley Company; 1988 Dec.

Nutrition clinic, v. 3 (6): p. 1-13. ill., charts; 1988 Dec. Literature review. Includes 32 references.

Language: English

Descriptors: Elderly nutrition; Brain; Senility; Nutrient deficiencies; Vitamin B12; Thiamine; Iron; Zinc; Mental ability; Nutritional state

Abstract: This article begins with a brief description of the brain and how it works. Normal aging of the brain is discussed as well as abnormal preventable aging due to moderate long-term nutrient deficiencies. Nutrient deficiencies associated with loss of memory and impaired cognitive function are discussed.

174

Nutrition and the occurrence and prevention of cardiovascular disease in the elderly.

Kannel, W.B.

Washington, D.C. : Nutrition Foundation; 1988 Feb.

Nutrition reviews v. 46 (2): p. 68-78. ill., charts; 1988 Feb. Includes 35 references.

Language: English

Descriptors: Cardiovascular diseases; Elderly; Disease prevention

Abstract: A technical overview discusses recent evidence concerning the effect of diet on the prevention of cardiovascular (CV) disease in elderly subjects. Attention is given to: dietary practices and nutrient requirements of the elderly; CV risk factors in the elderly; blood lipid trends with age; the effect of obesity on CV disease risk; the role of blood pressure and its control for reducing CV disease risk; and dietary effects on blood lipid composition. Controlled clinical trials linking dietary nutrients to serum lipid composition and CV disease incidence also are discussed

175

Nutrition education and changes in nutrition behavior among the 65-74-year-old population of eastern Finland.

Kivela, S.L.; Nissinen, A.

Philadelphia, Pa. : George F. Stickley Company; 1987 Apr.

Journal of nutrition education v. 19 (2): p. 77-82. charts; 1987 Apr. Includes 32 references.

Language: English

Quick Bibliography Series

Descriptors: Finland; Nutrition education; Elderly nutrition; Health promotion; Behavior modification; Consumption patterns; Regional surveys

Abstract: A 1982 survey assessed the sources, quality, and usefulness of nutrition information available in eastern Finland, and evaluated reported changes in nutrition behavior in the 65-74 elderly age group of the population. The results indicated that: the principal nutrition information sources were television and radio; women obtained more nutrition information from reading materials than did men; while the elderly were able to understand the information provided, only about half were able to follow the guidelines provided; and that changes in improved nutrition habits occurred less in men (up to 35% of the men) than in women (up to 57%). It also was determined that the elderly decreased their dietary fat intake and experienced positive health changes, and that non-smokers had more positive nutritional health changes.(wz)

176 NAL Call No: TX361.A3J63
A nutrition education workshop for managers of nutrition centers for the elderly utilizing videotaped simulations for assessment.
Stanek, K.L.; Fox, H.M.

Binghamton, N.Y. : Haworth Press; 1989.
Journal of nutrition for the elderly v. 8 (3,4): p. 101-106; 1989. Includes 6 references.

Language: English

Descriptors: Nutrition education; Workshops (programs); Program effectiveness; Videotapes; Elderly

Abstract: As funds become increasingly scarce, an effective nutrition education method which reaches large numbers of elderly individuals is appropriate. A nutrition workshop was provided for 30 managers of government-sponsored nutrition centers for the elderly in a 5-county area. The workshop curriculum, determined after interests of both center managers and participants at the nutrition centers were assessed, was presented in a packet of materials to each workshop attendee and supplemented with explanations and demonstrations. A pre- and post-test evaluation instrument, which included videotaped simulations of typical nutritional situations encountered with elderly persons at the centers, was used to measure attitudes and knowledge before and after the workshop. As a result of the workshop, knowledge increased significantly as measured by the test instrument and 75% of the managers reported use of materials and

ideas during the 5 months following the workshop.

177 NAL Call No: TX341.N838
Nutrition for senior citizens.

Rozovski, S.J.
New York, N.Y. : Columbia University, Institute of Human Nutrition; 1987.
Nutrition and health v. 9 (4): p. 1-6. charts; 1987.

Language: English

Descriptors: Elderly nutrition; Nutrient requirements; Nutrient intake; Guidelines

Abstract: A report provides nutritional guidelines for physicians and other health professionals treating elderly patients. The need for specific nutrients is described, covering intakes of energy, protein, fat, carbohydrate, fiber vitamins and minerals (Ca, P, Fe). Data on dietary recommendations for elderly men and women are included.(wz)

178 NAL Call No: 389.8 N953
Nutrition, immunity, aging, and cancer.

Good, R.A.; Lorenz, E.
Washington, D.C. : Nutrition Foundation; 1988 Feb.
Nutrition reviews v. 46 (2): p. 62-67; 1988 Feb.
Literature review. Includes 69 references.

Language: English

Descriptors: Carcinoma; Immunology; Diet; Aging; Nutritional state; Protein energy malnutrition; Zinc

Abstract: A technical report summarizes and discusses the results of recent studies concerning the effect of nutritional status on immunity, aging, and cancer risk. Attention is given to: associations between protein-energy malnutrition and decreased immune function; the role of zinc in immunocompetence; the effect of nutrition on immune function and diseases of aging; and rodent studies of dietary prevention of breast cancer. Recent approaches to understanding the effect of restricted energy intake on immune function also are discussed.

179 NAL Call No: QP141.A1N88
Nutrition, immunity, and outcome; past, present, and future. 11th Gopalan Gold Medal Oration.

Chandra, R.K.
Elmsford, N.Y. : Pergamon Press; 1988 Mar.
Nutrition research v. 8 (3): p. 225-237. ill; 1988 Mar. Includes 37 references.

Language: English

Descriptors: Nutrition; Cell mediated immunity;

Nutrition and the Elderly

Neonates; Elderly

Abstract: Extract: Nutrition is a critical determinant of immunocompetence and risk of illness. Young children with protein-energy malnutrition exhibit increased mortality and morbidity, due largely to infectious disease. Recent work has demonstrated that undernourished individuals have impaired immune responses. The most consistent abnormalities are seen in cell-mediated immunity, complement system, phagocytes, mucosal secretory antibody response, and antibody affinity. These changes, together with other handicapping factors observed in underprivileged societies, lead to more infections, which in turn produce physiological changes that worsen nutritional status. It is now established that deficiencies of single nutrients also impair immune responses. The best studied are zinc, iron, vitamin B6, vitamin A, copper, and selenium. If malnutrition occurs during fetal life, as epitomized by small-for-gestational age infants, the effects on cell-mediated immunity are very significant and long lasting. There is much recent evidence to suggest that at the other end of the age spectrum, namely in old age, nutrition plays an important role in maintenance of optimum immunity. Response to influenza virus and other vaccines is improved when the nutritional deficiencies in the elderly are corrected. These interactions of nutrition and immunity have several practical applications. (author)

180 NAL Call No: TX361.A3J6
Nutrition knowledge test for older adults.
Mwonya, R.; Ralston, P.A.; Beavers, I.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (4): p. 3-16.
charts; 1987. Includes 16 references.

Language: English

Descriptors: Nutrition knowledge, Elderly; Questionnaires; Regional surveys; Nutrition education, Socioeconomic status

Abstract: Extract: Purposes were to assess the effectiveness of an instrument designed to measure nutrition knowledge of older adults, and to use this instrument to compare nutrition knowledge of older adults according to age, sex, education, occupation and sources of nutrition information. A 24-item nutrition knowledge test was adapted for use with older adults, and was administered to 104 subjects (60 years of age and older) from two senior citizen centers in a medium-sized Midwestern community. Data were analyzed using an item analysis to determine the best and poorest items,

and one-way analyses of variance (ANOVA) to test for significant differences in nutrition knowledge by selected socio-demographic variables. The results revealed that the raw scores of the nutrition knowledge test ranged from 4 to 20 with a mean of 13.17 and standard deviation of 3.74. Fifteen items met the item analysis criteria, while the remaining six items were judged as poor. The results of the ANOVA showed significant differences in nutrition knowledge by educational level and occupation, with those with college education backgrounds and professional occupations having higher scores than other groups. Findings suggest that objective tests can be developed and used with older adult audiences, and that nutrition education efforts need to be focused on older adults from lower socioeconomic backgrounds. (author)

181 NAL Call No: TX341.N836
Nutrition myths about the elderly.
Van Nuys, Calif. : The Journal; 1989 Feb.
Nutrition & the M.D. v. 15 (2): p. 3; 1989 Feb.

Language: English

Descriptors: Elderly nutrition; Food beliefs; Caloric intake; Nutrient deficiencies; Food preferences

Abstract: Five myths about elderly nutrition are listed. The article discusses the amounts of food consumed, the need for soft foods, the need for vitamin supplement, food choices by the elderly and the relationship between nutrition and income of the elderly.

182 NAL Call No: 389.8 Z33
Nutrition of the elderly in a Mediterranean city in Spain: effects of life-style patterns.
Fernandez-Ballart, J.; Gordillo, B.; Arija, V.; Marti-Henneberg, C.
Bern : Hans Huber; 1989.
International journal for vitamin and nutrition research v. 59 (1): p. 8-13. charts; 1989. Includes 20 references.

Language: English

Descriptors: Spain; Food intake; Dietary surveys, Energy intake; Nutrient intake; Life style; Elderly

Abstract: The authors studied food intake of elderly men and women living in Reus, Spain. All of the subjects lived in non-institutional homes but with different aged companions. The type of life-style and age group of living companions were found to be the major contributing factors in the daily intake of nutrients and calories from proteins, lipids, and carbohydrates. The authors found that the greatest

Quick Bibliography Series

risk of energy deficiency exists among the elderly who live with the same age group especially in the case of men.

183 NAL CrII No: 389.8 AM34
A nutrition screening and assessment system for use with elderly in extended care.
Tramposch, T.S.; Blue, L.S.
Chicago, Ill. : The Association; 1987 Sep.
Journal of the American Dietetic Association v. 87 (9): p. 1207-1210. ill; 1987 Sep. Includes 17 references.

Language: English

Descriptors: Nutritional surveillance; Elderly; Nutritional assessment; Patient care

Abstract: Extract: One area of particular interest to health care professionals that is undergoing considerable investigation is the nutritional status of the elderly. This article describes the development, implementation, and evaluation of a procedure used to nutritionally screen and assess geriatric patients in extended care at the Veterans Administration Medical Center in Houston. As we developed the system, we considered the special needs and characteristics of geriatric individuals and incorporated them into two forms. Designs were selected to make our method comprehensive, concise, and ideally, time-efficient. The evaluation of the system included a time utilization study. This study was conducted to evaluate the efficiency of the new system compared with the efficiency of methods used hospital-wide in the past. Results suggest that the new system for reassessing the patient's nutritional status and reevaluating the nutrition care was three times more efficient.(author)

134 NAL Call No: 389.8 AM34
Nutrition screening for long-term care residents.
Noel, M.B.; Wojnarowski, S.M.
Chicago, Ill. : The Association; 1987 Nov.
Journal of the American Dietetic Association v. 87 (11): p. 1557-1558. forms; 1987 Nov. Includes 10 references.

Language: English

Descriptors: Nutritional surveillance; Patient care; Elderly nutrition; Hospitals

Abstract: A brief report describes the rationale for and development of nutritional screening in a 320-bed geriatric-psychiatric state hospital serving residents having an average length of stay of 806 days. Criteria are specified for an appropriate nutrition

screening tool, which was developed by a critical review of the literature concerning screening tools with consultation with a consulting firm of registered dietitians. This tool, a patient-oriented data form, was found to be totally consistent with standard methods of nutritional evaluation in identifying patients requiring nutrition services.(wz)

185 NAL Call No: TX354.N8
Nutrition throughout the life cycle.
Schlenker, Eleanor D.; Worthington-Roberts, Bonnie S.,1943-
St. Louis : Times Mirror/Mosby Colege Pub: 1988. xxiv, 597 p. : ill. ; 24 cm. Charts on lining paper. Includes bibliographies and index.

Language: English

Descriptors: Nutrition; Teaching materials; Textbooks; Nutrition; Life cycles; Growth stages; Post secondary education; Professional education

Abstract: This comprehensive text is designed primarily for students in upper division courses in life cycle nutrition and for health professionals working in both individual and community programs. Focus is on human growth and development needs at each stage of life: pregnancy and lactation, infancy, childhood, adolescence, adults and the aging process, and special needs of the elderly in the aging population. Discussions are provided on the changing body at each stage in the human life cycle; ways of determining nutritional status and needs along the way; the special needs of persons carrying increased risk at different ages and stressful life situations, and new approaches to nutrition education for health promotion. This text includes illustrations, appendices, charts, learning aids, and a glossary.

186 NAL Call No: TX361.A3J6
Nutritional and dental status of title III meal recipients.
Kunkel, M.E.; Chesnut, C.K.; Hoover, J.L.B.; Roughead, Z.K.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (3): p. 17-32. charts; 1987. Includes 34 references.

Language: English

Descriptors: South Carolina; Elderly nutrition; Supplemental feeding programs; Nutritional state; Dental health; Thiamine; Niacin; Calcium; Ethnicity; Urine analysis; Anthropometric dimensions; Regional surveys

Abstract: Seventy-one subjects aged 60 years and

Nutrition and the Elderly

older were recruited through Title III congregate dining sites and one church in northwestern South Carolina. Each participant was given a dental examination; had height, weight and triceps skinfold thickness measurements made; completed a 3-day dietary record; and collected a 24-hour urine sample. Urine samples were analyzed for creatinine, thiamin, niacin, phosphorus and nitrogen. The mean age of the participants was 73.5 +/- 5.8 years with a range of 62-89 years. Of the 37 dentulous subjects, all but 1 had periodontal disease. The only nutrient for which mean intakes were less than the RDA was calcium for black subjects. Thiamin excretion was adequate for all subjects while niacin excretion for 20 subjects was in the deficient range. There were no statistically significant relationships between dietary intake and oral health in these elderly subjects. (author)

187 NAL Call No: RC682.N87
Nutritional and pathological aspects of atherosclerosis in aging adults.
Guzman, M.A.; Strong, J.P.
Boca Raton, Fla. : CRC Press; 1987.
Nutrition and heart disease / editor, Ronald R. Watson. v. 1 p. 117-137. charts; 1987. Literature review. Includes 83 references.

Language: English

Descriptors: Louisiana; Puerto rico; Hawaii; Atherosclerosis, Dietary factors; Risks; Mortality, Epidemiology; Lipid metabolism

Abstract: A technical report summarizes and discusses the results of several major studies focused on the effects of dietary risk factors on pathological aspects of the etiology of atherosclerosis in aging adults. Specific attention is given to the methodology, dietary patterns, and results of 4 studies (with particular emphasis on the latter 3 studies), viz.: the international atherosclerosis autopsy project, conducted in 14 countries; the New Orleans autopsy studies, the Puerto Rico heart health program; and the Honolulu heart study. Statistical correlation results are presented and discussed.

188 NAL Call No: 389.8 J824
Nutritional factors in bone health in elderly subjects: methodological and contextual problems.
Heaney, R.P.
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.
American journal of clinical nutrition v. 50 (5): p. 1182-1189. charts; 1989 Nov. Literature review. In-

cludes 31 references.

Language: English

Descriptors: Bone strength, Calcium, Vitamin D, Mineral metabolism, Estrogens, Osteoporosis, Research methodology; Elderly

Abstract: Five problem areas relating to the assessment of nutritional influences on bone health in elderly subjects are described. These are 1) complexity of the domain of bone health, 2) difficulty in detecting bone fragility in vivo, 3) difficulty in detecting useful changes in bone mass, 4) remodeling adequacy of vitamin D nutrition in elderly subjects. The first four are inherent to the tissue being studied and the last is definitional. Strategies are available for dealing with some of these problems; others seem inherently intractable. However, understanding the basis for all of them can at least help avoid both entering upon futile research and misinterpretation of experimental results.

189 NAL Call No: TX361.A3J63
Nutritional management of the elderly trauma patient: a case report.
Tramosch, T.S.; Hester, D.D.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 8 (1): p. 45-57; 1988. Includes 38 references.

Language: English

Descriptors: Trauma, Patient care; Nutritional support; Nutrient requirements; Enteral feeding; Parenteral feeding; Case studies; Elderly

Abstract: The nutritional assessment and management of the elderly trauma patient is discussed. Nutritional requirements for protein, vitamins, minerals and fluids are included. A case report of a 65-year-old woman motor vehicle accident victim is presented.

190 NAL Call No: RA421.F35
Nutritional needs of the elderly: issues, guidelines, and responsibilities.
Roe, D.A.
Frederick, Md. : Aspen Publishers; 1989 May.
Family & community health v. 12 (1): p. 59-65; 1989 May. Includes 13 references.

Language: English

Descriptors: Elderly nutrition; Nutrition policy; Nutrient requirements; Physiology; Nutrient-drug interactions; Elderly

Abstract: The elderly have unique nutritional needs

Quick Bibliography Series

and problems influenced by environmental, genetic, physiologic and pharmacologic factors. Their food choices also affect their nutritional status and these choices may be influenced by income status and physical handicaps.

191 NAL Call No: TX341.C6
Nutritional problems common among elderly.
Washington, D.C. : Community Nutrition Institute; 1987 Apr16.
Nutrition week v. 17 (16): p. 4-5; 1987 Apr16.

Language: English

Descriptors: USA; Elderly nutrition; Federal programs; Food and nutrition controversies; Malnutrition; Socioeconomic status

Abstract: A review of assessments of US elderly at nutritional risk, compounding low income factors, and low food stamp program participation, concludes that elderly nutrition programs have received no significant funding increases for 5 years, despite long waiting lists reflecting the presence of a continuing unmet need. It is argued that the provision of food and resources to purchase food can ameliorate the undernutrition and illness that is associated with poor life quality and increased health care costs in old age.(wz)

192 NAL Call No: R11.C6
Nutritional problems in the elderly.
Morley, J.E.
Minneapolis, Minn. : General Mills, Inc; 1987.
Contemporary nutrition v. 12 (6): p. 1-2; 1987. Includes 20 references.

Language: English

Descriptors: USA; Elderly nutrition; Risks; Protein energy malnutrition; Anorexia; Vitamin supplements; Mineral deficiencies; Zinc; Reviews

Abstract: A brief review focuses on 3 areas of particular nutritional concern to the elderly, viz.: body mass issues related to longevity enhancement (including protein energy malnutrition (PEM) and the anorexia of aging); the use/abuse of vitamin supplements; and the role of trace elements in elderly nutrition. It is concluded that there is an increasing need for awareness of elderly nutritional problems, that elderly have higher risks for PEM and low zinc status, and that megadose vitamin supplements pose toxic risks.(mjs)

193 NAL Call No: TP368.C7
Nutritional requirements of the aged.
Bidlack, W.R.; Smith, C.H.

Boca Raton, Fla. : CRC Press; 1988.
Critical reviews in food science and nutrition v. 27 (3): p. 189-218. charts; 1988. Literature review. Includes 238 references.

Language: English

Descriptors: Elderly nutrition; Physiological functions; Diseases; Income; Nutritional state; Nutritional surveys; Food consumption; Recommended dietary allowances; Dietary guidelines; Dietary proteins; Dietary carbohydrates; Fats; Vitamins; Minerals

Abstract: The factors which influence the nutritional status of the elderly include physiological decline, health, medication, economics and nutrition knowledge. This article presents the results of nutrient intake studies and the recommended dietary allowances for the elderly.

194 NAL Call No: QP141.A1A63
Nutritional requirements of the elderly.
Munro, H.N.; Suter, P.M.; Russell, R.M.
Palo Alto, Calif. : Annual Reviews; 1987.
Annual review of nutrition v. 7: p. 23-49; 1987.
Literature review. Includes 171 references.

Language: English

Descriptors: Elderly nutrition; Nutrient requirements; Nutrient intake; Nutritional state; Osteoporosis; Trace elements; Vitamins; Nutrient-drug interactions

Abstract: A technical review summarizes and discusses the current knowledge and understanding of nutrient needs and intakes of elderly people, and the degree of their nutritional status risk. This review highlights the weaknesses of current recommended nutrient intakes for the elderly, noting the lack of solid data for such recommendations. Particular attention is given to separate discussions of: energy and protein intakes and requirements; Ca and other nutrients in osteoporosis; requirements for individual trace elements (Fe, Zn, Cu) and vitamins (A, C, D, E, K, B1, B2, B6, B12, folate, niacin, biotin, pantothenic acid); and data on the nutritional status of the elderly, as gleaned from national clinical surveys, nutrient-drug interaction studies, and nutritional intervention strategies.(wz)

195 NAL Call No: HQ1060.J68
Nutritional status and dietary intake in institutionalized patients with Alzheimer's disease and multiinfarct dementia.
Sandman, P.O.; Adolfsson, R.; Nygren, C.; Hallmans, G.; Winblad, B.

Nutrition and the Elderly

New York : Elsevier Science Publishing Company, 1987 Jan.

Journal of the American Geriatrics Society v. 35 (1): p. 31-38. charts; 1987 Jan. Includes 43 references.

Language: English

Descriptors: Elderly; Nutritional state; Nutrient intake; Protein energy malnutrition; Weight losses; Anthropometric dimensions; Alzheimer's disease

Abstract: A 5-week study of the nutritional status (anthropometry and clinical measurements) and dietary intake (weighed food diary) of 44 elderly patients (23 women, 21 men; mean age, 76) with Alzheimer's disease or multi-infarct dementia revealed the presence of energy and/or protein malnutrition in half the patients even though the mean dietary intake was adequate for energy, protein, vitamins, and minerals. A 4-fold higher incidence of infections was noted for malnourished than for non-malnourished patients and 89 per cent of the patients lost weight during hospitalization, but no association was found among weight loss, hospitalization duration, and length of illness.(wz)

196

NAL Call No: 389.8 Z33

Nutritional status, grip strength, and immune function in institutionalized elderly.

Suboticanec, K.; Stavljenic, A.; Bilic-Pesic, L.; Gorajscan, M.; Gorajscan, D.; Brubacher, G., Buzina, R.

Bern : Hans Huber; 1989.

International journal for vitamin and nutrition research v. 59 (1): p. 20-28. charts; 1989. Includes 28 references.

Language: English

Descriptors: Nutritional state; Nutrient intake; Blood plasma; Vitamins; Zinc; Immune response; Vitamin supplements; Strength; Men

Abstract. Elderly people are prone to vitamin deficiencies and malnutrition. This is probably due to a decline of basal metabolic rate and lean body mass as well as reduction in physical activity and total food consumption. The authors studied a group of elderly institutionalized men and women. Their food intake was recorded and certain essential minerals and vitamins found in their blood were measured. The effects of vitamin supplementation on grip strength and immunocompetence were analyzed.

197

NAL Call No: TX361.A3J6

Nutritional status of healthy middle class elderly women.

Bergstrom, N.; Wilson, S.E.; Smith, J.L. Binghamton, N.Y. : Haworth Press; 1987.

Journal of nutrition for the elderly v. 7 (1). p. 3-21. charts; 1987. Includes 48 references.

Language: English

Descriptors: Women; Elderly; Nutritional state, Socioeconomic status

Abstract: Extract: The purposes of this study were to determine the adequacy of dietary intake and blood/serum concentrations of nutrients of middle class elderly women. Thirty-six middle class women volunteered to participate in the study. An interview, a 24-hour dietary recall, anthropometric measurements and a blood sample were obtained from all subjects. Using the 1980 RDA as a basis for evaluation, less than two-thirds of the RDA of the following nutrients were ingested for a percentage of the subjects: calcium (41%), vitamin A (16%), thiamin (8%), and riboflavin and vitamin C (14%). Blood and serum levels of nutrients below normal clinical ranges included vitamin B6 (31% of subjects), carotene (23%), vitamin B12 (14%) and folate (11%). No correlations were found between intake and blood and serum concentrations.

198

NAL Call No: 389.8 Z33

Nutritional status of non institutionalized elderly people in North Italy.

Porini, M.; Testolin, G.; Simonetti, P.; Moneta, A.; Kovati, P.; Aguzzi, F.

Bern : Hans Huber; 1987.

International journal for vitamin and nutrition research v. 57 (2): p. 203-216. charts; 1987. Includes 19 references.

Language: English

Descriptors: Italy; Elderly; Nutritional state; Nutrient intake; Nutrient deficiencies; Anthropometric dimensions; Clinical investigations

Abstract: An anthropometric, dietary, and biochemical study of the nutritional status of 313 free-living and 37 publically-assisted elderly subjects revealed the assisted elderly to have lower body weight, percent body fat, muscle and fat areas, and a lower body mass index. The assisted elderly also had mean dietary intakes that were below recommended levels for 8 of 9 nutrients studied. While 10-15% of the free-living subjects had an inadequate thiamin and riboflavin status, these nutritional status deficiencies in the assisted elderly

Quick Bibliography Series

were greater. It also was noted that the free-living subjects had higher protein and fat intakes and lower thiamin and retinol intakes.(wz)

199 NAL Call No: 389.8 Z33
Nutritional status of the elderly in Naples.

Contaldo, F.; Di Biase, G.; Lanzetta, R.; Lanzilli, A.; Simonetti, M.S.; Fidanza, F.

Bern : Hans Huber; 1987.

International journal for vitamin and nutrition research v. 57 (2): p. 193-202. charts; 1987. Includes 23 references.

Language: English

Descriptors: Italy; Diet; Elderly nutrition; Mineral deficiencies; Vitamin Deficiency; Calcium; Iron; Niacin; Thiamine; Nutritional state; Nutritional adequacy; Urban areas

Abstract: A study of the nutritional status of 136 free-living, healthy elderly subjects (66 women, 70 men; ages 65-69) residing in an urban area revealed low calcium, iron, niacin, and thiamin intakes in 10-50% of the subjects and excess body fat in 85% of the women and in half of the men. Their diets also were slightly below recommended levels for carbohydrate and fat, but were high in protein. Food habits and differences in mineral deficiency incidence between sexes also are discussed.(wz)

200 NAL Call No: RA784.A1C3
Nutritional status, socio-economic environment and the lifestyle of the elderly in August Town, Kingston, Jamaica, 1984.

Mesfin, E.; Sinha, D.P.; Jutsum, P.J., Simmons, W.K.; Eldemire, D.

Kingston : Caribbean Food and Nutrition Institute, 1987.

Cajanus v. 20 (1): p. 23-47. charts; 1987. Includes 8 references.

Language: English

Descriptors: Jamaica; Elderly nutrition; Nutritional state; Anthropometric dimensions; Blood pressure, Health; Life style

Abstract: A survey assessed the nutritional status, socioeconomic environment, and lifestyle of 154 of 197 elderly residing in a small urban area of Kingston, Jamaica. Extensive data tabulations are provided on: age and sex distribution; sources of income; illnesses and illness patterns; anthropometric measurements; percentage of standard weight for age by sex; iron nutritional status; and nutritional status (judged by weight/height

ratio) vs. income sources and vs. high diastolic and systolic blood pressure, arthritis, and heart disease prevalence. The study design and methodology are detailed, and the survey results are discussed. Specific and general recommendations derived from the survey are included.(wz)

201 NAL Call No: RM222.5.N87
Nutritional support in long-term care facilities.

Sullivan, D.; Chernoff, R.; Lipschitz, D.A.

Baltimore, Md. : Williams & Wilkins ;, 1987 Feb. Nutrition in clinical practice : official publication of the American Society for Parenteral and Enteral Nutrition v. 2 (1): p. 6-13; 1987 Feb. Includes 45 references.

Language: English

Descriptors: Diet; Nutritional adequacy; Nursing homes; Elderly nutrition; Patient care; Guidelines

Abstract: An overview article examines and discusses various aspects of the provision of nutritional support in long-term care facilities. Attention is given to: the role of nursing homes in controlling patient malnutrition and its consequences; and factors involved in ensuring that patients in long-term care facilities receive adequate nutritional support (including mechanisms for nutritional status monitoring, identifying patients at nutritional risk, identifying adequate nutritional goals, and for delivering nutritional support). The inadequacy of professional staff caused by cost considerations is addressed as a key problem area that needs resolution.(wz)

202 NAL Call No: RA784.N8
Nutritional surveillance of the elderly: methods to determine program impact and unmet need.

Roe, D.A.

Baltimore, Md. : Williams & Wilkins; 1989 Oct. Nutrition today v. 24 (5): p. 24-29; 1989 Oct. Includes 9 references.

Language: English

Descriptors: Nutritional surveillance; Nutrition programs; Nutritional assessment; Federal programs; Public agencies; Home delivered meals; Elderly

Abstract: Nutritional surveillance of the elderly provides ongoing evidence that nutrition programs, both federal and state, can help to reduce health costs for those elderly whose care is otherwise so costly to the taxpayer. This article examines the development of one such program in New York State.

Nutrition and the Elderly

203 NAL Call No: TX341.D4
Nutrition-related problems in the institutionalized elderly.
Welch, T.
Columbus, Ohio : Ross Laboratories; 1989.
Dietetic currents v. 16 (1): p. 1-4; 1989. Includes 33 references.

Language: English

Descriptors: Elderly nutrition; Physiological functions; Weight; Albumins; Protein energy malnutrition; Iron deficiency anemia; Digestive disorders; Nursing homes

Abstract: The physical and biochemical changes which occur with aging create special nutritional problems for the geriatric patient. Health care professionals need to recognize the importance of nutrition in maintaining and improving the health of the elderly patient.

204 NAL Call No: RA784.N8
omega-3 fatty acids in growth and development and in health and disease. I. The role of omega-3 fatty acids in growth and development.
Simopoulos, A.P.
Baltimore, Md. : Williams & Wilkins; 1988 Apr.
Nutrition today v. 23 (2): p. 10-19. ill., charts; 1988 Apr. Includes 23 references.

Language: English

Descriptors: Polyunsaturated fatty acids; Nutrient requirements; Fish oils; Health promotion; Disease prevention; Lipid metabolism; Prostaglandins

Abstract: The first of a 2-part technical report summarizes and discusses the current scientific understanding of the role of fish and vegetable oil omega-3 fatty acids in growth and development. Specifically, the report addresses the sources and metabolism of these fatty acids, the elongation and desaturation of alpha-linolenic acid, and the role of these acids in pregnancy and fetal development, the nutrition of breast-fed infants, child health, and aging. The essentiality of these omega-3 fatty acids is discussed. The results of animal and human studies are included.

205 NAL Call No: 389.8 SCH6
Optimism abounds at New York's LAC.
Truax, M.
Denver, Colo. : American School Food Service Association; 1987 Jun.
School foodservice journal v. 41 (6): p. 50-51. ill, 1987 Jun.

Language: English

Descriptors: New York; School food service; Legislation; Conferences

Abstract: The New York State School Food Service Association held its 14th Legislative Action Conference in March 1987. Conference speakers included Stanley Lundine, New York's Lieutenant Governor; John Raftery, director of Child Nutrition Programs for Massachusetts; Reverend Kenneth Dean, a 1960s civil rights activist; Patricia Ayres, chairman of the Committee to Promote New York Agriculture; and Catherine Tillman, chief of the Bureau of Government Donated Foods for New York. Members of the Public Policy and Legislation Committee also presented the 1987 Legislative Issue Paper, which focused on the following topics: 1) general assistance support, 2) the non-nutritious food bill, 3) training program assistance, 4) the competitive foods bill, 5) nutrition education, 6) equipment assistance, 7) the access-to-lunch mandate, and 8) nutrition programs for the elderly.

206 NAL Call No: 389.8 J824
Osteopenia in the men of a Veterans Administration nursing home.
Nagraj, H.S.; Gergans, G.A.; Mattson, D.E.; Rudman, I.W.; Rudman, D.
Baltimore, Md. : American Society for Clinical Nutrition; 1990 Jan.
American journal of clinical nutrition v. 51 (1): p. 100-106. charts; 1990 Jan. Includes 32 references.

Language: English

Descriptors: Bone density; Bone fractures; Incidence; Osteoporosis; Body weight; Nursing homes; Elderly; Men

Abstract: We measured by photon absorptiometry the bone density at six sites in 65 nursing home men aged 57-58 y and in 25 independent community men aged 57-80 y. Average bone density in the community men ranged from 97% to 105% of age-matched normal men. In the nursing home men these values ranged from 71% to 92% of age-matched normal men (p less than 0.05 for comparison with the community men). About 50% of the nursing home men but none of the community men had a value less than 70% of age-matched normal men at one or more sites. Among the institutionalized men bone densities at all six sites (in g/cm²) were significantly (p less than 0.05) and directly correlated with body weight but were not significantly correlated with height, age, principal or

Quick Bibliography Series

secondary diagnoses, continuing medications, or functional level

207 NAL Call No: RC931.O73O76
Osteoporosis a patient education sampler.
United States, Food and Drug Administration
National Conference on Women's Health Series
1987 : United States. Food and Drug Administration
Washington : D.C. : U.S. Dept. of Health and
Human Services, Public Health Service, Food and
Drug Administration; 1987.
[24] p. : ill. ; 28 cm. Cover title. At head of title:
A Special Topic Conference. National Conference
on Women's Health Series; sponsored by the Food
and Drug Administration, October 30, 1987, Bet-
hesda, Maryland. Includes bibliographical refer-
ences.

Language: English

Descriptors: Osteoporosis--Prevention, Women's
health services; Teaching materials; Osteoporosis;
Patient education; Women; Study guides; Disease
prevention; Diagnosis; Guidelines

Abstract: An educational document for health care
providers, patients, and women in general provides
resource information for becoming more informed
and active in osteoporosis prevention and treat-
ment. The document includes articles from the
FDA Consumer publication addressing key health
messages for women in their teens and twenties;
middle-aged women, and older women. The arti-
cles include: osteoporosis, a women's guide; os-
teoporosis: cause, treatment, prevention;
osteoporosis, calcium and estrogens; and the
menopause time of life. An article encouraging ex-
ercising by the National Institute on Aging, sugges-
tions for further reading, and space for patients to
enter questions to ask their doctor also are in-
cluded. The pages of the document are perforated
for convenient duplication and sharing of the ma-
terials.

208 NAL Call No: TX341.B75
Osteoporosis and calcium supplementation.
Bailey, A.J
London : The Foundation; 1989 May.
BNF Nutrition bulletin - British Nutrition Founda-
tion v. 14 (2): p. 126-134. ill; 1989 May. Includes 5
references.

Language: English

Descriptors: Osteoporosis; Calcium; Supplements;
Bone density; Collagen; Elderly

Abstract: The use of calcium supplements for the
prevention of osteoporosis or for regeneration of
bone is controversial. This article argues that cal-
cium deficiency is rare and that bone strength does
not depend solely on mineralization. It is proposed
that the quality of the collagenous framework is
important to bone strength because replacement
of the collagen matrix framework by hormone
therapy results in an increase in bone thickness and
a reduction in the incidence of fractures.

209 NAL Call No: RC931.O73R6 1986
Osteoporosis current concepts : report of the
Seventh Ross Conference on Medical Research.
Roche, Alex F., 1921-; Gussler, Judith D.
Ross Laboratories
Ross Conference on Medical Research 7th : 1986
: Charleston, S.C.
Columbus, Ohio : Ross Laboratories; 1987.
vii, 97 p. : ill. ; 26 cm. "Held at Charleston, South
Carolina, April 23-25, 1986"--P. iii. Includes
bibliographies.

Language: English

Descriptors: Osteoporosis--Congresses; Reference
works; Osteoporosis; Therapy; Calcium; Vitamin
D; Metabolism; Disease prevention; Patient care
Abstract: The proceedings of a 1986 medical re-
search conference for physicians and scientists
working on various aspects of osteoporosis
presents 31 papers addressing aspects of osteoporosis
studies and issues concerning this disease. Topics
include: the epidemiology of osteoporosis etiology;
skeletal status in children, adolescents, and the el-
derly; hormonal and calcium metabolism; risks fac-
tors; hormonal and calcitonin therapy; bone
metabolism and bone mass; cellular abnormalities;
bone loss prevention; bone mass measurements;
general osteoporosis treatment; and ethical issues
and fraud. Discussions are appended to each of the
papers.

210 NAL Call No: TX361.A3J6
The PaCS survey--a renewed commitment to good
care.
Matthews, I E.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (4): p. 69-
73; 1987. Includes references.

Language: English

Descriptors: USA; Nursing homes; Patient care; El-
derly nutrition; Surveys; Facilities; Medicaid;
Medicare

Nutrition and the Elderly

Abstract: A brief report discusses the characteristics and requirements of the new Patient Care and Services (PaCS) survey that all nursing homes for elderly patients participating in the US federal Medicare/Medicaid program must experience by the end of 1987 as part of their annual state licensure inspection. The 3 major components of the PaCS survey format focus on: (1) new facilities, emphasizing traditional concerns (facility structure, staffing, written policies/procedures, etc.); (2) a comprehensive facilities inspection, including the dining area and quality of eating assistance provided; and (3) fire and safety compliance. Detailed discussion is given to elderly nutritional assessment and satisfying nutritional needs. It is noted that the PaCS survey system places an increased emphasis on aggressive intervention rather than passive care of elderly subjects.

211 NAL Call No: TX361.A3J6
Peer support and nutrition education for older adults with diabetes.
Pratt, C.; Wilson, W.; Leklem, J.; Kingsley, L.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (4): p. 31-43. charts; 1987. Includes 22 references.

Language: English

Descriptors: Diabetes; Elderly; Nutrition education; Weight reduction; Food habits; Psychological factors; Peer relationships; Support systems; Patient compliance; Beliefs; Dietary surveys
Abstract: Extract: The impact of nutrition education, peer support and several pre-intervention psycho-social variables upon weight change and glycemic control were assessed in older, noninsulin-dependent diabetic patients. Significant weight reductions were observed in persons who participated in the combined nutrition education and peer support program. No significant changes were observed in glycemic control. Several psycho-social variables were significantly correlated with reductions in weight and improvement in glycemic control. Implications for future research and intervention programs for older persons with nutrition related diseases are presented. (author)

212 NAL Call No: RM214.T66
Perceived relationships between taste and smell acuity and food intake in the elderly.
Jurdi-Haldeman, D.; Napier, A.K.
Gaithersburg, Md. : Aspen Publishers; 1988 Oct.
Topics in clinical nutrition v. 3 (4): p. 4-8; 1988 Oct. Includes 38 references.

Language: English

Descriptors: Food intake; Flavors; Taste; Odors; Elderly; Nutritional intervention

Abstract: The improvement in the nutritional status of the elderly patient is discussed in terms of improving the appetite, resulting in increased food intake. Dietitians are encouraged to make necessary modification in food preparation and presentation to improve food palatability and flavor for the elderly.

213 NAL Call No: TX361.A3J6
Perceptions of elderly residents--geriatric not pediatric.
Matthews, L.E.
Binghamton, N.Y. : Haworth Press; 1987.
Journal of nutrition for the elderly v. 6 (3): p. 57-60; 1987.

Language: English

Descriptors: Institutional food service
Abstract: A technical report stresses that the nutritional and food needs of institutional elderly subjects residing in health care facilities should be considered in terms of their needs as elderly subjects, rather than "infanticizing" such subjects, treating them as infants or young children. Practical guidelines for long term professionals and staff treating such elderly are included on appropriate menu selections, texture modifications, and for treating elderly as respected adults in resident interviews.(wz)

214 NAL Call No: RM214.T66
Perspectives on nutrition care for the geriatric hospice patient.
Weddle, D.O.; Prewitt, T.E.; Gordon, A.K.
Gaithersburg, Md. : Aspen Publishers; 1988 Oct.
Topics in clinical nutrition v. 3 (4): p. 20-26; 1988 Oct. Includes 15 references.

Language: English

Descriptors: Nutritional support, Elderly; Hospices; Diet counseling
Abstract: This article provides general information about hospice care for the geriatric patient as well as information about the current role of nutrition and dietetic services in hospices. The implications for nutrition services in future hospice environment also discussed.

215 NAL Call No: 389.8 J824
Phylloquinone in plasma from elderly and young

Quick Bibliography Series

adults: factors influencing its concentration.
Sadowski, J.A.; Hood, S.J.; Dallal, G.E.; Garry, P.J.
Baltimore, Md : American Society for Clinical Nutrition; 1989 Jul.
American journal of clinical nutrition v. 50 (1): p. 100-108 charts; 1989 Jul. Includes 40 references.

Language: English

Descriptors: Vitamin K; Blood plasma; Triglycerides; Cholesterol; High density lipoprotein; Regression analysis; Young adults; Elderly

Abstract: Fasting plasma phylloquinone concentrations were determined by reverse-phase high-performance liquid chromatography in a population of young (n = 131) and elderly (n = 195) human subjects. The distribution of values was non-Gaussian but was converted to Gaussian form after a logarithmic transformation of the data. The normal range derived from the lognormal distribution was 0.29-2.64 nmol/L (geometric average = 0.87 nmol/L, median = 0.86 nmol/L). Young females had significantly decreased plasma phylloquinone concentrations when compared with the elderly females whereas concentrations in young males were slightly higher than those in elderly males. A regression model was constructed and revealed that plasma phylloquinone concentrations were positively correlated with plasma triglycerides and alpha-tocopherol. Although the elderly subjects (average = 1.05 nmol/L) had higher levels of phylloquinone than the young subjects (0.94 nmol/L), when the concentration of phylloquinone in plasma is expressed as nanomoles phylloquinone per millimole of triglyceride, the elderly subjects (0.62×10^{-6}) showed decreased levels for phylloquinone compared with the young subjects (0.82×10^{-6}) (p less than 0.001).

216 NAL Call No: 389.8 Z33
Plasma and leucocyte zinc concentrations and their response to zinc supplementation in an elderly population.
Field, H.P.; Whitley, A.J.; Srinivasan, T.R.; Walker, B.E.; Kelleher, J.
Bern : Hans. Huber; 1987.
International journal for vitamin and nutrition research v. 57 (3): p. 311-317. charts; 1987. Includes 24 references.

Language: English

Descriptors: Zinc; Elderly nutrition; Blood plasma; Leukocytes; Mineral supplements

Abstract: A study of plasma and leucocyte zinc (Zn) levels relative to different levels of Zn supplementation (50-150 mg/day) in 15 institutionalized elderly women revealed that plasma Zn levels rose while granulocyte Zn levels declined. Mononuclear cell Zn levels were unaffected by Zn supplementation. The clinical implications of these results are discussed.

217 NAL Call No: P 620.A1N8
Plasma calcium and trace metals in human subjects with mature cataract.

Bhat, K.S.

Stoneham, Mass. : Butterworth Publishers; 1988 Jan.

Nutrition reports international v. 37 (1): p. 157-163. ill., charts; 1988 Jan. Includes 23 references.

Language: English

Descriptors: India; Men; Women; Cataract; Blood plasma; Calcium; Magnesium; Chromium, Copper; Zinc; Nutritional state

Abstract: Extract: Plasma samples from 39 human subjects with mature cataract and 15 subjects without the eye disease were analysed for calcium, magnesium, chromium, copper and zinc by atomic absorption spectrometry. Copper and zinc levels in plasma were significantly lower in subjects with cataract as compared to control subjects. The concentrations of calcium, magnesium and chromium in plasma from subject in the former group were similar to those from the latter group. Age appeared to have no effect on plasma copper, but zinc levels were significantly lowered in elderly persons with cataract as compared to younger subjects with the eye disease. The results suggest an association between cataract and nutritional status of copper and zinc in human subjects. (author)

218 NAL Call No: J89.8 J824
Plasma pyridoxal 5'-phosphate concentration and dietary vitamin B-6 intake in free-living, low-income elderly people.

Manore, M.M.; Vaughan, L.A.; Carroll, S.S.; Leklen, J.E.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Aug.

American journal of clinical nutrition v. 50 (2): p. 339-345. charts; 1989 Aug. Includes 32 references.

Language: English

Descriptors: Pyridoxine; Nutrient intake; Blood plasma; Vitamin supplements; Low income groups;

Nutrition and the Elderly

Elderly

Abstract: Free living, elderly persons (aged greater than or equal to 60 y, n = 198) were recruited to determine the effects of age, sex, health status, dietary vitamin B-6 intakes, and B-6 supplement use on plasma pyridoxal 5'-phosphate (PLP). Vitamin B-6 intakes were determined from 3-d diet records; supplementation was based on self-reported brand and frequency data. Fasting blood samples were analyzed for PLP. Subjects were primarily low-income Caucasians. There was no linear relationship between dietary vitamin B-6 intake, age, sex or health status, and PLP while accounting for supplemental vitamin B-6 use. PLP, however, was negatively correlated with age (p less than 0.001) in individuals with PLP values between 32 and 90 nmol/L. Vitamin B-6 status was low (PLP less than nmol/L) in 32% of this elderly population (n = 198) and could be attributed to low dietary vitamin B-6 intakes and/or the presence of health problems reported to alter vitamin B-6 status. This research suggests that low vitamin B-6 status is prevalent in low-income, elderly persons, especially those with multiple health problems.

219 NAL Call No: 389.8 AM34
Pleasantness perception of salt in young vs. elderly adults.
Chauhan, J.
Chicago, Ill. : The Association; 1989 Jun.
Journal of the American Dietetic Association v. 89 (6): p. 834-835. charts; 1989 Jun. Includes 9 references.

Language: English

Descriptors: Taste sensitivity, Sodium chloride, Age; Sex differences; Young adults; Elderly
Abstract: This study examined age effects on suprathreshold taste intensity and pleasantness perception of salt (NaCl) in both aqueous and food systems using large population samples in three discrete age groups, the young (20-29 yrs), the old (70-79 yrs) and the very old (80-99) yrs.

220 NAL Call No: 389.8 N953
Practical nutrition for the elderly.
Davies, L.
Washington, D.C. : Nutrition Foundation; 1988 Feb.
Nutrition reviews v. 46 (2): p. 83-87. ill., charts; 1988 Feb Includes 7 references.

Language: English

Descriptors: Elderly nutrition, Malnutrition, Diag-

nosis; Patient care, Nutrition education; Guidelines

Abstract: A brief report discusses practical approaches to the diagnosis of malnutrition in the elderly and provides suggestions for enhancing the understanding of nutrition by the elderly and by health professionals dealing with the elderly. Specific attention is given to myths and realities concerning elderly diets; and the general classification of malnutrition in the elderly and easily recognizable warning signs of malnutrition. Practical examples of nutrition teaching for the elderly and for physicians and nutritionists caring for the elderly are included.

221 NAL Call No: TX361.A3J63
Predicting dependent feeding behaviors in the institutionalized elderly.
Nolen, N.R.; Garrard, J.
Binghamton, N.Y. : Haworth Press, 1988
Journal of nutrition for the elderly v. 7 (3) p. 17-25. charts; 1988. Includes 12 references

Language: English

Descriptors: Eating habits; Food consumption; Liquids; Self feeding; Prediction; Nursing homes; Patient care; Elderly

Abstract: Dependence in feeding behaviors in the institutionalized elderly results in increased staff costs and loss of self-esteem. This study analyzed the conditions which appeared to be related to the loss of independent feeding skills in 106 individuals in a skilled care setting. At highest risk for loss of self-feeding skills are passive, preoccupied individuals and those with cognitive disorders. Staff has a tendency to feed people when intake begins to fall. Consistency in seating and routine, as well as creative alternatives between total independence and dependence are needed in order to maximize self-feeding behaviors

222 NAL Call No: 389.8 AM34
Prediction of body weight for the nonambulatory elderly from anthropometry.
Chumlea, W.C.; Guo, S.; Roche, A.F.; Steinbaugh, M.L.
Chicago, Ill. : The Association; 1988 May.
Journal of the American Dietetic Association v. 88 (5): p. 564-568. charts; 1988 May. Includes 25 references.

Language: English

Descriptors: Body weight, Elderly; Prediction, Anthropometric dimensions, Nutritional state

Quick Bibliography Series

Abstract: Extract: Body weight is one of the most important measurements in assessing nutritional status. Weight is also an important variable in equations that are needed to predict caloric expenditure and in indexes of body composition. Equations to predict the body weight of nonambulatory elderly persons with known errors should make use of body measurements that are indexes of the actual constituents of body composition and that can be collected regardless of an elderly person's level of mobility. The study samples consisted of a validation sample of 228 ambulatory white elderly persons, a cross-validation sample, and a clinical-validation sample of nonambulatory white elderly persons. Equations to predict body weight in the validation sample were derived from recumbent measures of arm and calf circumferences, triceps and subscapular skinfolds, and knee height. The accuracy, sensitivity, and validity of the equations were independently tested in the cross-validation sample; mean signed differences in actual and predicted weight ranged from differences in actual and predicted weight ranged from 0.1 to 1.8 kg. In the clinical-validation sample, the mean differences were larger. Prediction equations are presented using two, three, and four body measurements to allow the selection of an equation based upon those measurements that are possible to collect from a nonambulatory person on an individual basis.

223 NAL Call No: 389.8 J824
Predictive ability of various nutritional variables for mortality in elderly people.
Agarwal, N.; Acevedo, F.; Leighton, L.S.; Cayten, C.G.; Pitchumoni, C.S.
Baltimore, Md. : American Society for Clinical Nutrition; 1988 Nov.
American journal of clinical nutrition v. 48 (5): p. 1173-1178. charts; 1988 Nov. Includes 31 references.

Language: English

Descriptors: Nutritional assessment; Prediction; Malnutrition; Diagnosis; Blood serum protein; Albumins; Body weight; Mortality

Abstract: Nutritional indices (percentage ideal body weight [IBW], serum albumin, serum transferrin, total lymphocyte count [TLC] and delayed cutaneous hypersensitivity [DH] response) were assessed in 80 consecutive patients (aged 85-100 y) within 24 h of admission to determine their predictive value for mortality. Nine patients died. Pearson correlation analysis demonstrated that death

was significantly (p less than 0.05 to less than 0.01) associated with sepsis, serum albumin less than 30 g/L, TLC less than or equal to 1500 cells/mm³, and percentage IBW less than or equal to 90%. However, when serum albumin was controlled for, logit regression analyses demonstrated that the impact of other nutritional indices on death was insignificant. The effect of serum albumin remained significant (p less than 0.05 to less than 0.01) even when age and physician's diagnosis were held constant. With the logit model, serum albumin greater than or equal to 30 g/L had a sensitivity of 0.33, specificity of 0.99, and overall predictive power of 0.91. Serum albumin is thus the simplest and best single predictor of mortality and can provide early identification of elderly people at increased risk of death.

224 NAL Call No: TX361.A3J63
Predictors of nutrition supplement use in the elderly part II: the role of beliefs, attitude, subjective norm and intention.
Cotugna, N.
Binghamton, N.Y. : Haworth Press; 1989.
Journal of nutrition for the elderly v. 8 (3,4): p. 15-33. charts; 1989. Includes 17 references.

Language: English

Descriptors: Consumer attitudes; Consumer behavior; Food beliefs; Prediction; Elderly; Women; Men

Abstract: This study was able to demonstrate the predictive role of beliefs in determining the intention to use nutrient supplements in an elderly population. It provided insight into the beliefs which formulated positive attitudes toward nutrition supplement use, and the relevant weights of those beliefs.

225 NAL Call No: TX361.A3J63
Predictors of nutrition supplement use in the elderly part I: A review of the literature.
Cotugna, N.
Binghamton, N.Y. : Haworth Press; 1989.
Journal of nutrition for the elderly v. 8 (3,4): p. 1-1989. Literature review. Includes 41 references.

Language: English

Descriptors: Vitamin supplements; Mineral supplements; Consumer behavior; Consumer surveys; Literature reviews; Elderly; Women; Men

Abstract: The use of nutrition supplements is a common health behavior in Americans especially

Nutrition and the Elderly

among the elderly. The author studied reasons as to why the elderly are so vulnerable to the use of nutrition supplements. Level of education, age differentiation, and socioeconomic status do not have an impact on supplement use by the elderly, although they do in the general population. Race or ethnic background differences have not been adequately studied. The one factor which does appear to be significant is gender, with women being more frequent users of nutrition supplements than men.

226

NAL Call No: 448.9 AM37

The prevalence of high blood cholesterol levels among adults in the United States.

Sempos, C.; Fulwood, R.; Haines, C.; Carroll, M.; Anda, R.; Williamson, D.F.; Remington, P.; Cleeman, J.

Chicago, Ill. : The Association; 1989 Jul07.

JAMA : Journal of the American Medical Association v. 262 (1) p. 45-52 charts; 1989 Jul07. Includes 49 references.

Language: English

Descriptors: USA; Cholesterol, Surveys; Blood lipids, Risks; Cardiovascular diseases, Guidelines, Patient education, Nutritional intervention; Men, Women

Abstract: Using the National Cholesterol Education Program's Guidelines for the Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults and the most recent nationally representative serum total cholesterol and lipoprotein data for adults (ages 20 to 74 years) from the second National Health and Nutrition Examination Survey (1976 to 1980), it is estimated that 41% of adults should have lipoprotein analysis after an initial measurement of serum total cholesterol. Furthermore, it is estimated that 88% of those who need lipoprotein analysis, or 36% of all adults aged 20 to 74 years, are candidates for medical advice and intervention for high blood cholesterol levels. We estimate, based on 1986 population data, that approximately 40 million Americans between the ages of 20 and 59 years are candidates for medical advice and intervention. An additional 24 million Americans aged 60 years and older are candidates. Overall, we estimate that about 60 million Americans aged 20 years and older are candidates for medical advice and intervention for high levels of blood cholesterol, although a less intensive approach might be appropriate for elderly patients.

227

NAL Call No: 151.65 P96

Prospective study of pneumonia hospitalizations and mortality of U.S. older people: the role of chronic conditions, health behaviors, and nutritional status.

LaCroix, A.Z.; Lipson, S.; Miles, T.P.; White, L. Washington, D.C. : Public Health Service; 1989 Jul. Public health reports v. 104 (4): p. 350-360. charts, 1989 Jul. Includes 43 references.

Language: English

Descriptors: Pneumonia; Morbidity; Mortality; Nutritional state; Risks; National surveys; Men; Women; Elderly

Abstract: Mortality and hospitalization rates for pneumonia have increased among older Americans during recent years (1979-86), despite a national commitment to the reduction of premature deaths from pneumonia. A prospective study of deaths and hospitalizations attributable to pneumonia was conducted among 5,474 subjects ages 55 and older who participated in the NHANES I Epidemiologic Followup Study. Prevalent chronic conditions, health behaviors, and nutritional status indicators, measured at baseline, were examined in relation to pneumonia hospitalization and death during 12 years of followup. Mortality and hospitalization rates for pneumonia were higher among men than women, and higher among those ages 65 and older than among those 55-64 of both sexes. Risk of pneumonia death was higher among subjects with a history of congestive heart failure, stroke, cancer, or diabetes. Risk of pneumonia hospitalization was higher among subjects with a history of chronic obstructive pulmonary disease and among men who were current smokers. Daily alcohol consumption did not increase risk of pneumonia in this study population. Four measures of nutritional status were examined taking age, prevalent chronic conditions, and cigarette smoking into account. body mass index, arm muscle area, and serum albumin and hemoglobin levels. Risk of pneumonia death was 2.6 times higher in men in the lowest quartile, compared with men in the highest quartile, of body mass index. Similarly, the risk was 4.5 times higher among men in the lowest quartile of arm muscle area. Risk of death from pneumonia was 3.6 times higher among women in the lowest quartile of serum albumin levels compared with women in the highest quartile. Relative risks for these nutritional status indicators remained elevated after adjusting for age and the medical history risk factors.

228

NAL Call No: RC620.A1N8

Quick Bibliography Series

Protein calorie malnutrition in elderly chronic care institutions in Hong Kong.

Woo, J.; Ho, S.C.; Cheung, C.K.; Mak, Y.T.; Swaminathan, R.

Stoneham, Mass. : Butterworth; 1989 Nov.

Nutrition reports international v. 40 (5): p. 1011-1018. charts; 1989 Nov. Includes 18 references.

Language: English

Descriptors: Hong kong; Protein energy malnutrition; Nutritional state; Protein intake; Caloric intake; Anthropometric dimensions; Long term care; Elderly

Abstract: Protein nutritional status was assessed in 208 elderly residents of four long term care institutions in Hong Kong, by measuring protein and energy intake, anthropometric indices, and concentrations of plasma proteins. Compared to free-living elderly subjects of a similar age group, institutionalized subjects had lower daily protein and protein calorie intake in spite of similar intake total energy, lower anthropometric measurements and lower plasma protein concentrations. Subjects who were unable to feed themselves and those who died within a three month period had poorer nutritional status. This survey shows evidence of protein calorie malnutrition among elderly residents of long term care institutions in Hong Kong, and that it may contribute to short term mortality.

229

NAL Call No: 389.8 J824

Protein nutriture of a group of free-living elderly.

Munro, H.N.; McGandy, R.B.; Hartz, S.C.; Russell, R.M.; Jacob, R.A.; Otradovec, C.L.

Baltimore, Md. : American Society for Clinical Nutrition; 1987 Oct.

American journal of clinical nutrition v. 46 (4): p. 586-592. charts; 1987 Oct. Includes 25 references.

Language: English

Descriptors: Protein intake; Energy intake; Pyridoxine; Muscle tissue; Skin fold thickness; Elderly

Abstract: The adequacy of the protein intakes of elderly people without overt debilitating diseases was investigated on 691 free-living men and women divided into those aged 60-75 y and those greater than 75 y. In both age groups men and women had average protein intakes of 1.02-1.06 g/kg body weight, values well above the safe level of 0.75 g/kg recommended in a WHO/FAO/UNU report. Although plasma concentrations of albumin, prealbumin, and transferrin declined with age, these were not related to low intakes of protein by individual elderly people. Similarly, upper-arm muscle mass was not less in those elderly people at the lower end of the range of protein intakes. Thus in this population of overtly healthy elderly men and women, there was no evidence of protein deficiency in contrast to other surveys where elderly people with chronic diseases were included. (author)

vidual elderly people. Similarly, upper-arm muscle mass was not less in those elderly people at the lower end of the range of protein intakes. Thus in this population of overtly healthy elderly men and women, there was no evidence of protein deficiency in contrast to other surveys where elderly people with chronic diseases were included.

230

NAL Call No: 389.8 J824

Protein nutriture of a group of free-living elderly.

Munro, H.N.; McGandy, R.B.; Hartz, S.C.; Russell, R.M.; Jacob, R.A.; Otradovec, C.L.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Oct.

American journal of clinical nutrition v. 46 (4): p. 586-592. charts; 1987 Oct. Includes 25 references.

Language: English

Descriptors: Elderly nutrition; Protein intake; Nutritional state; Blood composition; Anthropometric dimensions; Pyridoxine

Abstract: Extract: The adequacy of the protein intakes of elderly people without overt debilitating diseases was investigated on 691 free-living men and women divided into those aged 60-75 y and those greater than 75 y. In both age groups men and women had average protein intakes of 1.02-1.06 g/kg body weight, values well above the safe level of 0.75 g/kg recommended in a WHO/FAO/UNU report. Although plasma concentrations of albumin, prealbumin, and transferrin declined with age, these were not related to low intakes of protein by individual elderly people. Similarly, upper-arm muscle mass was not less in those elderly people at the lower end of the range of protein intakes. Thus in this population of overtly healthy elderly men and women, there was no evidence of protein deficiency in contrast to other surveys where elderly people with chronic diseases were included. (author)

231

NAL Call No: TX361.A3J63

The provision of weekend home delivered meals by state and a pilot study indicating the need for weekend home delivered meals.

Walden, O.; Hayes, P.A.; Lee, D.Y.; Montgomery, D.H.

Binghamton, N.Y. : Haworth Press; 1988.

Journal of nutrition for the elderly v. 8 (1). p. 31-43. charts; 1988. Includes 16 references.

Language: English

Descriptors: USA; Home delivered meals; Workable days; Program evaluation; Nutrient intake;

Nutrition and the Elderly

Surveys; Elderly

Abstract: A survey of state units on aging revealed that 43 states and the District of Columbia provide some weekend home delivered meals. In most states meals were provided in only a few areas. There is no national system for reporting weekend delivery. Very few states have data on the extent of weekend meal service. A pilot study to determine the need for home delivered meals indicated that persons receiving home delivered meals 5 days a week were more likely to have an insufficient dietary intake of protein, thiamin, riboflavin, calcium, iron, and phosphorus on weekends than weekdays.

232

NAL Call No: 389.8 J824

Pyridoxine supplementation: effect on lymphocyte responses in elderly persons.

Talbott, M.C.; Miller, L.T.; Kerkvliet, N.I.
Baltimore, Md. : American Society for Clinical Nutrition; 1987 Oct.

American journal of clinical nutrition v. 46 (4): p. 659-664. charts; 1987 Oct. Includes 21 references.

Language: English

Descriptors. Pyridoxine, Vitamin supplements, Lymphocytes; Immune response, Elderly

Abstract. The effect of pyridoxine supplementation on lymphocyte responsiveness was investigated in 15 persons aged 65-81 y. Eleven subjects received 50 mg/d pyridoxine HCl (PN). Four subjects received a placebo. Lymphocyte proliferation to T and B cell mitogens, lymphocyte subpopulations with monoclonal antibodies, and plasma pyridoxal 5'-phosphate (PLP) were measured before and after 1 and 2 mo of supplementation. After 1 and 2 mo plasma PLP levels increased by 195 +/- 88 nM and 201 +/- 84 nM, respectively, in subjects receiving PN. With PN supplementation, lymphocyte proliferation increased significantly in response to phytohemagglutinin (p less than 0.01), pokeweed mitogen (p less than 0.01), and Staphylococcus aureus (Cowain I) (p less than 0.05). For PN-treated subjects with low presupplement plasma PLP levels, lymphocyte blastogenesis also increased significantly (p less than 0.01) in response to concanavalin A. Percentages of T3+ and T4+ but not T8+ cells increased significantly (p less than 0.05) in PN-treated subjects. These results suggest that improving vitamin B-6 status is important in stimulating immunocompetence in the elderly.

233

NAL Call No: 389.8 J824

Pyridoxine supplementation: effect on lymphocyte responses in elderly persons.

Talbott, M.C.; Miller, L.T.; Kerkvliet, N.I.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Oct.

American journal of clinical nutrition v. 46 (4) p. 659-664. ill., charts; 1987 Oct. Includes 21 references.

Language: English

Descriptors: Pyridoxine; Pyridoxal phosphate; Vitamin supplements; Lymphocytes, Elderly nutrition

Abstract: Extract. The effect of pyridoxine supplementation on lymphocyte responsiveness was investigated in 15 persons aged 65-81 y. Eleven subjects received 50 mg/d pyridoxine HCl (PN). Four subjects received a placebo. Lymphocyte proliferation to T and B cell mitogens, lymphocyte subpopulations with monoclonal antibodies, and plasma pyridoxal 5'-phosphate (PLP) were measured before and after 1 and 2 mo of supplementation. After 1 and 2 mo plasma PLP levels increased by 195 +/- 88 nM and 201 +/- nM, respectively, in subjects receiving PN. With PN supplementation, lymphocyte proliferation increased significantly in response to phytohemagglutinin (p less than 0.01), pokeweed mitogen (p less than 0.01), and Staphylococcus aureus (Cowain I) (p less than 0.05). For PN-treated subjects with low presupplement plasma PLP levels, lymphocyte blastogenesis also increased significantly (p less than 0.01) in response to concanavalin A. Percentages of T3+ and T4+ but not T8+ cells increased significantly (p less than 0.05) in PN-treated subjects. These results suggest that improving vitamin B-6 status is important in stimulating immuno-competence in the elderly.(author)

234

NAL Call No: TX341.C6

RDA study for vitamin A defines need, eyes risks.
Washington, D.C. : Community Nutrition Institute, 1987 May21.

Nutrition week v. 17 (21): p. 4-5. charts; 1987 May21.

Language: English

Descriptors. USA, Vitamin A, Nutrient requirements; Recommended dietary allowances; Food and nutrition controversies, Vitamin supplements

Abstract: A brief technical report provides an excerpt of the recommended dietary intake (RDI) for vitamin A proposed for the 1985 10th edition of the Recommended Dietary Allowances report rejected by the US National Academy of Sciences.

Quick Bibliography Series

Information is provided on vitamin A dietary sources, intakes, essentiality, deficiency states, and the vitamin A nutritional status and requirements for infants, children, adults, pregnant and lactating women, and the elderly. Ancillary information is given on vitamin A nutrient-nutrient interactions and vitamin A toxicity at high ingestion doses. The clinical rationale for lowering the vitamin A RDA for infants and adults also is discussed.(wz)

235 NAL Call No: aHD9001.N275
Recent trends in domestic food programs.

Matsumoto, M.

Washington, D.C. : The Service; 1987.

National food review NFR - United States Dept. of Agriculture, Economic Research Service (NFR-36): p. 24-25. charts; 1987.

Language: English

Descriptors: USA; Supplemental feeding programs; Federal programs; Child nutrition; Program participants; Cost benefit analysis

Abstract: A brief statistical report compares the participation in and costs of US domestic food programs between the 4th quarter of fiscal year 1986 (July-September) and the same period during 1985. Attention is focused on 3 categories of these programs, viz.: child nutrition programs (e.g., school lunch program); supplemental food programs (e.g., WIC program); and food distribution programs (e.g., nutrition programs for the elderly). Tabular data are included on the average participation in USDA food programs and on the benefit cost of the various USDA food programs.(wz)

236 NAL Call No: 389.8 J824
Recommended dietary intakes of elderly subjects.

Hegsted, D.M.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.

American journal of clinical nutrition v. 50 (5): p. 1190-1194; 1989 Nov. Includes 17 references.

Language: English

Descriptors: Nutrient requirements, Recommended dietary allowances; Individual characteristics; Elderly

Abstract: Various authors have criticized the Recommended Dietary Allowances (RDA) because they do not include values for elderly subjects. It is argued here that such values would not be useful. The large majority of elderly subjects have one or more diagnosed diseases, drug consumption is high, etc, and it is unlikely that specific dietary

standards for such a group could be developed or would be useful. More importantly, I argue that the RDA do not and cannot do most of the things which dietary standards ought to do, and their use generates considerable misinformation. It is time to reconsider the whole issue of dietary standards. It is clear that elderly subjects who do not require therapeutic diets need to consume a diet which differs from that of the rest of the population.

237 NAL Call No: 389.8 J824
Recommended dietary intakes (RDI) of vitamin C in humans.

Olson, J.A.; Hodges, R.E.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Apr.

American journal of clinical nutrition v. 45 (4): p. 693-703. charts; 1987 Apr. Literature review. Includes 100 references.

Language: English

Descriptors: Ascorbic acid; Nutrient requirements; Recommended dietary intakes; Absorption; Human nutrition research

Abstract: Extract: Vitamin C is unevenly distributed throughout all body cells and fluids. A total-body pool of 900 mg (5.11 mmol) in an adult male meets the following criteria of a satisfactory vitamin C status: 1) it is threefold higher than one that prevents scurvy, 2) no known health advantages accrue at higher pool sizes, 3) absorption efficiency is high and urinary loss low at appropriate intakes, and 4) a 1-mo reserve allows for periods of low intake or stress. To maintain a suitable body pool in healthy 76-kg men and 62-kg women requires daily intakes (RDI) of 40 mg (227 micromoles) and 30 mg (170 micromoles), respectively. Reasons for not increasing RDI values to enhance iron absorption and to prevent nitrosamine formation are given. The specific association of normal intakes of dietary vitamin C with cancer is very weak and not quantifiable. Advisable intakes for children, pregnant and lactating women, and the elderly are considered. The present recommendations are in better accord with current information than are the 1980 RDA values.(author)

238 NAL Call No: 389.8 J824
Recommended dietary intakes (RDI) of vitamin A in humans.

Olson, J.A.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Apr.

Nutrition and the Elderly

American journal of clinical nutrition v. 45 (4): p. 704-716. charts; 1987 Apr. Includes 80 references.

Language: English

Descriptors: Vitamin A; Nutrient requirements; Recommended dietary intakes; Human nutrition research

Abstract: Extract: Mean and median dietary intakes of vitamin A in the US are approximately 1000 and 624 retinol equivalents, respectively. In this paper a satisfactory vitamin A status is defined as a total-body pool that provides adequate vitamin A to meet all known physiological needs and a reserve for 4 mo on low intakes or during stress. Mean dietary intakes required to maintain an adequate body pool (ie, 20 micrograms vitamin A/g liver [0.07 micromoles/g]) in reference 76-kg men and 62-kg women are 506 and 413 micrograms retinol, respectively. Recommended dietary intakes (RDI) for nearly all reference men and women are 700 micrograms and 600 micrograms, respectively. Vitamin A needs in infants, children, the elderly, and pregnant and lactating women are quantified. The toxicity of vitamin A in early pregnancy, for which a safe intake level is not known, is stressed. On many grounds these suggested lower RDI values are preferable to the 1980 RDA values.(author)

239 NAL Call No: TX361.A3J63
The relationship between a nutrition education program and nutrition knowledge and eating behaviors of the elderly.
Bedell, B.A.; Shackleton, P.A
Binghamton, N.Y. : Haworth Press; 1989.
Journal of nutrition for the elderly v. 8 (3,4): p. 35-45; 1989. Includes 25 references.

Language: English

Descriptors: Nutrition knowledge; Nutrition information, Consumer education, Program evaluation
Abstract: An experimental study was conducted to examine the relationship between a nutrition education program and nutrition knowledge and eating behaviors of older adults. Twenty-four subjects were randomly assigned to an experimental or a control group. Prior to any intervention, all subjects completed a demographic data sheet, a 25-item True/False questionnaire and a 24-Hour Dietary Recall interview. The experimental group attended four one-hour nutrition education classes over a two-week period. Following the educational intervention, both groups of subjects again completed the True/False questionnaire and the 24-

Hour Dietary Recall interview. Controlling for pretest, posttest differences on the two dependent variables were tested using repeated measures analysis of variance. No significant differences were found between the two groups.

240 NAL Call No: 389.8 AM34
Relationship of socioeconomic status and living arrangements to nutritional intake of the older person.
Ryan, V.C.; Bower, M.E.
Chicago, Ill. : The Association; 1989 Dec.
Journal of the American Dietetic Association v. 89 (12): p. 1805-1807. charts; 1989 Dec. Includes 35 references.

Language: English

Descriptors: South Carolina; Nutrient intake; Nutritional surveys; Socioeconomic status; Living conditions; Nutritional assessment; Psychological factors; Elderly

Abstract: The number and proportion of older Americans have grown rapidly in recent years, and this trend is expected to continue. Adequate nutritional intake is essential for optimal physical and mental activity. The purpose of this study was to examine the relationship of socioeconomic status and living arrangements to nutritional intake of persons 55 years of age or older living in South Carolina. Data on 268 subjects who participated in the South Carolina Nutrition Survey, which was based on a true probability sample that represents the state's adult population, were analyzed. Eighty-nine percent of the subjects were found to have poor nutritional intake on the basis of a combination of four index nutrients: iron, vitamin B-6, calcium, and vitamin A. The data did not show a relationship between living arrangements and nutrient intake but affirmed the literature findings that support a positive relationship between low socioeconomic status and inadequate nutritional intake.

241 NAL Call No: 389.8 J824
Relationship of vitamin A and vitamin E intake to fasting plasma retinol, retinol-binding protein, retinyl esters, carotene, alpha-tocopherol, and cholesterol among elderly people and young adults: increased plasma retinyl esters among vitamin A-supplement users.
Krasinski, S.D.; Russell, R.M.; Otradovec, C.L.; Sadowski, J.A.; Hartz, S.C.; Jacob, R.A.; McGandy, R.B.
Baltimore, Md. : American Society for Clinical

Quick Bibliography Series

Nutrition; 1989 Jan.
American journal of clinical nutrition v. 49 (1): p. 112-120. charts; 1989 Jan. Includes 50 references

Language: English

Descriptors: Vitamin supplements; Nutritional assessment; Blood plasma; Carrier proteins; Carotenes; Age; Adults; Elderly; Men; Women

Abstract: We studied the relationships of supplemental and total vitamin A and supplemental vitamin E intake with fasting plasma biochemical indicators of vitamin A and vitamin E nutritional status among 562 healthy elderly people (aged 60-98 y) and 194 healthy young adult (aged 19-59 y) volunteers. All subjects were nonsmokers. For the young adults, plasma retinol was significantly greater in males than in females (p less than 0.01); retinol was not related to supplemental vitamin A intake for either group. Fasting plasma retinyl esters demonstrated a significant increase with vitamin A supplement use. For supplemental vitamin A intakes of 5001-10 000 IU/d, a 2.5-fold increase over nonusers in fasting plasma retinyl esters was observed for elderly people (p less than 0.05) and a 1.5-fold increase for young adults (p greater than 0.20). For elderly people, greater fasting plasma retinyl esters were associated with long-term vitamin A supplement use (greater than 5 y) and biochemical evidence of liver damage. Elderly people who take vitamin A supplements may be at increased risk for vitamin A overload.

242 NAL Call No: 389.8 AM34
Relationships between sour and salt taste perception and selected subject attributes.
Chauhan, J.
Chicago, Ill. : The Association; 1989 May.
Journal of the American Dietetic Association v. 89 (5): p. 652-658. charts; 1989 May. Includes 32 references.

Language: English

Descriptors: Taste sensitivity; Salt; Acidity; Age; Alcoholic beverages; Tobacco smoking

Abstract: The relationships between perceived sour and salt taste intensity and selected subject attributes--taste/smell deficits, salt use, avoidance of salt/sour tasting foods, smoking, alcohol use, dentures, education, and prescription drug use--were investigated in 180 subjects (30 men and 30 women/age group: 20 to 29 years [young], 70 to 79 years [old], 80 to 99 years [very old]). Multiple regression analyses showed that the taste intensity

estimates were significantly related to age, sex, stimulus concentration, self-reported taste/smell deficits, avoidance of salt/sour tasting foods, salt use, alcohol, smoking, education, and use of certain drugs (drugs for the treatment of arthritis, drug-acting on the gastrointestinal tract and heart system, and diuretics/drugs for the treatment of hypertension). The very old showed deficits in perception of the low suprathreshold tastant levels, with near normal perception of the high levels, whereas the old exhibited deficits at all levels. Wearing dentures was the only single variable not selected in any of the final regression models, due to the high prevalence and homogeneity in use of dentures in the elderly groups. Further work using a stratified sample and a large number of subjects is warranted to elucidate the influence of subject attributes on taste perception.

243 NAL Call No: RC262.C5N8
The relative validity of a retrospective estimate of food consumption based on a current dietary history and a food frequency list.

Bakkum, A.; Bloemberg, B.; Staveren, W.A. van; Verschuren, M.; West, C.E.
Hillsdale, N.J. : Lawrence Erlbaum Associates, Inc; 1988.

Nutrition and cancer v. 11 (1): p. 41-53. charts; 1988. Includes 24 references.

Language: English

Descriptors: Man; Food consumption; Validity; Accuracy; Dietary history; Etiology; Nutrient intake; Frequency; Nutritional assessment

Abstract: Extract: The relative validity of information and food consumption in the distant past was assessed by combining a dietary history (referring to the recent past) with a food frequency list (monitoring major changes over the past 12-14 years). This approach was evaluated in a study of two groups of apparently healthy elderly people (mean age 80 years) who had participated in a food consumption study 12-14 years before the start of the present study. One group consisted of 18 harbor employees who retired subsequent to the initial assessment of food intake. On the average, each member of this group had reduced his food consumption by about 1,000 kcal. The other group consisted of 46 elderly men and women who had retired before their food consumption was measured initially. This group had not markedly changed their food intake. The results showed that both groups overestimated changes in their food intake and that the systematic overestimation and

Nutrition and the Elderly

random error were similar for both groups. If the men in both groups were combined to form one group, a valid ranking of subjects in small and large consumers of energy and most of the selected nutrients was possible. However, current food intake influenced the accuracy of the measurement of past food intake.(author)

244 NAL Call No: 389.8 J824
Riboflavin status of adolescent vs elderly Gambian subjects before and during supplementation.

Bates, C.J.; Powers, H.J.; Downes, R.; Brubacher, D., Sutcliffe, V.; Thurnhill, A.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Oct.

American journal of clinical nutrition v 50 (4): p. 825-829. charts; 1989 Oct. Includes 24 references

Language: English

Descriptors: Gambia; Riboflavin; Nutritional state; Nutrient intake; Blood plasma; Adolescents; Elderly

Abstract: Riboflavin status was measured by activation coefficient of erythrocyte glutathione reductase (EGRAC) in elderly and adolescent rural Gambians whose intakes were low. Fifty-one adolescents and 52 elderly subjects were each subdivided into six supplementation groups to receive 0.25, 0.5, 0.75, 1.0, 1.5, or 2.5 mg riboflavin/d on 5 consecutive days weekly for 5 wk. Before supplementation, EGRAC values were grossly at normal and differed (p less than 0.005) between the two female groups [elderly subjects 1.94 ± 0.33 (average \pm SD); adolescents, 1.72 ± 0.27] but not between the two male groups. With supplementation greater than 0.25 mg/d the values became nearly equal in the two age groups. Only at total intakes between 1.8 and 2.5 mg riboflavin/d did the subjects attain acceptable status, with a mean EGRAC of 1.3-1.4. Clearly there is a need for further study of the discrepancy between observed index values during supplementation and the accepted ranges of normality at intakes close to the recommended amounts.

245 NAL Call No: 389.8 J82

The role of folate and vitamin B12 in neurotransmitter metabolism and degenerative neurological changes associated with aging: proceedings of a workshop.

Green, R.

Bethesda, Md. : American Institute of Nutrition; 1989 Jun.

The Journal of nutrition v. 119 (6): p. 841-842; 1989

Jun. This is a summary of a workshop held May 19-20, 1988.

Language: English

Descriptors: Folic acid; Neurotransmitters; Human nutrition research

Abstract: Under the sponsorship of the National Institute on Aging and the National Institute of Diabetes and Digestive and Kidney Diseases, the Life Sciences Research Office of the Federation of American Societies for Experimental Biology held a workshop entitled, "The Role of Folate and Vitamin B-12 in Neurotransmitter Metabolism and Degenerative Neurological Changes Associated with Aging." The purpose of the May 1988 workshop was to bring together scientists from various disciplines to identify opportunities for research on an important topic relating to neuroscience, nutrition and aging.

246 NAL Call No: RC682.N87
The role of oxidized lipids in heart disease and aging.

Smith, T.L.; Kummerow, F.A.

Boca Raton, Fla. : CRC Press; 1987.

Nutrition and heart disease / editor, Ronald R. Watson. v. 1 p. 45-64. ill., charts; 1987. Literature review. Includes 132 references.

Language: English

Descriptors: Cardiovascular diseases; Lipid peroxidation; Risks; Peroxides; Lipid metabolism

Abstract: A technical report reviews the ways lipid peroxidation (LP) may be involved in atherosclerosis, describes a model of how peroxidation could be atherogenic, and discusses the possible relationships between aging, LP, and heart disease. Attention is given to: the incidence of LP in heart disease patients; cell injury attributed to LP; interactions among endothelial cells, LP, and atherosclerosis; LP and other factors implicated in atherosclerosis at the cellular level; the source of LP products and cellular protection against LP; and the association of LP with aging and heart disease. The hypothesis is presented that LP plays a part in the initiation of heart disease.

247 NAL Call No: RA784.F6

The role of taste and smell in nutrition. Effects of aging, disease state, and drug.

Schiffman, S.S.

London : Libbey; 1987.

Food and health : issues and directions / edited by Mark L. Wahlqvist ... [et al.]. p. 85-91. charts; 1987.

Quick Bibliography Series

(International perspectives in food, diet and health, 9951-869x ; 1). Includes 26 references.

Language: English

Descriptors: Taste sensitivity, Smell, Organoleptic traits; Aging; Obesity; Carcinoma; Therapy

Abstract: An overview discusses the status of the chemical senses, which play an important role in nutrition and food intake, throughout the lifespan, including taste and smell abnormalities due to normal aging, disease status, and ethical drug usage. The role of taste and olfaction in obesity and cancer also are addressed. A wide variety of disorders affecting taste and smell are tabulated. Treatments of taste and smell disorders are described.(wz)

248 NAL Call No: 389.1 W892

The role of zinc in human nutrition.

Ronaghy, H.A.

Basel : S. Karger; 1987.

World review of nutrition and dietetics v. 54: p. 237-254. charts; 1987. Literature review. Includes 70 references.

Language: English

Descriptors: Zinc; Mineral deficiencies; Nutrient requirements; Patient care; Surgery; Mineral supplements

Abstract: A variety of studies are discussed concerning various aspects of zinc (Zn) nutrition in humans. Attention is given to: the biological essentiality of Zn; its metabolic turnover; the causes and clinical manifestations of Zn deficiency; the effects of Zn deficiency on the developing fetus; Zn deficiency and requirements in infants, adolescents, and the elderly; the clinical importance of Zn in surgical patients; and Zn supplementation.

249 NAL Call No: 389.8 J824

The roles of inflammation and iron deficiency as causes of anemia.

Yip, R.; Dallman, P.R.

Baltimore, Md. : American Society for Clinical Nutrition; 1988 Nov.

American journal of clinical nutrition v. 48 (5): p. 1295-1300. charts; 1988 Nov. Includes 32 references.

Language: English

Descriptors: Iron deficiency anemia; Inflammation; Erythrocyte sedimentation rate; Socioeconomic status; Children; Men; Women

Abstract: Inflammatory disease as well as iron deficiency may play an important role in the cause

of anemia in the United States. We evaluated the relationships between Fe deficiency, inflammatory disease, and anemia using data from the First National Health and Nutritional Examination Survey (NHANES I). Fe nutrition index was based on the ratio of serum Fe to Fe-binding capacity (Fe:TIBC) and inflammatory index was based on erythrocyte sedimentation rate (ESR). Groups with the highest prevalence of anemia were younger children, young women, and elderly men. Fe deficiency (low Fe:TIBC) was most common among the anemic children and young women but rare in anemic elderly men. Conversely, inflammation (high ESR) was most common among anemic elderly individuals. The prevalence of anemia was more than twice as high in the lowest than in the highest income group. Relative contributions of Fe deficiency and inflammation to anemia did not differ substantially among income groups.

250 NAL Call No: RM214.N87

Selection of supplements.

Debruyne, L.K.; Rolfes, S.R.

Philadelphia, Pa. : George F. Stickley Company; 1987 Apr.

Nutrition clinics v. 2 (2): p. 1-23. charts; 1987 Apr. Includes 15 references.

Language: English

Descriptors: USA; Vitamin supplements; Mineral supplements, Food and nutrition controversies; Age groups; Risks; Nutrient requirements; Guidelines

Abstract: A technical report discusses public misconceptions and facts concerning the need for vitamin-mineral supplements, focusing on the hype of the supplement industry and food faddists, and arguing that such needs, if any, be diagnosed and treated by a physician, not be self-prescription. Specific attention is given to the supplement needs of infants, teenagers, women of childbearing age, pregnant women, the elderly, strict dieters, vegetarians, people with chronic diseases, and people who smoke, drink, or take regular medications. Guidelines are included for selecting the right supplement, and data on safe nutrient ranges and information on certain nutrients and foods not needed in supplements are presented. Basic information and tabular data on vitamin and mineral needs, supplements for infants, children, adults, and elderly, and food group selection plans are appended.(wz)

251

NAL Call No: 389.8 B773

Nutrition and the Elderly

Selenium balance studies in apparently healthy and housebound elderly people eating self-selected diets.

Bunker, V.W.; Lawson, M.S.; Stansfield, M.F., Clayton, B.E

Cambridge : Cambridge University Press; 1988 Mar.

The British journal of nutrition v. 59 (2): p. 171-180. ill., charts; 1988 Mar. Includes 41 references.

Language: English

Descriptors: United Kingdom; Balance studies; Mineral metabolism; Selenium; Nutrient intake

Abstract: Extract: Metabolic balance studies (5 d) for Se were conducted in twenty-four apparently healthy elderly people (age 69.6-85.4 years), and twenty housebound subjects (age 69.9-85.1 years) with chronic disease. During the study the subjects lived in their own homes, ate self-selected diets and continued their normal daily activities. Geometric mean daily dietary intakes of the two groups were significantly different (P less than 0.01), being 819 (range 310-1631) nmol for the healthy and 475 (range 233-1136) nmol for the household elderly. Daily intake of Se significantly correlated with balance in both groups. Solution of the regression equations gave theoretical daily requirements of 447 nmol for the healthy and 419 nmol for the housebound subjects. The healthy elderly were in positive balance of 148 nmol/d for Se and the housebound were in equilibrium with an overall mean retention of 43 nmol/d. Mean levels of Se in blood fractions for the healthy and housebound subjects were significantly different, being 1.65 and 1.40 micromole/l whole blood, 1.45 and 1.21 micromole/l plasma and 5.72 and 5.30 nmol/g haemoglobin in erythrocytes respectively. There was no difference in the whole-blood glutathione peroxidase (EC 1.11.1.9) activities between the groups. There were clear differences in Se status between the two groups of elderly people. The cause of the positive balance in the healthy subjects remains unexplained.(author)

252 NAL Call No: 448.9 AM37

'Senile' osteoporosis reconsidered.

Resnick, N.M.; Greenspan, S.L.

Chicago, Ill. : The Association; 1989 Feb 17.

JAMA : Journal of the American Medical Association v. 261 (7): p. 1025-1029; 1989 Feb 17 Literature review. Includes 78 references.

Language: English

Descriptors: Osteoporosis; Vitamin D; Vitamin

Deficiency; Therapy; Calcium; Estrogens; Exercise; Elderly

Abstract: Osteoporosis is a devastating, morbid, and costly condition whose ravages are felt most profoundly by women over age 70 years. Yet most research on its prevention and treatment has focused on perimenopausal women, although there are significant differences between perimenopausal and older women in factors related to bone mineral metabolism, rate of bone loss, the structural integrity of remaining bone, risk factors for fractures, and the types of fractures sustained. Currently recommended therapies, which slow bone loss in perimenopausal women, may be of less benefit for older women whose loss of bone has already slowed or ceased and whose remaining bone may be of inadequate quantity and quality to prevent fracture. Thus, the application of currently available modalities is unlikely to mitigate significantly the consequences of osteoporosis in this population. Further research is urgently needed, and some directions for future investigation are suggested.

253

NAL Call No: QP141.A1N88

Serum magnesium levels in nursing home patients.

Dave, D.M.; Katz, P.R.; Gutman, S.

Elmsford, N.Y. : Pergamon Press; 1987 Sep.

Nutrition research v. 7 (9): p. 981-984. charts; 1987 Sep. Includes 13 references.

Language: English

Descriptors: New York; Nutritional state; Magnesium; Nursing homes; Elderly

Abstract: Extract: The elderly nursing home patient is at risk for hypomagnesemia due to poor nutritional intake, multiple chronic disease states, and polypharmacy. In order to assess the need to screen for magnesium deficiency in a long-term care setting, serum magnesium levels were determined in 75 nursing home patients. Mild hypomagnesemia was observed in 3 of the patients, all with poor eating habits. Diuretic use, although common, was not associated with magnesium deficiency. Routine screening for hypomagnesemia in nursing home patients with adequate nutritional intake does not appear warranted.(author)

254

NAL Call No: 389.8 J824

Serum protein concentrations and bone mineral content in aging normal men.

Orwoll, E.S.; Weigel, R.M.; Oviatt, S.K.; Meier, D.E.; McClung, M.R.

Quick Bibliography Series

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Oct.

American journal of clinical nutrition v. 46 (4): p. 614-621. ill., charts; 1987 Oct. Includes 32 references.

Language: English

Descriptors: Men, Aging; Bone mineralization; Protein intake; Blood composition; Calcium

Abstract: Extract: Low serum protein levels and low dietary protein intakes possibly influence bone metabolism. To evaluate the relationships among serum protein concentrations, protein intakes, and bone mineral content (BMC), we have examined two populations of normal men. BMC was measured at two radial sites and a vertebral site. Aging was associated with a fall in BMC at all sites. Serum albumin levels also fell with age and were related to BMC. Albumin concentrations were associated with BMC when other variables (calcium and protein intake, vitamin D levels, parathyroid hormone, and urinary calcium) also were considered. Thus, alterations in protein metabolism may affect BMC and may play a role in the genesis of senile osteopenia.(author)

255 NAL Call No: 389.8 J824
Serum retinol levels of persons aged 4-74 years from three Hispanic groups.

Looker, A.C.; Johnson, C.L.; Underwood, B.A. Baltimore, Md. . American Society for Clinical Nutrition; 1988 Dec.

American journal of clinical nutrition v. 48 (6). p. 1490-1496. charts, 1988 Dec. Includes 29 references.

Language: English

Descriptors: Nutritional state; Blood serum; Nutrition surveys; Ethnicity; Hispanics; Children; Adolescents; Adults; Elderly

Abstract: Previous research suggests that Hispanics in this country may have poor vitamin A status. Using serum retinol data from the Hispanic Health and Nutrition Examination Survey, we examined the vitamin A status of Mexican Americans (MA), Cubans, and Puerto Ricans (PR) aged 4-74 y. MA had lower mean serum retinol levels and higher prevalences of serum retinol in the range 0.70-1.01 micromole/L than did Cubans in several age-sex groups. The prevalence (or percentage) of serum retinol in a range indicating possible risk of functional impairment was not elevated in any of the Hispanic groups except the females aged 18-44 . However, a high percentage of children an

adolescents in the three Hispanic groups had serum retinol values in ranges that might indicate less-than-optimal vitamin A status. Determination of vitamin A status requires a more definitive assessment than by serum vitamin A alone.

256 NAL Call No: TX341.C6
Service innovations in elderly nutrition programs. Washington, D.C. : Community Nutrition Institute; 1988 Nov17.

Nutrition week v. 18 (45): p. 4-5; 1988 Nov17.

Language: English

Descriptors: Elderly nutrition; Nutrition programs, Federal programs; Community feeding centers, Home delivered mea's; Program evaluation

Abstract: This article is an excerpt from a report entitled "Service Innovations in Elderly Nutrition Program: Strategies for Meeting Unmet Needs." This report, funded by the American Association of Retired Persons, evaluates the federal Elderly Nutrition Program and makes recommendations for improving services.

257 NAL Call No: HV696.F6B32
Service innovations in the elderly nutrition program strategies for meeting unmet needs.

Balsam, Alan L.; Rogers, Beatrice Lorge AARP Andrus Foundation, Tufts University, School of Nutrition Medford, Mass. . Tufts University School of Nutrition; 1988.

106 p. : ill. ; 28 cm. Cover title. July, 1988. Prepared under the auspices of the Tufts University School of Nutrition for the American Association of Retired Persons Andrus Foundation.

Language: English

Descriptors: Food relief--United States; Aged--Nutrition; Elderly; Senior citizens; Nutrition programs; Public agencies

Abstract: This report discusses the extent to which the federally funded Elderly Nutrition Program provides services to socially impaired elders, homeless elders, ethnic and linguistic minorities, and extremely impaired individuals with needs for more than one meal daily and/or weekend meals. Data on the extent to which these innovative services have been implemented was collected by means of a mailed questionnaire to a random sample of 568 nutrition project directors nationwide. Other information about the programs was collected, including: program size, mechanics of foodservice, project structure, and nutrition project

Nutrition and the Elderly

director characteristics.

258 NAL Call No: QP141.A1N88
Sex and age changes in serum zinc levels.
Rea, I.M.
Elmsford, N.Y. : Pergamon Press; 1989 Jan.
Nutrition research v. 9 (1): p. 121-125. charts; 1989
Jan. Includes 16 references.

Language: English

Descriptors: Zinc. Blood serum; Age groups; Elderly; Young adults; Males; Females; Sex differences

Abstract: Serum zinc was measured in 126 fit healthy subjects living at home, 31 elderly males (mean age 78.3 years), 30 young males (mean age 21.6 years), 33 elderly females (mean age 81.6 years) and 32 young females (mean age 22.1 years). There were highly significant differences for serum zinc between the four groups with respect to age and sex (p less than 0.0002). Mean serum zinc was significantly lower in the elderly male group compared to the young male group with a similar but non significant trend in female groups. Males, old and young, had higher serum zinc levels than females of the same age and differences were significant between young males and females. Serum zinc levels are therefore lower in very elderly subjects compared to young subjects irrespective of sex, in keeping with findings noted in less than 75 year olds. Sex differences in serum zinc persist even in very old age, with elderly males showing higher levels than females of the same age.

259 NAL Call No: 448.8 B77
Simple test of intestinal calcium absorption measured by stable strontium.
Milsom, S.; Ibbertson, K.; Hannan, S.; Shaw, D.; Pybus, J.
London : British Medical Association; 1987 Jul25.
British medical journal v. 295 (6592): p. 231-234.
ill., charts; 1987 Jul25. Includes 14 references..

Language: English

Descriptors: Calcium; Absorption; Digestive system diseases; Strontium; Tests

Abstract: Extract: A clinical test of intestinal calcium absorption has been developed using non-radioactive stable strontium as a calcium tracer. In nine elderly subjects there was a close correlation between the fractional absorption of strontium and radioactive calcium (^{45}Ca) during a five hour period after the simultaneous oral administration of the two tracers. Comparable precision was

achieved with each tracer in six subjects in whom the test was repeated after two weeks. The effect of food on strontium absorption was examined in a further 33 normal subjects (age 21-60 years), and the administration of the strontium with a standard breakfast was shown to reduce the variance at individual time points. A simplified test in which serum strontium concentration was measured four hours after the oral dose given with a standard breakfast was adopted as the routine procedure. The normal range (mean (2 SD)), established over 97 tests in 53 patients, was 7.0-18.0% of the dose in the extracellular fluid. A further 30 patients with possible disorders of calcium absorption (10 with primary hyperparathyroidism and 20 with coeliac disease) were studied by this standard test. In both groups of patients the mean four hour strontium values were significantly different from normal. This standard strontium absorption test allows assessment of calcium absorption with sufficient sensitivity and precision to have a wide application in clinical practice.(author)

260 NAL Call No: 389.8 J824
Social, psychological and physical factors affecting the nutritional status of elderly subjects: separating cause and effect.
Goodwin, J.S.
Baltimore, Md. : American Society for Clinical Nutrition; 1989 Nov.
American journal of clinical nutrition v. 50 (5): p. 1201-1209. charts; 1989 Nov. Includes 70 references.

Language: English

Descriptors: Nutritional state; Mental health; Intelligence; Alcoholic beverages; Nursing homes; Psychological factors; Elderly

Abstract: The fact that most of the important changes in nutritional status with age are not secondary to aging per se in no way detracts from their importance. This paper reviews three factors that can affect nutritional status in elderly adults: ethanol intake, cognitive status, and institutionalization. The discussion focuses on the difficulties in obtaining reliable information on the interaction of nutritional status with these factors.

261 NAL Call No: TX361.A3J63
Some aspects of sensory properties of food that relate to food habits and associated problems of elderly consumers.
Madeira, K.
Binghamton, N.Y. : Haworth Press; 1988.

Quick Bibliography Series

Journal of nutrition for the elderly v. 8 (4): p 3-24. charts; 1988. Includes 13 references.

Language: English

Descriptors: Food habits; Food preferences; Nutrition surveys; Senses; Consumer behavior

Abstract: A questionnaire was developed and administered to 30 elderly consumers to explore perceptions about sensory properties of food that relate to food habits and food-related problems. Fifty percent of the survey sample believed that their food habits had changed considerably in the past 10 years, and 47% believed that health problems influenced their food selection. Few reported changes in food preparation because of mastication problems. Approximately half believed that foods smelled and tasted as good to them now as foods did when the consumers were young. Sensory properties of food were significantly (p less than or equal to .05) more important than several other factors in making purchase decisions for food preparation in three specific social contexts.

262 NAL Call No: RA784.A1C3
Special considerations in meal planning for the elderly.

Kingston : Caribbean Food and Nutrition Institute, 1987.

Cajanus v. 20 (1): p. 17-22. ill; 1987.

Language: English

Descriptors: Developing countries; Elderly nutrition; Menu planning; Nutritional adequacy; Food preferences; Nutrient requirements; Guidelines

Abstract: Practical guidelines are given for planning meals for the elderly to ensure the nutritional adequacy of the elderly and reduce the potential for elderly malnutrition. Specific attention is focused on: the consistency of foods; the importance of their palatability and digestability; meal frequency and the attractiveness of meal presentation; the physical status of elderly subjects with respect to specific nutrient requirements; and guidelines on the eating environment, food variety, and meal planning. Menu compositions for elderly women and men are included.(wz)

263 NAL Call No: 389.8 J824
Status of anthropometry and body composition data in elderly subjects.

Chumlea, W.C.; Baumgartner, R.N.

Baltimore Md. : American Society for Clinical Nutrition; 1989 Nov.

American journal of clinical nutrition v. 50 (5). p.

1158-1166. ill; 1989 Nov. Includes 91 references.

Language: English

Descriptors: Body composition; Anthropometric dimensions; Laboratory methods; Impedance, Ultrasound; Ethnicity; Elderly; Men; Women

Abstract: Understanding the normal changes in the body and its composition with increasing old age and their health implications are important to the health care and nutritional support of elderly subjects. Distribution statistics for selected body measurements of persons aged 65-80 y are available from the national surveys. Recumbent anthropometric techniques and B-mode ultrasound may be applicable to measuring those greater than 80 y who have difficulty standing or are chair- or bedfast. The problems of estimating body composition in elderly subjects could be improved by using a four-compartment model. Noninvasive methods, such as anthropometry and bioelectric impedance, could be used to predict body composition in elderly subjects if appropriate equations were available and validated against direct methods. The most pressing need is for the development of suitable reference data for anthropometry and body composition in large representative samples of black, white, Hispanic and Oriental elderly persons in the US.

264 NAL Call No: 389.8 SCH6
Study to decide if calcium curbs bone loss.

Denver, Colo. : American School Food Service Association; 1987 Mar.

School foodservice journal v. 41 (3): p. 30. ill; 1987 Mar.

Language: English

Descriptors: Calcium, Osteoporosis; Mineral supplements; Menopause; Bone density

Abstract: The USDA Human Nutrition Research Center on Aging at Tufts University is studying the possible benefits of women taking calcium supplements after menopause. The study will note if there are benefits to supplementation; the long-term effects of supplementation; and any unsuspected risks associated with supplementation. The methodology of the study is discussed.

265 NAL Call No: 389.8 J824
Subnormal concentrations of serum selenium and plasma carnitine in chronically tube-fed patients.
Feller, A.G.; Rudman, D.; Erve, P.R.; Johnson, R.C.; Boswell, J.; Jackson, D.L.; Mattson, D.E.
Bethesda, Md. : American Society for Clinical

Nutrition and the Elderly

Nutrition; 1987 Feb.

American journal of clinical nutrition v. 45 (2): p. 476-483. charts; 1987 Feb. Includes 36 references.

Language: English

Descriptors: Selenium; Carnitine; Enteral feeding; Elderly; Nutritional requirements; Nursing homes

Abstract: Extract: Forty-seven tube-fed nursing home patients were investigated with regard to serum or plasma selenium (Se), carnitine, and red blood cell (RBC) glutathione peroxidase (GSH-Px). Thirty-six patients were tube fed with Isocal, and 11 were tube fed with Compleat B, an L-carnitine-containing formula. Eighteen elderly nursing home patients and 10 young adults served as controls. Serum Se and plasma carnitine were lowest (p less than 0.05) in the Isocal patients. In all 36 Isocal subjects, Se was below normal, and in 26% of Isocal patients RBC GSH-Px was also below normal. Free and total carnitine were below normal in most Isocal subjects. All 11 Compleat B patients had subnormal serum Se, but most had normal carnitine concentrations. These data suggest that enteral formulae in nursing homes should contain greater than 100 micrograms Se and on the order of a mmol carnitine/1600 kcal.(author)

266 NAL Call No: LB1025.14

Suggestions for extension home economists programming at congregate nutrition sites.

Newhouse, J.K.

Champaign, Ill. : Home Economics Education, University of Illinois at Urbana-Champaign; 1989 Nov.

Illinois teacher of home economics v. 33 (2): p. 76-78; 1989 Nov. Includes 9 references.

Language: English

Descriptors: Community feeding centers, Learning activities; Cooperative extension service; Nutrition programs; Public relations; Elderly

Abstract: The congregate nutrition site which provide group meals for older adults is an appropriate forum for educational outreach to the elderly by the Cooperative Extension Service. This article gives guidelines for planning programs which fit the needs and interests of the people who use these sites.

267 NAL Call No: 389.8 J82

Summary of the NATO advanced research workshop on dietary omega-3 and omega-6 fatty acids: biological effects and nutritional essen-

tiality.

Simopoulos, A.P.

Bethesda, Md. : American Institute of Nutrition; 1989 Apr.

The Journal of nutrition v. 119 (4): p. 521-528; 1989 Apr. Literature review. Includes 56 references.

Language: English

Descriptors: Essential fatty acids; Docosenoic acid; Eicosapentaenoic acid; Nutrient deficiencies; Lipid metabolism; Infant feeding; Diseases; Parenteral feeding; Elderly nutrition; Human nutrition research; Nato

Abstract: A number of human studies presented at the workshop indicate that the premature infant at birth is biochemically deficient in docosahexaenoic acid (DHA) in both the brain and liver phospholipids, and that DHA is essential for normal visual acuity. The amount of DHA necessary to maintain normal amounts of the liver and brain phospholipids postnatally is 11 mg/kg daily. Elderly patients on long-term gastric tube feedings and others on long-term intravenous fluids and on total parenteral nutrition are particularly prone to deficiencies of alpha-linolenic acid, eicosapentaenoic acid (EPA) and DHA. The amounts estimated to prevent deficiencies in the elderly are 800-1100 mg/d of alpha-linolenic acid and 300-400 mg/d of EPA and DHA combined. Preliminary data indicate that children with malnutrition and mucoviscidosis, women with toxemia, and elderly people have decreased amounts of DHA in plasma phospholipids. The omega-3 fatty acids lower triglycerides and, at high levels, lower cholesterol. The anti-aggregatory, antithrombotic and anti-inflammatory properties of omega-3 fatty acids have been confirmed, and a dose-response curve is emerging. Despite the increase in bleeding time, no clinical evidence of bleeding has been noted by the investigators in any of the studies. Clinical trials are necessary in order to precisely define the dose and mechanisms involved in defining the essentiality of omega-3 fatty acids in growth and development and their beneficial effects in coronary heart disease, hypertension, inflammation, arthritis, psoriasis, other autoimmune disorders, and cancer.

268 NAL Call No: 389.8 SCH6

Survey shows elderly receive good grades for nutrition.

Denver, Colo. : American School Food Service Association; 1987 Apr.

School foodservice journal v. 41 (1): p. 26. ill; 1987 Apr.

Quick Bibliography Series

Language: English

Descriptors: Massachusetts; Food habits; Elderly nutrition; Vitamin And mineral supplements; Nutritional assessment; Nutritional surveys

Abstract: Scientists at USDA's Human Nutrition Research Center on Aging at Tufts University recently reported that many U.S. elderly persons are well-nourished in spite of diminished appetites. Reports were based on a survey of nearly 700 healthy Boston residents over the age of 60 years, the first study to examine the nutritional status of this particular age group. Major findings of the study were that 1) daily calorie intakes averaged 200 to 400 calories lower for this age group than for middle-age people, 2) vitamin and mineral levels in blood samples were within normal ranges in spite of low dietary intakes of some nutrients, and 3) over 50% of the survey population took vitamin and/or mineral supplements, most commonly vitamins C and E, iron, zinc, and calcium. Another survey of 350 elderly people confined to nursing homes found that their nutritional status was comparable to that of the healthy elderly population. Additional analyses of survey findings are currently being conducted.(ajc)

269 NAL Call No: RC620.A1J6
Thermogenic response to an oral glucose load in man: comparison between young and elderly subjects.

Bloesch, D.; Schutz, Y.; Breitenstein, E.; Jequier, E.; Felber, J.P.

New York, N.Y. : John Wiley & Sons; 1988 Dec. Journal of the American College of Nutrition v. 7 (6) p. 471-483. charts; 1988 Dec. Includes 48 references.

Language: English

Descriptors: Glucose; Carbohydrate loading, Physiological functions; Heat production; Energy expenditure; Anthropometric dimensions; Blood glucose; Insulin; Blood lipids; Young adults; Elderly

Abstract: This article compares the glucose-induced thermogenesis of young adults and elderly adults and correlates the changes in thermogenesis to age and body composition.

270 NAL Call No: TX361.A3J63
A tool for screening nutritional and oral health of elderly residents in long-term care facilities.

Gilbride, J.A.; Bachman, R.

Binghamton, N.Y. : Haworth Press; 1988. Journal of nutrition for the elderly v. 7 (4). p. 9-21;

1988 Includes 34 references.

Language: English

Descriptors: Nutritional assessment; Dental health. Screening tests; Long term care; Elderly

Abstract: Nutritional and dental services are two components of the total medical care in nursing homes. Appropriate nutritional and dental care affect the level of functioning and quality of life for older people who reside in nursing homes. Coordination between the dietitian and the dentist may help to identify, solve, and even prevent nutritional as well as dental problems. A screening tool, such as the one presented in this paper, can identify those residents at high risk for nutritional deficiencies caused by poor oral health status. Screening identifies the level of risk and which residents should be further investigated by the dietitian, the dentist, or both.

271 NAL Call No: 389.8 J824
Total body potassium in aging humans: a longitudinal study.

Flynn, M.A.; Nolph, G.B.; Baker, A.S.; Martin, W.M.; Krause, G.

Baltimore, Md. : American Society for Clinical Nutrition; 1989 Oct

American journal of clinical nutrition v. 50 (4): p 713-717. charts; 1989 Oct. Includes 27 references.

Language: English

Descriptors: Body composition; Potassium; Body fat; Body lean mass; Aging; Longitudinal studies; Men; Women

Abstract: Total body potassium (TBK) data calculated from longitudinal measurements over 18 y of 40K by whole-body counting of 564 male and 61 female healthy humans in a 2-pi liquid scintillation counter show little change in females younger than 50 y compared with males of those ages. Males show less TBK from 41 y onward as they age, with most rapid rate of loss between 41 and 60 y. Females have a rapid loss of TBK when they are older than 60 y; the loss is at a greater rate than that of males. Percent total body fat calculated from total body weight and lean body mass (LBM) derived from TBK document greater adiposity in females at all ages except ages 51-60 y when females are similar to males in change in percent fat per year per centimeter.

272 NAL Call No: RM216.M342
Trace elements and immune response.

Chandra, R.K.

Nutrition and the Elderly

St. Louis, Mo. : C. V. Mosby Company; 1987 May.
Clinical nutrition v. 6 (3): p. 118-125. ill., charts;
1987 May. Includes 43 references.

Language: English

Descriptors: Trace elements; Immunity; Infection; Mineral deficiencies; Mineral supplements; Iron; Zinc; Copper; Selenium; Iodine; Blood composition

Abstract: Extract: Nutrition is a critical determinant of immunocompetence and risk of infection. Protein-energy malnutrition results in impaired cell-mediated immunity, decreased levels of serum complement components, phagocyte dysfunction, and reduced secretory antibody response on mucosal surfaces. Deficiencies of many micronutrients also impair immunity. Obesity and excessive intake of nutrients are associated with decreased immune responses. The interactions of nutrition and immunity have several practical implications, particularly for young children and elderly persons.(author)

273 NAL Call No: 389 8 AM34
Training in geriatrics for future dietitians.
Shoaf, L.R.; Kotanchev, N.S.
Chicago, Ill. : The Association; 1987 Mar.
Journal of the American Dietetic Association v. 87
(3): p. 330-333. ill., charts; 1987 Mar. Includes 41 references.

Language: English

Descriptors: USA; Geriatrics; Training; Dietitians; Nutrition education; Elderly nutrition; National surveys

Abstract: Extract: All ADA-accredited dietetic internship programs and coordinated undergraduate programs (CUPs) were surveyed to determine the inclusion of geriatric nutrition and education about the aged population. All responding programs included some exposure to geriatrics; however, CUPs provided more experiences than internship programs. Skilled- and intermediate-care facilities were the most prevalent sites of training.(author)

274 NAL Call No: 389.8 AM34
Upper arm anthropometric norms in elderly white subjects.
Falciglia, G., O'Connor, J.; Gedling, E.
Chicago, Ill. : The Association; 1988 May.
Journal of the American Dietetic Association v. 88
(5): p. 569-574. ill., charts; 1988 May. Includes 14 references.

Language: English

Descriptors: Nutritional assessment; Anthropometry - dimensions; Elderly; Standards
Abstract: Extract: The purpose of this study was to develop percentile norms for upper arm anthropometry for use in the nutrition assessment of the elderly. This survey involved the measurement of four parameters: triceps skinfold thickness, mid-upper arm circumference, mid-upper arm muscle circumference, and mid-upper arm muscle area. Measurements were collected from 746 white, noninstitutionalized ambulatory elderly persons living in the Cincinnati area. Age- and sex-specific percentile distributions for the four parameters indicated age-related trends. All four parameters were lowered with advancing age in elderly women and men aged 60 to 89 years. The degree of change with age varied among parameters but was consistently greater in elderly women than in men. Elderly women had larger triceps skinfold thickness but smaller mid-upper arm circumference, mid-upper arm muscle circumference, and mid-upper arm muscle area measurements than did elderly men. These anthropometric findings provide useful standards for ambulatory white elderly people. It is evident that upper arm anthropometry changes with age. As a result, and elderly person's nutritional status should not be evaluated with a single standard for the elderly. Surveys of the elderly should use appropriate age standards to ensure and accurate estimation of obesity or undernutrition in the elderly.(author)

275 NAL Call No: TX361.A3J63
The use of dietary fibre in a long-stay geriatric ward.
Finlay, M.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 8 (1). p. 19-30. charts; 1988. Includes 16 references.

Language: English

Descriptors: Food fiber; Wheat bran; Calcium; Digestive tract; Intestine transit time; Long term care; Elderly

Abstract: The effect of added dietary fibre on bowel function and calcium balance has been studied in a group of elderly long-stay patients. The results show that a regime of a basic fibre-enriched diet with supplementary bran improves bowel habit in patients with chronic constipation and diarrhoea without effect on serum calcium, phosphate and alkaline phosphatase levels. Unacceptable gastrointestinal side effects had been encountered

Quick Bibliography Series

but did not occur when the supplementary bran was added at ward level.

276 NAL Call No: TX361.A3J63
Use of supplemental feedings for nursing home residents.

Matthews, L.E.

Binghamton, N.Y. : Haworth Press, 1988.

Journal of nutrition for the elderly v. 7 (4): p. 35-41. charts; 1988.

Language: English

Descriptors: Supplements; Liquid diet; Protein content; Caloric intake, Nutritional intervention; Food costs; Long term care; Elderly

Abstract: Many elderly residents of nursing homes need food supplements because they do not get enough nutrients from regular meals. This article describes several types of supplements and gives protein and caloric contents of each. The relative costs of the supplements are also considered.

277 NAL Call No: TX361.A3J63
Using high fiber foods to meet resident needs.

Matthews, L.E.

Binghamton, N.Y. : Haworth Press; 1988.

Journal of nutrition for the elderly v. 7 (3): p. 47-52. charts; 1988. Includes 2 references.

Language: English

Descriptors: Food fiber; Bran, Whole grains; Fruit juices; Fruits; Vegetables; Blended foods, Constipation; Long term care; Elderly

Abstract: The use of dietary fiber to maintain bowel function for elderly patients in long-term care is discussed. Possible detrimental effects of high fiber foods are included.

278 NAL Call No: 389.S Z33
Vitamin A, E and C nutriture of elderly people in North Italy.

Porrini, M.; Simonetti, P.; Ciappellano, S.; Testolin, G.

Bern : Hans Huber; 1987.

International journal for vitamin and nutrition research v. 57 (4): p. 349-355. charts; 1987. Includes 27 references.

Language: English

Descriptors: Italy; Elderly nutrition; Nutritional state; Ascorbic acid; Vitamin A

Abstract: A study of the nutritional vitamin A, C, and E status of 313 non-institutionalized elderly residing in a small town and an agricultural village

in northern Italy revealed no vitamin A deficiency in either community, but a vitamin C deficiency was noted in over half of the people. An incidence of vitamin E deficiency in 10-20% of elderly in the small town also was noted. Copious clinical data on blood composition are included.

279 NAL Call No: QP141.A1N88

Vitamin and food supplement practices and nutrition beliefs of the elderly in seven western states. Sheehan, E.T.; Delett, A.; Read, M.H.; Bendel, B.; Bhalla, V.; Bock, M.A.; Harrill, I., Mitchell, M. Elmsford, N.Y. : Pergamon Press; 1989 Mar.

Nutrition research v. 9 (3): p. 251-258; 1989 Mar. Includes 15 references.

Language: English

Descriptors: Arizona; California; Colorado; Hawaii; New Mexico; Nevada; Washington; Vitamin And mineral supplements; Questionnaires; Nutrition information; Information sources; Information dissemination; Elderly

Abstract: A mail survey of 2,451 non-institutionalized subjects, including 595 sixty years or older living in seven Western states (Arizona, California, Colorado, Hawaii, New Mexico, Nevada and Washington) reported frequencies and factors associated with vitamin and mineral supplementation. Sixty percent of subjects over 60 years reported using one or more supplements with multiple vitamin, vitamin D and vitamin E being most frequently used (greater than 50%). Reasons for use ranged from lack of vitamin in food to energy and/or tiredness. Sources of nutrition information cited most frequently were newspapers and books. The level of usage in the elderly is a major concern for health providers as well as the lack of information from professional resources.

280 NAL Call No: TX341.H82

The vitamin and mineral intake of a free-living young elderly Australian population in relation to total diet and supplementation practices.

Baghurst, K.L.; Record, S.J.

London : J. Libbey; 1987 Oct.

Human nutrition : applied nutrition v. 41A (5): p. 327-337. charts; 1987 Oct. Includes 17 references.

Language: English

Descriptors: Australia; Nutrient intake; Vitamins; Minerals; Elderly; Nutritional state

Abstract: A survey assessed the vitamin and mineral intakes of a randomly-selected group of

Nutrition and the Elderly

free-living Australian elderly subjects in 2 age groups (65-69 and 70-75). The results indicated that the mean intakes of most nutrients were equivalent to or greater than recommended levels, with a lower percentage of elderly at risks for vitamin (folate, vitamin B6) and mineral (Ca, Zn, Mg, Cu) deficiencies than reported for less able elderly subjects. Differences and similarities in intakes between sexes also are discussed.(wz)

281 NAL Call No: 389.8 Z33
Vitamin B6 status of Finnish elderly. Comparison with Dutch younger adults and elderly. The effect of supplementation.

Tolonen, M.; Schrijver, J., Westermarck, T.; Halme, M.; Tuominen, S.E.J.; Frilander, A.; Keinonen, M.; Sarna, S.
Bern : Hans Hube.; 1988.

International journal for vitamin and nutrition research v. 58 (1): p. 73-77. charts; 1988. Includes 14 references.

Language: English

Descriptors: Finland; Netherlands; Pyridoxine; Nutritional state; Vitamin supplements

Abstract: A study assessed the vitamin B6 status of 45 Finnish elderly (31 women, 14 men; ages, 66-96), compared this with that reported earlier for Dutch elderly and younger adults, and examined the efficacy of a 1-year, 2 mg/day pyridoxine supplement in a double-blind, placebo controlled design. It was found that ca. 25% of the Finnish and Dutch elderly were deficient in vitamin B6, and that pyridoxine supplementation corrected this.(wz)

282 NAL Call No: 389.8 J824
Vitamin C status and nutrient interactions in a healthy elderly population.

Jacob, R.A.; Otradovec, C.L.; Russell, R.M.; Munro, H.N.; Hartz, S.C.; McGandy, R.B.; Morrow, F.D.; Sadowski, J.A.

Baltimore, Md. : American Society for Clinical Nutrition; 1988 Dec.

American journal of clinical nutrition v. 48 (6): p. 1436-1442. charts; 1988 Dec. Includes 54 references.

Language: English

Descriptors: Ascorbic acid; Blood plasma; Nutrient-nutrient interactions; Vitamin supplements; Elderly; Men; Women

Abstract: Vitamin C status and interactions with other nutrients were studied in 677 healthy, non-

institutionalized elderly people aged 60-98 y. Although 6% of the males and 3% of the females showed marginal vitamin C status (plasma ascorbic acid 11 to less than 23 micromole/L), only one person had a plasma ascorbic acid (AA) level less than 11 micromole/L. At all levels of total vitamin C intake, mean plasma AA levels were higher in females than males. Vitamin C supplement use was associated with generally higher blood levels of vitamins B-6, B-12, and E and folate in both sexes and with higher levels of retinol in females. However, after both age and the total dietary intake of the specific nutrient being examined were controlled for, plasma AA levels were significantly correlated only with plasma levels of vitamin F and folate in females.

283 NAL Call No: RC620.A1N8
Vitamin C status of free-living, cognitively impaired and unimpaired elders.

Spindler, A.A.; Renvall, M.A.; Ramsdell, J.W.; Mofidi, S.

Stoneham, Mass. : Butterworth; 1989 Apr.

Nutrition reports international v. 39 (4): p. 713-719. charts; 1989 Apr. Includes 15 references.

Language: English

Descriptors: Ascorbic acid; Nutrient intake; Dietary history; Nutritional state; Leukocytes; Mental ability; Elderly

Abstract: We compared the dietary intake of Vitamin C with level of leucocytic ascorbic acid in 16 cognitively impaired and 26 unimpaired elders. Intake of Vitamin C in men was higher in impaired than in the unimpaired subjects, while the converse was true in women, producing a significant (p less than 0.008) sex by cognition interaction. Ascorbic acid level in leucocytes, which was higher in women than in men, was significantly (p less than 0.01) affected by degree of impairment only in women. Low or deficient biochemical status was found in 44% of the impaired and 27% of the unimpaired subjects. Vitamin C consumption exceeded 60 mg/day in 2/4 elders with deficient and 7/10 elders with low biochemical status.

284 NAL Call No: 389.8 J824

Vitamin D status in the elderly: seasonal substrate deficiency causes 1,25-dihydroxycholecalciferol deficiency.

Bouillon, R.A.; Auwerx, J.H.; Lissens, W.D.; Pelemans, W.K.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Apr.

Quick Bibliography Series

American journal of clinical nutrition v. 45 (4): p. 755-763. ill., charts, 1987 Apr. Includes 37 references.

Language: English

Descriptors: Belgium; Vitamin D; Elderly nutrition; Nutritional state; Seasonal variation; Bone disorders; Metabolism; Calcium; Parathyroid hormone

Abstract: Extract: The seasonal variation of 25-hydroxycholecalciferol and 1,25-dihydroxycholecalciferol was analyzed in 240 elderly subjects (mean age: 78 yr) in Belgium. Serum 25-hydroxycholecalciferol was lowest from February until May (mean levels less than 25 nmol/L). Summer peak levels were, however, not higher than nadir levels in younger control subjects. A seasonal variation in total and free 1,25-dihydroxycholecalciferol concentrations was also observed in the geriatric population with a nadir in February and March (50 +/- 24 pmol/L). The peak values in summer (110 +/- 33 pmol/L) were not different from those of the younger controls. Serum calcium and phosphate were decreased whereas alkaline phosphatase and parathyroid hormone were increased throughout the year in the geriatric patients. Oral 25-hydroxycholecalciferol treatment rapidly normalized serum 1,25-dihydroxycholecalciferol concentrations in vitamin D-deficient subjects. Deficiency of both the vitamin D substrate and hormone is frequent in the elderly population in Belgium.(author)

285 NAL Call No: 448.8 L22
Vitamin D supplementation in the elderly.
Boston, Mass. : Little, Brown and Company; 1987 Feb07.

The Lancet v. 1 (8528): p. 306-307; 1987 Feb07. Includes 36 references.

Language: English

Descriptors: Vitamin D; Vitamin supplements; Bone resorption; Bone fractures; Disease prevention; Elderly

Abstract: A technical review examines published studies to assess whether vitamin D supplements should be given to elderly subjects. Further studies are needed to (1) verify that such supplementation reduces hip fracture incidence or reduces or prevents age-related bone loss in the femoral neck, and (2) the appropriate time for supplementation initiation and its duration. At minimum, it is argued that since vitamin D supplementation is easy and safe, evidence strongly favors such supplementation for high-risk patients.(wz)

286 NAL Call No: 389.8 J824
Vitamin requirements of the elderly.

Suter, P.M.; Russell, R.M.

Bethesda, Md. : American Society for Clinical Nutrition; 1987 Mar.

American journal of clinical nutrition v. 45 (3): p. 501-512; 1987 Mar. Literature review. Includes 156 references.

Language: English

Descriptors: Elderly nutrition; Vitamins; Nutrient requirements; Nutritional state

Abstract: Extract: In general, low dietary intakes can account for much of poor vitamin nutrition reported among various elderly populations. Despite problems in assessing vitamin nutrition in the elderly, the 1980 RDAs for thiamin, riboflavin, and ascorbic acid seem appropriate for those populations. However, RDAs for vitamin A and folate may be too high and the RDAs for vitamin D, vitamin B-6, and vitamin B-12 may be too low, due to specific age-related changes in the metabolism of these vitamins. For vitamin E, vitamin K, niacin, biotin, and pantothenic acid, the data is conflicting and/or insufficient to make a judgement about the appropriateness of the RDAs or to estimate safe and adequate daily intakes for the elderly.(author)

287 NAL Call No: RC620.A1N8
Vitamin A intake and vitamin A status in an elderly, institutionalized population.

Asciutti-Moura, L.S.; Guillard, J.C.; Fuchs, F.; Klepping, J.; Richard, D.

Stoneham, Mass. : Butterworth; 1989 Jun.

Nutrition reports international v. 39 (6): p. 1107-1115. charts; 1989 Jun. includes 32 references.

Language: English

Descriptors: Nutritional state; Blood plasma; Nutrient intake; Caloric intake; Nursing homes

Abstract: Vitamin A status of 30 institutionalized elderly subjects (7 men, mean age = 79 y and 23 women, mean age = 87 y) was evaluated using plasma retinol level, plasma retinol-binding protein level and the retinol dose response. The kilocalorie, protein, total fat and vitamin A intakes were also estimated using a five-day food record. Regarding energy, protein and total fat, the diets may be generally considered satisfactory in respect to the French Recommended Dietary Allowances (RDA). The mean dietary intake of vitamin A was greater than RDA. Only 5% of males and 2% of females consumed less than their individual vita-

Nutrition and the Elderly

min A requirements as determined by a probability method. Despite the adequacy of dietary vitamin A, 14% of male subjects and 21% of female subjects had a retinol dose response greater than 20% indicating a great number of elderly institutionalized people with depleted liver reserves.

288 NAL Call No: 389.8 AM34
Weight loss and senile dementia in an institutionalized elderly population.

Franklin, C.A.; Karkeck, J.
Chicago, Ill. : The Association; 1989 Jun.
Journal of the American Dietetic Association v. 89 (6): p. 790-792. charts; 1989 Jun. Includes 28 references.

Language: English

Descriptors: Alzheimer's disease; Senility; Weight losses; Underweight; Nutritional state; Nursing homes

Abstract: The purpose of this study was to determine whether patients with Alzheimer's disease or senile dementia have a higher incidence of underweight and weight loss than non-demented patients. In a retrospective study, the admission weight and height and the current weight of 36 patients with Alzheimer's disease and senile dementia of the Alzheimer's type were compared with those of 31 non-demented patients. Patients with Alzheimer's disease or senile dementia were significantly (p less than .05) further below minimum acceptable body weight on admission and were significantly (p less than .05) further below minimum acceptable body weight after an average of 17 months in the institution in comparison with the control group. Additional research is needed to determine whether the weight loss seen in patients with Alzheimer's disease or senile dementia is due to a metabolic aberration inherent in the disease or to management issues in feeding those patients.

289 NAL Call No: 389.8 AM34
What characterizes elderly women who overuse vitamin and mineral supplements?

Ranno, B.S.; Wardlaw, G.M.; Geiger, C.J.
Chicago, Ill. : The Association; 1988 Mar.
Journal of the American Dietetic Association v. 88 (3): p. 347-348. charts; 1988 Mar. Includes 10 references.

Language: English

Descriptors: Overfeeding; Women; Elderly nutrition; Nutritional state

Abstract: A regional survey assessed the socioeconomic, demographic, and health behavior characteristics of 72 elderly women regarding their use and misuse of vitamin and mineral supplements. The study identified some potentially important characteristics of overusers. It is suggested that screening of overusers first be conducted with elderly women with health problems and who live alone.(wz)

290 NAL Call No: aTX501.F66
When an older person needs help in the kitchen.

Parmley, M.A.
Washington, D.C. : U.S. Department of Agriculture; 1988.
Food news for consumers v. 5 (1): p. 8-9. ill; 1988.

Language: English

Descriptors: Food preparation; Elderly; Nutritional adequacy; Food safety; Risks; Guidelines

Abstract: A special report provides information and guidelines for aiding elderly people in safe and nutritious food preparations. Attention is focused on: the susceptibility of older people to food poisoning; nutritional problems of older people due to poor food selections; food handling problems associated with the elderly; the effect of money problems and attitudes on waste on proper food selection and food safety; and signs to look for concerning improper food preparation by the elderly. Ten specific tips on nutritional value and food safety are summarized.(wz)

291 NAL Call No: 389.8 J824
Zinc and immunocompetence in the elderly: baseline data on zinc nutriture and immunity in unsupplemented subjects.

Bogden, J.D.; Oleske, J.M.; Munves, E.M.; Lavenhar, M.A.; Bruening, K.S.; Kemp, F.W.; Holding, K.J.; Denny, T.N.; Louria, D.B.
Bethesda, Md. : American Society for Clinical Nutrition; 1987 Jul.
American journal of clinical nutrition v. 46 (1): p. 101-109. charts; 1987 Jul. Includes 54 references.

Language: English

Descriptors: Diet; Zinc; Immunity; Nutritional state; Mitogens

Abstract: Extract: Zinc nutriture and immune function were studied in 100 subjects, age 60-89 yr. Mean (+/- SD) zinc concentrations found were 84.8 +/- 15.5 micrograms/dL (13.0 +/- 2.4 microM) for plasma. 1.04 +/- 0.24 micrograms

Quick Bibliography Series

(0.016 +/- 0.004 micromoles)/10(9) cells for erythrocytes, 4.06 +/- 1.85 micrograms (0.062 +/- 0.028 micromoles)/10(9) cells for mononuclear cells, 3.91 +/- 1.77 micrograms (0.060 +/- 0.027 micromol.s)/10(9) cells for polymorphonuclear leukocytes, 0.53 +/- 0.39 micrograms (0.0081 +/- 0.0060 micromoles)/10(9) cells for platelets, and 222 +/- 101 micrograms (3.39 +/- 1.54 micromoles)/g for hair. Zinc ingestion was below the RDA in more than 90% of study subjects. The incidence of anergy to a panel of seven skin test antigens was 41%; responses to these antigens were significantly associated with the plasma zinc concentration. Subjects with depressed lymphocyte responses to mitogens had significantly lower platelet and significantly higher mononuclear cell zinc concentrations than those with normal responses.(author)

292 NAL Call No: TX361.A3J63
Zinc nutriture and cell mediated immunity in institutionalized elderly.
Soltesz, K.S.; Williford, J.H.; Renker, L.P.; Meserve, L.A.
Binghamton, N.Y. : Haworth Press; 1988.
Journal of nutrition for the elderly v. 8 (1): p. 3-17. cl ts; 1988. Includes 24 references.

Language: English

Descriptors: Zinc; Nutritional assessment; Immune response; Anthropometric dimensions; Mineral supplements; Nursing homes; Elderly
Abstract: This study examined the effects of increasing zinc nutriture on the cell mediated immune response of an institutionalized elderly population. A volunteer institutionalized population was identified and nutritional assessment consisting of a weighed food intake record and anthropometric measurements was conducted. Zinc status and delayed cutaneous hypersensitivity (DCH) skin tests were administered to determine immune responsiveness. Subjects were supplemented with 15 mg zinc gluconate for 28 days, after which the zinc status and immune status of each subject were reassessed. Results indicated that dietary zinc intakes were inadequate in 67% of the test population. After supplementation, serum zinc levels increased significantly (p less than .006), while results of the alkaline phosphatase test indicated no significant change. DCH reactions improved significantly (p less than .034) after zinc supplementation.

293 NAL Call No: TX341.H8

Zinc status and dark adaptation in patients subjected total gastrectomy: effect of zinc supplement 'ion.

Sandstrom, B.; Davidsson, L.; Lundell, L.; Olbe, L
London : J. Libbey; 1987 May.

Human nutrition : clinical nutrition v. 41C (3): p. 235-242. ill., charts; 1987 May. Includes 19 references.

Language: English

Descriptors: Zinc; Nutritional state; Elderly; Mineral deficiencies; Mineral supplements; Blood composition; Urine analysis

Abstract: A study involving measurements of serum zinc (Zn), urinary Zn excretion, and a simple dark adaptation test estimated Zn status in 10 totally gastrectomized elderly patients (1 woman, 9 men; ages 61-80) vs. an age-matched control group, before and after 4 weeks of Zn supplementation. Serum Zn (but not urinary Zn) was lower in the patients than in the controls before supplementation; Zn supplementation did not affect dark adaptation, but increased both serum and urinary Zn levels in the patients.(wz)

294 NAL Call No. TX341.B75
Zinc--a precious metal.

Turnbull, A.J.; Thompson, R.. .A.

London : The Foundation; 1989 Jan.

BNF Nutrition bulletin - British Nutrition Foundation v. 14 (1): p. 23-35. charts; 1989 Jan. Includes 46 references.

Language. English

Descriptors: Zinc; Nutrient intake; Recommended dietary allowances; Physiological functions; Nutrient sources; Food fiber; Bioavailability; Mineral deficiencies; Human nutrition research

Abstract: Zinc is an essential trace element and is required for such processes as growth, immunity and reproduction. Many people, even in this country, have a marginal intake of zinc and several common dietary constituents can impair its absorption. Despite homeostatic regulation of zinc absorption, subclinical zinc deficiency may be common, although it is difficult to detect. Attention to the zinc nutrition of at-risk groups such as the malnourished, children, pregnant women and the elderly is therefore required. Zinc is relatively non-toxic and should be used in the prevention and treatment of zinc deficiency and possibly other disorders.

Author Index

- AARP Andrus Foundation, Tufts University,
School of Nutrition 257
- Acevedo, F. 223
- Adolfsson, R. 195
- Agarwal, N. 223
- Aguzzi, F. 198
- Alfthan, G. 27
- American Dietetic Association 84
- Ames, B.D. 21
- Anda, R. 226
- Anderson, E.J. 135
- Anderson, J.J.B. 99
- Andres, R. 160
- Arabatzis, K. 75
- Arija, V. 182
- Aro, A. 158
- Arthur, E. 49
- Asciutti-Moura, L.S. 107, 287
- Asp, E.H. 128
- August, D. 59
- Auwerx, J.H. 284
- Bachiman, R. 270
- Baghurst, K.I. 280
- Bailey, A.J. 208
- Bailey, L.B. 102, 141
- Baker, A.S. 271
- Bakkum, A. 243
- Balsam, A. 60
- Balsam, A.L. 54
- Balsam, Alan L. 257
- Baptista, R.J. 58
- Barbosa, J.C. 75
- Barrett-Connor, E. 72
- Bartlett, B.J. 50
- Basu, T.K. 6
- Bates, C.J. 244
- Baumgartner, R.N. 263
- Beare-Rogers, Joyce 66
- Beavers, I. 180
- Bedell, B.A. 239
- Bendel, B. 279
- Beresteyn, E.C.H. van 46
- Berg, H. van den 81
- Bergstrom, N. 197
- Betts, N.M. 22
- Bhagavan, H.N. 15, 18
- Bhalla, V. 279
- Bhat, K.S. 217
- Bidlack, W.R. 193
- Bilic-Pesic, L. 196
- Bistrrian, B.R. 58
- Black, A.E. 25
- Blackburn, G.L. 58
- Bloemberg, B. 243
- Bloemberg, B.P.M. 39
- Bloesch, D. 269
- Blue, L.S. 183
- Blumberg, J.B. 85
- Bock, M.A. 279
- Bogden, J.D. 291
- Boss, G.R. 131
- Boswell, J. 265
- Bouchard, C. 53
- Bouillon, R.A. 284
- Bowen, P. 24
- Bower, M.E. 240
- Bracker, M.D. 105
- Breitenstein, E. 269
- Bridgewater, R. 10
- Briefel, R.R. 156
- Brody, Jane E. 143
- Brubacher, D. 244
- Brubacher, G. 196
- Bruening, K.S. 291
- Bunker, V.W. 96, 150, 157, 251
- Buzina, R. 196
- Caindec, N. 117
- Campbell, A.J. 74
- Carlos, J.P. 151
- Carroll, M. 226
- Carroll, S.S. 142, 218
- Cayten, C.G. 223
- Ceccanti, M. 24
- Cerda, J.J. 141
- Champagne, C.D. 58
- Chandra, R.K. 179, 272
- Channing L. Bete Co 2
- Chapuy, M.C. 35
- Chapuy, P. 35
- Chauhan, J. 6, 219, 242
- Chen, L.H. 12
- Chernoff, R. 127, 201
- Chesnut, C.K. 186
- Cheung, C.K. 228
- Choi, E. 17
- Chumlea, W.C. 222, 263
- Ciappellano, S. 278
- Clark, V.A. 32
- Clay, G. 53
- Clayton, B.E. 96, 150, 157, 251
- Cleeman, J. 226
- Clifton-Bligh, P. 166
- Cole, T.J. 25
- Coleman, P. 7
- Contaldo, F. 199
- Cook-Newell, M.E. 12

Quick Bibliography Series

- Cooper, J.K. 79
 Coper, H. 110
 Cotter, R. 106
 Cotugna, N. 224, 225
 Coulson, A.H. 32
 Courpron, P. 51
 Dallal, G.E. 17, 215
 Dallman, P.R. 249
 Darling, M.E. 128
 Dave, D.M. 253
 Davidsson, L. 293
 Davies, H.E.F. 16
 Davies, L. 220
 Davis, L.G. 104
 Davis, R. 86
 De Looy, A.E. 37
 De Lorenzo, A. 24
 DeBruyne, L.K. 173
 Debruyne, L.K. 250
 Debry, G. 171
 Dechavanne, M. 51
 Delett, A. 279
 Delhey, D.M. 135
 DeMicco, F.J. 89
 Denny, T.N. 291
 Di Biase, G. 199
 Dillon, J.C. 97
 Donald, E.A. 6
 Downes, R. 244
 Doyle, B. 77
 Draughn, P. 4
 Duursma, S.A. 46
 Echizen, H. 26, 98
 Eldemire, D. 200
 Elfving, S. 158
 Erve, P.R. 265
 Ervin, R.B. 140
 Falciglia, G. 274
 Fanelli, M.T. 1, 93
 Felber, J.P. 269
 Feldman, E.B. 118
 Feller, A.G. 106, 265
 Fernandez-Ballart, J. 182
 Ferro-Luzzi, A. 24
 Fidanza, F. 199
 Field, H.P. 216
 Finlay, M. 275
 Fishman, P. 49
 Flendrig, J.A. 138
 Florencio, C.A. 111
 Flynn, M.A. 271
 Foltz, M.B. 8
 Food Research and Action Center 154
 Forette, B. 43
 Fosmire, G.J. 140
 Fox, H.M. 29, 176
 Francis, R.M. 145
 Franklin, C.A. 288
 Friedman, H. 24
 Frier, H.I. 144
 Frilander, A. 281
 Frongillo, E.A. Jr 57
 Fryberger, M.F. 131
 Fuchs, F. 107, 287
 Fulwood, R. 226
 Garrard, J. 221
 Garry, P.J. 11, 15, 18, 47, 215
 Gates, A.D. 161
 Gedling, E. 274
 Gee, M. 6
 Geiger, C.J. 289
 Gerber, J. 129
 Gergans, G.A. 206
 Gershoff, S.N. 94
 Gettner, S. 24
 Gilbride, J.A. 270
 Gilligan, C. 65
 Goddijn, E. 39
 Goldberg, J.P. 94
 Good, R.A. 178
 Goodwin, J.M. 11
 Goodwin, J.S. 11, 260
 Gorajscan, D. 196
 Gorajscan, M. 196
 Gordillo, B. 182
 Gordon, A.K. 214
 Goulding, A. 74
 Gouws, E. 159
 Gray, G.F. 167
 Green, R. 245
 Greenspan, S.L. 252
 Gref, C.G. 158
 Gruffudd, S. 16
 Guillard, J.C. 107, 287
 Guo, S. 222
 Gussler, Judith D. 209
 Gutman, S. 253
 Guzman, M.A. 187
 Habicht, J.P. 115, 116
 Haines, C. 226
 Hale, W.E. 92
 Hallfrisch, J. 160
 Hallmans, G. 15
 Halme, M. 281
 Hanes, F.A. 37
 Hankin, J.H. 62

Author Index

- Hannan, S. 259
Harrill, I. 279
Harris, L.J. 52
Harris, S.B. 73
Harris, T. 156
Hartz, S.C. 70, 91, 94, 153, 229, 230, 241, 282
Hawrysh, Z.J. 6
Hayes, P.A. 231
Heaney, R.P. 188
Hegsted, D.M. 236
Hegsted, M. 4
Hemphill, K. 53
Henderson, C. 32
Hermus, R.J.J. 165
Hester, D.D. 189
Heymsfield, S.B. 31
Hibbard, J.H. 104
Hildreth, G.J. 4
Hinks, L.J. 96, 150
Ho, S.C. 228
Hodges, P.A.M. 52, 113
Hodges, R.E. 237
Hodkinson, H.M. 63
Holding, K.J. 291
Hollingsworth, J.W. 105, 131
Holt, V. 40, 114
Hood, S.J. 215
Hoover, J.L.B. 186
Horton, H.W. 56
Horwath, C.C. 76
Hughes, R.E. 16
Hulshof, K.F.A.M. 90, 165
Hunt, C. 11
Hunt, I.F. 32
Hunt, W.C. 47
Ibbertson, K. 259
Imbach, A. 78, 122
Itoh, R. 26, 98
Jackson, D.L. 117, 265
Jacob, R.A. 17, 70, 91, 229, 230, 241, 282
Jacobs, R.M. 32
Jacques, P. 17
Jacques, P.F. 91, 153
James, O.F.W. 10
Janghorbani, M. 59
Janicke, B. 110
Jensen, H.M. 136
Jequier, E. 269
Johnson, C.L. 255
Johnson, J.M. 52
Johnson, M. 75
Johnson, R.C. 106, 265
Johnston, P.K. 32
Jones, E. 16
Juan, D. 130
Jung, D. 130
Jurdi-Haldeman, D. 212
Jutsum, P.J. 200
Kamen, Y. 31
Kampen-Donker, M. van 39
Kannel, W.B. 174
Karas, G. Brian 88
Karreck, J. 288
Kasim, S. 44
Katz, P.R. 253
Kehayias, J. 31
Keinonen, M. 281
Kelleher, J. 216
Kemp, F.W. 291
Kergoat, M.J. 78
Kervliet, N.I. 232, 233
Khaw, K.T. 72
Kime, R.E. 104
King, P. 147, 148
Kingsley, L. 211
Kistemaker, C. 90, 165
Kivela, S.L. 27, 175
Klauber, M.R. 105
Kleinman, J.C. 156
Klepping, J. 107, 287
Kohrs, M.B. 40, 114
Kostiainen, E. 158
Kotanchek, N.S. 273
Krasinski, S.D. 241
Krause, G. 271
Kromhout, D. 39
Kronl, M. 3, 7, 100, 119, 124
Kuczmariski, R.J. 155
Kummerow, F.A. 246
Kunkel, M.E. 186
Kurtz, C.S. 14
Kyllastinen, M. 158
Labadarios, D. 159
LaCroix, A.Z. 227
Lagarde, M. 51
Lanzetta, R. 199
Lanzilli, A. 199
Laramée, S.H. 135
Lavenhar, M.A. 291
Lawson, M.S. 150, 157, 251
Leaf, A. 9
Lee, D.Y. 231
Lehman, W. 61
Lei, K.Y. 67
Leigh, S. 10
Leighton, L.S. 223

Quick Bibliography Series

- Leklem, J. 211
Leklem, J.E. 218
Leonard, R. 109
Lerclerc, B.S. 78
Lezenne Coulander, C. de 39
Lichtman, S. 31
Lieberman, L.S. 14
Liebman, B. 86
Lieux, E.M. 19
Lipschitz, D.A. 201
Lipson, S. 227
Lissens, W.D. 284
Litchford, M.D. 162
Locniskar, M. 169, 13
Looker, A.C. 255
Lopez, L.M. 115, 116
Lorenz, E. 178
Louria, D.B. 291
Lowik, M.R.H. 81, 90, 165
Lugo, N.J. 105
Lundell, L. 293
Lyle, R.M. 28
Macatangay, L.A. 111
Madeira, K. 261
Maenpaa, P.H. 27
Maiani, G. 24
Mak, Y.T. 228
Maññ, N.L. 4
Manore, M.M. 61, 71, 142, 218
Marcus, R. 36
Marti-Henneberg, C. 182
Martin, W.M. 271
Martinez, O.B. 125, 137
Massey, L.K. 133
Matsumoto, M. 80, 235
Matthews, L.E. 108, 210, 213, 276, 277
Mattson, D.E. 106, 117, 206, 265
May, F.E. 92
McAvay, G. 139
McBean, L.P. 172
McClung, M.R. 254
McGandy, R.B. 69, 70, 91, 94, 229, 230, 241, 282
McParland, B.E. 74
Meier, D.E. 254
Melby, C.L. 28
Mendy, F. 51
Merz, B. 113
Meserve, L.A. 292
Mesfin, E.
Messina, M.S. 54
Meunier, P.J. 35
Miles, T.P. 227
Miller, D.G. 58
Miller, A. 144
Miller, L.T. 231, 233
Milsom, S. 259
Mitchell, M. 279
Mobarhan, S. 24
Mofidi, S. 283
Moneta, A. 198, 231
Moore, M.T. 92
Morgan, J.C. 132
Morley, J.E. 192
Morris, J.A. 25
Morrow, F.D. 282
Mullee, M.A. 96
Muncie, H.L. Jr 68
Munro, H.N. 194, 229, 230, 282
Munves, E.M. 291
Murakami, K.
Murphy, N.J. 32
Mwonya, R. 180
Nagraj, H.S. 117, 206
Napier, A.K. 212
Nelson, M. 25
Newhouse, J.K. 266
Nielsen, C.C. 103
Nieman, D.C. 75
Niewind, A. 119
Nissinen, A. 27, 175
Noel, M.B.
Noel, M.M.B. 21
Nolen, N.R. 221
Nolph, G.B. 271
Nordstrom, J. 40, 114
Nygren, C. 195
O'Connor, J. 274
Obermann-de Boer, G.L. 39
Ockhuizen, T. 81
Ohls, J.C. 132
Oka, J. 26, 98
Olbe, L.
Oleske, J.M. 291
Olivey, A. 113
Olson, J.A. 237, 238
Orwoll, E.S. 254
Osteraas, G. 60
Otradovec, C.L. 70, 229, 230, 241, 282
Otte, R.G. 131
Oviatt, S.K. 254
Papa, A. 48, 290
Peacock, M. 145
Pekkanen, J. 27
Pelemans, W.K. 284
Pennell, M.D. 163
Peters, H. 70

Author Index

- Peters, J.E. 34
PetitClerc, C. 78
Philadelphia Health Management Corporation,
Thomas Jefferson University, Dept. of Gerop-
sychiatry 87
Pierce, R.P. Jr 113
Pierson, R.N. Jr 31
Pirhonen, A.
Pirkanen, A. 27
Pitchumoni, C.S. 223
Poehlman, E.T. 28
Pollitzer, W.S. 99
Porrini, M. 198, 278
Posner, B.M. 132
Postmes, T.J. 138
Potter, J.F. 10
Potts, M.C. 101
Powers, H.J. 244
Pratt, C. 211, 214
Probart, C.K. 104
Pybus, J. 259
Ralston, P.A. 180
Ram.mohan, M. 130
Ramsdell, J.W. 283
Ranaldi, L. 24
Ranno, B.S. 289
Rao, V. 3
Raymakers, J.A. 46
Rea, I.M. 258
Read, M.H. 279, 280
Remington, P. 226
Renker, L.P. 292
Renvall, M.A. 283
Resnick, N.M. 252
Rezek, J. 22
Rheaume, Y. 149
Rhyne, R.L. 47
Richard, D. 107, 287
Riley, M.E. 149
Roche, A.F. 222, 13, 84, 121
Rogers, Beatrice Lorge 257
Rolfes, S.R. 250
Ronaghy, H.A. 248
Ross Laboratories 209
Rossouw, J.E. 159
Roughead, Z.K. 186
Rovati, P. 198
Rozovski, S.J. 177
Rudman, D. 106, 117, 206, 265
Rudman, I.W. 117, 206
Rudman, L. 126
Russell, R.M. 70, 91, 153, 194, 229, 230, 241, 282,
286
Ryan, A.S. 8
Ryan, V.C. 161, 240
Sadowski, J.A. 215, 241, 282
Sahyoun, N.R. 70
Saltman, P. 131
Saltman, P.D. 105
Sanchez, C.J. 11
Sandall, M.J. 133
Sandman, P.O. 195
Sandstrom, B. 293
Sarna, S. 281
Schaafsma, G. 46
Schaefer, E.J. 17
Schiffman, S.S. 134, 247
Schlenker, Eleanor D.
Schoeller, D.A. 41
Scholes, J.E. 57
Schrijver, J. 81, 281
Schutz, Y. 269
Scythes, C.A. 163
Sels, J.P. 138
Sempos, C. 226
Shackleton, P.A. 239
Shaw, D. 259
Sheehan, E.T. 279
Shephard, R.J. 20
Sherman, K.M. 75
Shifflett, P.A. 52
Shoaf, I 136, 273
Shrott, A. 119
Shultz, T.D. 75
Simmons, W.K. 200
Simonetti, M.S. 199
Simonetti, P. 198, 278
Simopoulos, A.P. 204, 267
Sinha, D.P. 152, 200
Smiciklas-Wright, H.A. 140
Smith, C.H. 193
Smith, E.L. 65
Smith, J.L. 197
Smith, P.E. 65
Smith, T.L. 246
Sobal, J. 68
Sodha, N. 96
Soltész, K.S. 292
Spindler, A.A. 283
Srinivasan, T.R. 216
Stanek, K.L. 29
Stansfield, M.F. 150, 157, 251
Staveren, W.A. van 243
Stavljenic, A. 196
Steen, B. 30
Steinbaugh, M.L. 222

Quick Bibliography Series

- Stephenson, S. 38
Stewart, R.B. 92
Strause, L.G. 105, 131
Strong, J.P. 187
Suboticanec, K. 196, 201
Sulsky, S. 153
Suri, Y.P. 33
Suski, N.S. 103
Sutcliffe, V. 244
Suter, P.M. 194, 286
Swaminathan, R. 228
Talbott, M.C. 232, 233
Taylor Kennedy, J. 100
Taylor, J. 3
Testolin, G. 198
Thomas, A.J. 96
Thomas, P. 11
Thomas, R.G. 92
Thompson, R.P.H. 294
Thurnhill, A. 244
Tolonen, M. 281
Tortrat, D. 43
Trampusch, T.S. 183, 189
Truax, M. 205
Tuominen, S.E.J. 281
Turnbull, A.J. 294
Underwood, B.A. 255
Underwood, B.C. 75
United States, Food and Drug Administration 207
Uusitalo, U. 158
Van Eck, M. 159
Van Marrelo, C. 49
VanderJagt, D.J. 15, 18, 47
Vaughan, L.A. 61, 71, 142, 218
Vericel, E. 51
Verschuren, M.
Volicer, L. 149
Wakefield, L.M. 162
Walden, O. 231
Walford, R.L. 73
Walker, B.E. 216
Wang, J. 31
Wardlaw, G.M. 289
Warwick, Z.S. 134
Watson, Ronald R. 170
Weddle, D.O. 214
Wedel, M. 81
Weigel, R.M.
Weindruch, R. 73
Weinstock, C.P. 123
Welch, T. 203
West, C.E. 243
Westenbrink, S. 81, 90, 165
Westermarck, T. 281
White, L. 227
White-O'Connor, B. 68
Whitley, A.J. 216
Willcox, K.K. 67
Williamson, D.F. 226
Williford, J.H. 292
Wilson, Jane Weston 88
Wilson, S.E. 197
Wilson, W. 211
Winblad, B. 195
Winkler, L.L. 19
Wojnarowski, S.M. 184
Wolfe, M.D. 151
Wolmark, Y. 43
Woo, J. 228, 156
Yamada, K. 26
Yeung, D.L. 122, 163
Yip, R. 249
Young, V.R. 59
Zimmerman, S. 124
Zimmerman, S.A. 163

Subject Index

- 1,25-dihydroxycholecalciferol 117
Absorption 145, 237, 259
Accuracy 243
Acetylsalicylic acid 3, 100
Acidity 242
Acyl-coa desaturase 107
Adaptation 110
Adolescence 141
Adolescents 123, 244, 255
Adult day care 38
Adults 37, 241, 255
Age 219, 241, 242
Age differences 130
Age groups 6, 250, 258
Aged--Nutrition 87, 88, 121, 154, 257
Aging 6, 7, 24, 30, 44, 46, 65, 67, 73, 86, 89, 102, 110, 118, 144, 155, 164, 168, 170, 171, 178, 247, 254, 271
Albumins 203, 223
Alcoholic beverages 10, 11, 27, 153, 242, 260
Alcoholism 167
Aluminum 64, 167
Alzheimer's disease 64, 103, 149, 162, 167, 195, 288
Amino acid metabolism 16
Amino acids 106
Animal experiments 73
Anorexia 192
Anthropometric dimensions 1, 14, 70, 78, 142, 155, 186, 195, 198, 200, 222, 228, 230, 263, 269, 274, 292
Anthropometry 13
Antibodies 105
Apolipoprotein 17
Appetite 83, 130
Arizona 279
Arthritis 68
Ascorbic acid 15, 16, 17, 18, 26, 100, 137, 158, 237, 278, 282, 285
Atherosclerosis 44, 187
Attitudes 89
Audiovisual aids 84
Australia 280
Balance studies 150, 157, 251
Behavior 118
Behavior modification 175
Belgium 284
Beliefs 211
Beta-carotene 24
Bioavailability 102, 294
Blended foods 277
Blood chemistry 47, 117
Blood composition 1, 18, 230, 254, 272, 293
Blood glucose 269
Blood lipids 24, 27, 158, 226, 269
Blood plasma 12, 51, 106, 137, 196, 215, 216, 217, 218, 241, 244, 282, 287
Blood pressure 28, 200
Blood serum 24, 27, 43, 107, 138, 142, 255, 258
Blood serum protein 223
Body composition 14, 30, 31, 41, 99, 155, 263, 271
Body density 31
Body fat 155, 271
Body lean mass 271
Body water 41
Body weight 29, 130, 206, 222, 223
Bone density 32, 99, 206, 208, 264
Bone disorders 284
Bone fractures 117, 206, 285
Bone mineralization 36, 46, 145, 166, 254
Bone resorption 46, 74, 285
Bone strength 188
Bone weight 99
Brain 173
Bran 33, 277
Brassica 119
Bread 138
C-amp 74
C-peptide 138
Calcium 22, 27, 32, 33, 35, 36, 99, 145, 186, 188, 199, 208, 209, 217, 252, 254, 259, 264, 275, 284
California 147, 279
Caloric intake 3, 39, 76, 122, 125, 153, 181, 228, 230, 287
Calorie-restricted diets 73
Campylobacter 112
Canada 122, 124, 163
Canada--fre 7
Carbohydrate loading 269
Carcinoma 43, 178, 247
Cardiovascular diseases 43, 65, 67, 82, 92, 118, 169, 170, 171, 174, 226, 246
Caribbean 152
Carnitine 16, 265
Carotenes 78, 241
Carrier proteins 241
Cartilage 68
Case studies 108, 113, 189
Cataract 217
Cell mediated immunity 179
Child nutrition 235
Children 14, 37, 84, 123, 146, 249, 255
Cholesterol 43, 44, 91, 138, 158, 215, 226
Choline 64
Chromium 45, 217

Quick Bibliography Series

- Clinical investigations 1, 46, 198
 Clinical nutrition 108
 Cognitive development 11
 Collagen 208
 Colorado 279
 Community action 126
 Community feeding centers 60, 256, 266
 Community nutrition 50
 Community programs 38
 Computer analysis 125
 Computer applications 54
 Computer graphics 61
 Computer software 61
 Conferences 205
 Congressional reports 5
 Constipation 277
 Consumer attitudes 22, 144, 172, 224
 Consumer behavior 48, 55, 56, 172, 224, 225, 261
 Consumer education 239
 Consumer preferences 42, 55, 56
 Consumer protection 55
 Consumer surveys 225
 Consumers 34, 52
 Consumption patterns 10, 11, 42, 48, 55, 101, 119, 175
 Convenience foods 172
 Cook-chill system 147, 148
 Cookery 88
 Cooperative extension service 266
 Copper 59, 96, 150, 217, 272
 Cost benefit analysis 235
 Costs 58, 80, 112
 Creatinine 78
 Curriculum 136
 Data analysis 25, 140
 Data collection 62
 Dehydroascorbic acid 15
 Demography 4, 9, 89
 Dental caries 151
 Dental health 151, 186, 270
 Depression 63
 Developing countries 262
 Diabetes 65, 83, 138, 169, 211
 Diagnosis 46, 78, 133, 135, 207, 220, 223
 Diet 2, 15, 28, 29, 33, 51, 63, 65, 72, 74, 98, 138, 143, 158, 178, 199, 201, 291
 Diet counseling 49, 214
 Diet planning 143
 Diet study techniques 61
 Dietary carbohydrates 76, 193
 Dietary factors 99, 170, 187
 Dietary fat 66
 Dietary guidelines 88, 168, 193
 Dietary history 25, 62, 139, 140, 165, 243, 283
 Dietary proteins 193
 Dietary surveys 39, 40, 76, 111, 152, 165, 182, 211
 Dietetic education 136
 Dietitians 8, 21, 54, 84, 273
 Digestive disorders 203
 Digestive system diseases 259
 Digestive tract 275
 Digoxin 82
 Dining facilities 77
 Disease prevention 46, 65, 118, 144, 164, 166, 170, 174, 204, 207, 209, 285
 Diseases 117, 193, 267
 Diuretics 82
 Docosenoic acid 267
 Drug combinations 83
 Drug interactions 84
 Drug therapy 82, 117, 125
 Drug-nutrient interactions 84
 Drugs 124
 Eating habits 39, 48, 103, 130, 221
 Eating out 34, 48
 Eating patterns 90, 123, 172
 Educational programs 136
 Eicosapentaenoic acid 267
 Elderly 1, 3, 4, 7, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 22, 24, 26, 27, 28, 29, 31, 32, 34, 36, 38, 40, 41, 43, 47, 48, 50, 51, 58, 59, 63, 64, 70, 71, 74, 75, 76, 77, 81, 82, 83, 84, 87, 89, 92, 93, 94, 96, 98, 101, 103, 104, 106, 107, 108, 113, 114, 115, 116, 117, 122, 123, 124, 125, 127, 129, 130, 133, 135, 137, 138, 139, 140, 141, 142, 146, 147, 148, 149, 150, 151, 152, 153, 155, 157, 158, 159, 161, 162, 163, 165, 167, 169, 174, 176, 179, 180, 182, 183, 188, 189, 190, 195, 197, 198, 202, 206, 208, 211, 212, 214, 215, 218, 219, 221, 222, 224, 225, 227, 228, 229, 231, 232, 236, 240, 241, 244, 252, 253, 255, 257, 258, 260, 263, 265, 266, 269, 270, 274, 275, 276, 277, 279, 280, 282, 283, 285, 290, 292, 293
 Elderly nutrition 5, 8, 9, 21, 33, 35, 45, 50, 52, 54, 57, 60, 62, 67, 69, 78, 80, 85, 86, 88, 91, 95, 97, 100, 109, 111, 116, 119, 121, 122, 128, 131, 132, 133, 135, 136, 154, 156, 160, 164, 171, 173, 175, 177, 181, 184, 186, 190, 191, 192, 193, 194, 199, 200, 201, 203, 210, 216, 220, 230, 233, 256, 262, 267, 268, 273, 278, 284, 286, 289
 Employed women 172
 Employer employee relationships 89
 Energy consumption 115
 Energy expenditure 149, 162, 269
 Energy intake 96, 101, 111, 149, 182, 229
 Energy requirements 20, 171

Subject Index

- England 101
Enteral feeding 189, 265
Environmental factors 113
Enzyme activity 26, 131
Enzyme deficiency 97
Epidemiology 151, 187
Erythrocyte sedimentation rate 249
Erythrocytes 26
Essential fatty acids 97, 267
Estrogens 36, 188, 252
Ethnic foods 129
Ethnic groups 99
Ethnicity 4, 17, 99, 186, 255, 263
Etiology 243
Exercise 2, 65, 86, 252
Experimental diets 24
Facilities 210
Family size 90, 172
Fasting 106
Fat 66
Fat consumption 76
Fat metabolism 60
Fats 193
Fatty acids 66, 107
Federal aid 5
Federal programs 5, 60, 80, 109, 126, 132, 191, 202, 235, 256
Feeding habits 123
Females 258
Ferritin 137, 142
Fetus 118
Fiber 33, 138
Finland 27, 175, 281
Fish 71
Fish oils 204
Flavor enhancers 134
Flavorings 56
Flavors 212
Fluid intake 110
Folates 141
Folic acid 102, 137, 245
Food acceptability 56
Food allergies 120, 169
Food and nutrition controversies 23, 33, 67, 69, 109, 191, 234, 250
Food beliefs 181, 224
Food consumption 25, 39, 68, 76, 90, 165, 193, 221, 243
Food costs 37, 276
Food distribution programs 146
Food fiber 76, 102, 169, 275, 277, 294
Food groups 40, 68
Food habits 40, 111, 114, 172, 211, 261, 268
Food industry 56, 129, 144
Food intake 24, 77, 81, 110, 125, 134, 157, 182, 212
Food intolerance 124
Food misinformation 104
Food policy 37
Food preferences 29, 40, 42, 48, 55, 75, 103, 113, 114, 120, 122, 169, 181, 261, 262
Food preparation 290
Food production 42
Food quality 52, 128
Food relief--United States 257
Food safety 290
Food service 147
Food service management 19, 54
Food serving methods 77
Food stamp program 80, 115, 116, 132
Food technology 56
Foodborne diseases 112
Frequency 243
Fruit juices 277
Fruits 277
Gambia 244
Games 94
Genetic factors 119
Genetic variation 120
Genotype nutrition interaction 120
Geriatrics 121, 273
Gingivitis 151
Glucose 138, 269
Glucose tolerance 45
Glutathione reductase 26
Groups 101
Growth stages 185
Guar gum 138
Guidelines 1, 34, 84, 85, 86, 177, 201, 207, 220, 226, 250, 262, 290
Hawaii 187, 279
Health 2, 124, 125, 152, 200
Health care 54, 133, 164
Health care costs 133, 135
Health hazards 84, 143
Health promotion 73, 86, 118, 168, 175, 204
Health services 52, 95, 104, 133
Heart--Diseases--Diet therapy 170
Heart--Diseases--Nutritional aspects 170
Heart--Diseases--Prevention 170
Heat production 269
Height 28
Hemoglobin a1 138
Heredity 99
High density lipoprotein 17, 44, 91, 138, 158, 215
Hispanics 255

Quick Bibliography Series

- Home care 50, 54, 127, 133
Home delivered meals 49, 50, 57, 124, 128, 202, 231, 256
Hong kong 228
Hospices 214
Hospital diets 159
Hospital food service 148
Hospitals 54, 111, 133, 135, 184
Household income 172
Households 90
Human nutrition research 139, 237, 238, 245, 267, 294
Hunger 146
Hydroxyproline 74
Hyperglycemia 83
Hypertension 29, 72, 82
Hypoglycemia 83
Immune response 196, 232, 292
Immunity 131, 272, 291
Immunology 178
Impedance 263
Incidence 124, 206
Income 69, 193
Indexes 28
India 217
Individual characteristics 236
Individual feeding 108
Infant feeding 267
Infants 14, 141
Infection 272
Inflammation 249
Influenzavirus 105
Information dissemination 279
Information sources 104, 161, 279
Institutional food service 111, 213
Institutions 113
Insulin 269
Intake 74
Intelligence 260
Interior design 77
Interviews 125
Intestinal absorption 24, 59, 102
Intestine transit time 24, 275
Iodine 272
Iron 12, 98, 116, 141, 142, 173, 199, 272
Iron deficiency anemia 12, 141, 203, 249
Italy 198, 199, 278
Jamaica 200
Japan 26, 98
Kentucky 12
Labor force 89
Labor productivity 19
Laboratory methods 31, 41, 263
Lactation 141
Learning activities 266
Legislation 5, 205
Leukocytes 150, 216, 283
Life cycles 141, 185
Life style 129, 144, 152, 182, 200
Linoleic acid 51
Linolenic acid 51
Lipid metabolism 67, 97, 187, 204, 246, 267
Lipid peroxidation 246
Lipids 51
Lipids in human nutrition 66
Liquid diet 276
Liquids 221
Literature reviews 225
Living conditions 240
Long term care 77, 108, 148, 228, 270, 275, 276, 277
Longevity 73
Longitudinal studies 39, 40, 43, 72, 139, 271
Louisiana 187
Low calorie foods 123, 172
Low income groups 37, 71, 115, 116, 142, 146, 218
Low-fat diet--Recipes 87
Lunch 101
Lymphocytes 15, 232, 233
Magnesium 217, 253
Males 258
Malnutrition 77, 78, 79, 95, 135, 191, 220, 223
Man 138, 157, 243
Marketing 8
Maryland 146
Massachusetts 17, 54, 60, 268
Maternal nutrition 118
Medicaid 210
Medicare 210
Men 3, 4, 15, 16, 17, 22, 27, 39, 41, 59, 76, 81, 99, 106, 117, 142, 153, 155, 163, 165, 196, 206, 217, 224, 225, 226, 227, 241, 249, 254, 263, 271, 282
Menopause 36, 264
Mental ability 173, 283
Mental health 63, 260
Menu planning 262
Menus 108
Metabolism 63, 209, 284
Methodology 14, 30, 152
Michigan 21, 126
Milk protein 102
Mineral content 32
Mineral deficiencies 192, 199, 248, 272, 293, 294
Mineral metabolism 188, 251

Subject Index

- Mineral supplements 35, 92, 105, 131, 216, 225, 248, 250, 264, 272, 292, 293
- Minerals 76, 153, 160, 193, 280
- Missouri 114, 148
- Mitogens 291
- Models 73
- Monocytes 15
- Morbidity 227
- Mortality 9, 43, 72, 187, 223, 227
- Mothers 126
- Muscle tissue 229
- National surveys 8, 27, 81, 112, 115, 116, 132, 156, 227, 273
- Nato 267
- Nebraska 22
- Neonates 179
- Netherlands 39, 81, 90, 138, 165, 281
- Neurotransmitters 63, 245
- Neutron activation analysis 31
- Nevada 279
- New Mexico 279
- New York 49, 57, 205, 253
- Niacin 45, 186, 199
- Nitrogen 157
- Nucleotidase 131
- Nursing homes 19, 107, 117, 133, 201, 203, 206, 210, 221, 253, 260, 265, 287, 288, 292
- Nutrition programs 87
- Nutrient availability 98
- Nutrient contents 61
- Nutrient data banks 61
- Nutrient deficiencies 64, 83, 173, 181, 198, 267
- Nutrient excesses 64
- Nutrient intake 1, 6, 12, 15, 18, 25, 26, 29, 32, 37, 39, 51, 59, 70, 71, 75, 76, 81, 99, 101, 103, 107, 111, 122, 123, 125, 128, 130, 132, 139, 140, 142, 153, 159, 161, 162, 163, 165, 167, 177, 182, 194, 195, 196, 198, 218, 231, 240, 243, 244, 251, 280, 283, 287, 294
- Nutrient requirements 23, 36, 60, 69, 95, 97, 110, 122, 160, 168, 170, 171, 177, 189, 190, 194, 204, 234, 236, 237, 238, 248, 250, 262, 286
- Nutrient sources 143, 294
- Nutrient-drug interactions 3, 82, 83, 84, 85, 100, 102, 190, 194
- Nutrient-nutrient interactions 33, 45, 145, 282
- Nutrition 66, 120, 143, 143, 151, 169, 179, 185, 185
- Nutrition education 21, 49, 52, 93, 94, 104, 175, 176, 180, 211, 220, 273
- Nutrition information 22, 94, 239, 279
- Nutrition knowledge 4, 21, 160, 161, 180, 239
- Nutrition labeling 129
- Nutrition physiology 7
- Nutrition policy 118, 190
- Nutrition programs 19, 38, 50, 54, 126, 146, 147, 202, 256, 257, 266
- Nutrition research 118
- Nutrition surveys 25, 75, 90, 154, 172, 255, 261
- Nutrition--Study and teaching 2
- Nutritional adequacy 111, 128, 165, 172, 199, 201, 262, 290
- Nutritional assessment 1, 14, 47, 53, 70, 71, 78, 121, 133, 135, 152, 155, 156, 167, 183, 202, 223, 240, 241, 243, 268, 270, 274, 292
- Nutritional disorders 7
- Nutritional intervention 53, 97, 132, 164, 212, 226, 276
- Nutritional requirements 121, 149, 265
- Nutritional state 1, 8, 12, 18, 26, 27, 69, 70, 71, 78, 79, 81, 90, 96, 106, 107, 116, 137, 141, 142, 150, 152, 159, 161, 163, 173, 178, 186, 193, 194, 195, 196, 197, 198, 199, 200, 217, 222, 227, 228, 230, 244, 253, 255, 260, 278, 280, 281, 283, 284, 286, 287, 288, 289, 291, 293
- Nutritional support 60, 189, 214
- Nutritional surveillance 17, 183, 184, 202
- Nutritional surveys 104, 153, 154, 156, 163, 193, 240, 268
- Nutritional value 144
- Obesity 168, 247
- Odors 212
- Ohio 50
- On-site food preparation 147, 148
- Ontario 137
- Ontario--fre 125
- Oregon 104
- Organoleptic traits 247
- Osteoarthritis 3
- Osteoporosis 46, 65, 100, 145, 166, 188, 194, 206, 207, 208, 209, 252, 264
- Osteoporosis--Congresses 209
- Osteoporosis--Prevention 207
- Overfeeding 289
- Paradental disease 151
- Parathyroid hormone 35, 284
- Parenteral feeding 58, 189, 267
- Parenteral feeding 127
- Participation 57, 115
- Patient care 33, 53, 78, 85, 121, 149, 170, 183, 184, 189, 201, 209, 210, 220, 221, 248
- Patient compliance 57, 211
- Patient education 49, 84, 207, 226
- Peer relationships 211
- Peroxides 246
- Personnel 89

Quick Bibliography Series

- Ph 102
 Philippines 111
 Phospholipids 51
 Phosphorus 166
 Physical activity 20, 68, 99, 125
 Physical environment 77
 Physically handicapped persons 34
 Physiological functions 193, 203, 269, 294
 Physiology 41, 190
 Platelets 51
 Pneumonia 227
 Policy 164
 Polyunsaturated fatty acids 97, 204
 Popular works 2, 87
 Post secondary education 185
 Potassium 72, 271
 Potassium deficiency 82
 Poultry meat 71
 Prediction 42, 221, 222, 223, 224
 Pregnancy 141
 Preventive nutrition 143
 Problem solving 21
 Product development 144
 Professional competence 21
 Professional education 185
 Program development 53
 Program effectiveness 60, 132, 176
 Program evaluation 19, 49, 94, 128, 231, 239, 256
 Program participants 80, 235
 Prostaglandins 204
 Protein content 276
 Protein energy malnutrition 178, 192, 195, 203, 228
 Protein intake 32, 76, 96, 228, 229, 230, 254
 Protein requirement 157
 Psychological factors 11, 211, 240, 260
 Public agencies 202, 257
 Public relations 266
 Public services 164
 Puerto rico 187
 Purgatives 82
 Pyridoxal phosphate 233
 Pyridoxine 81, 218, 229, 230, 232, 233, 281
 Questionnaires 20, 54, 62, 125, 180, 279
 Ratios 59
 Recipes 88, 143
 Recommended dietary allowances 23, 111, 141, 160, 193, 234, 236, 294
 Recommended dietary intakes 23, 96, 122, 237, 238
 Reference standards 47
 Reference works 121, 170, 209
 Regional surveys 114, 175, 180, 186
 Regression analysis 215
 Reports 60
 Research methodology 20, 25, 62, 151, 156, 188
 Restaurants 34, 48
 Reviews 36, 46, 67, 170, 192
 Riboflavin 26, 137, 244
 Risks 33, 43, 67, 90, 92, 170, 187, 192, 226, 227, 246, 250, 290
 Rural population 22
 Salmonella 112
 Salt 242
 School food service 89, 205
 School lunches 80
 Screening tests 270
 Seasonal variation 284
 Selenium 27, 251, 265, 272
 Self feeding 221
 Senility 64, 173, 288
 Senior citizens 257
 Senses 261
 Sensory disorders 77
 Seventh day adventists 28
 Sex differences 10, 11, 18, 104, 125, 130, 142, 162, 219, 258
 Skeleton development 36
 Skin fold thickness 229
 Smell 6, 134, 247
 Snacks 123
 Socioeconomic status 4, 8, 11, 95, 104, 180, 191, 197, 240, 249
 Sodium 29
 Sodium chloride 74, 219
 Source fat 67, 107
 South africa 159
 South australia 76
 South Carolina 161, 186, 240
 Spain 182
 Stable isotopes 59
 Standards 1, 274
 State government 146
 Strength 196
 Stroke 72
 Strontium 259
 Study guides 207
 Supplemental feeding programs 52, 57, 60, 69, 80, 109, 126, 132, 186, 235
 Supplements 22, 79, 97, 103, 104, 124, 140, 142, 167, 208, 276
 Support systems 211
 Surgery 79, 248
 Surveys 4, 21, 57, 92, 210, 226, 231
 Task analysis 19
 Taste 6, 119, 134, 212

Subject Index

- Taste sensitivity 219, 242, 247
Taste threshold 6, 120
Teaching materials 185, 207
Teaching methods 93
Television 94
Test construction 93
Tests 259
Textbooks 185
Therapeutic diets 46, 121, 127, 167
Therapy 209, 247, 252
Thermal processes 102
Thiamine 26, 173, 186, 199
Tobacco smoking 27, 242
Trace elements 194, 272
Training 89, 273
Transketolase 26
Trauma 189
Treatment 46
Trends 9, 42, 55, 80, 112, 118, 129
Triglycerides 24, 44, 138, 215
Ultrasound 263
Underweight 288
United Kingdom 10, 37, 96, 251
Urban areas 10, 199
Urban population 22
Urine analysis 186, 293
USA 5, 8, 23, 28, 69, 80, 95, 112, 115, 116, 123,
132, 191, 192, 210, 226, 231, 234, 235, 250, 273
Utilization 104
Vaccination 105
Validity 243
Variations 139, 140
Vegetables 119, 124, 277
Vegetarianism 75
Vegetarians 28, 32
Videotapes 176
Vitamin A 27, 100, 234, 238, 278
Vitamin And mineral supplements 71, 125, 268,
279
Vitamin B12 23, 137, 173
Vitamin D 27, 35, 36, 74, 145, 166, 188, 209, 252,
284, 285
Vitamin Deficiency 167, 199, 252
Vitamin E 27, 91
Vitamin K 215
Vitamin supplements 15, 18, 23, 35, 81, 91, 158,
192, 196, 218, 225, 232, 233, 234, 241, 250, 281,
282, 285
Vitamins 76, 153, 160, 193, 194, 196, 280, 286
Washington 133, 279
Weight 28, 203
Weight gain 68, 83
Weight losses 68, 195, 288
Weight reduction 211
Wheat bran 82, 102, 275
Whole grains 277
Women 4, 15, 16, 17, 22, 24, 32, 41, 43, 59, 74,
75, 76, 81, 99, 140, 142, 145, 153, 155, 163, 165,
197, 207, 217, 224, 225, 226, 227, 241, 249, 263,
271, 282, 289
Women's health services 207
Workable days 231
Workshops (programs) 176
Young adults 24, 59, 215, 219, 258, 269
Zinc 33, 59, 92, 96, 105, 131, 140, 150, 173, 178,
192, 196, 216, 217, 248, 258, 272, 291, 292, 293,
294