

ED 323 423

CG 022 745

AUTHOR Updegrove, Natalie A.
 TITLE Childhood Obesity and Cardiovascular Disease: January 1985-May 1990. Quick Bibliography Series.
 INSTITUTION National Agricultural Library, Beltsville, MD.
 REPORT NO QB-90-59
 PUB DATE Jun 90
 NOTE 57p.
 AVAILABLE FROM U.S. Department of Agriculture, National Agricultural Library. Public Services Division, Room 111, Beltsville, MD 20705.
 PUB TYPE Reference Materials - Bibliographies (131)
 EDRS PRICE MF01/PC03 Plus Postage.
 DESCRIPTORS *Adolescents; Cardiovascular System; *Child Health; *Children; Eating Habits; *Heart Disorders; *Obesity; *Physical Health
 IDENTIFIERS AGRICOLA; *National Agricultural Library DC

ABSTRACT

This bibliography consists of 212 recent citations (January 1985 through May 1990) from AGRICOLA, the National Agricultural Library (NAL) computerized database. The bibliography addresses issues concerning childhood obesity and cardiovascular disease. Each citation includes the NAL call number, the title, the author(s) the city of publication, the name and date of the journal in which it is published (if applicable), page numbers, language, descriptors, and abstract. The citations are listed in alphabetical order. An author index is included. (TE)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *


Childhood Obesity and Cardiovascular Disease

January 1985 - May 1990

Quick Bibliography Series: QB 90-59
Updates SRB 88-06 and SRB 88-05

212 Citations from AGRICOLA

Natalie A. Updegrove
Food and Nutrition Information Center



**Food &
Nutrition**

National Agricultural Library Cataloging Record:

Updegrave, Natalie A.

Childhood obesity and cardiovascular disease.

(Quick bibliography series ; 90-59)

1. Obesity in children – Bibliography. 2. Heart – Diseases – Bibliography. I. Title.
aZ5071.N3 no.90-59



DOCUMENT DELIVERY SERVICES TO INDIVIDUALS

The National Agricultural Library (NAL) has a unique responsibility to supply copies of agricultural materials not found elsewhere.

Filling requests for materials readily available from other sources would divert NAL's resources and diminish its ability to serve as a national source for agricultural and agriculturally related materials. Therefore, NAL should be viewed as a library of last resort and individuals should submit requests first to local or state sources prior to sending to NAL. In the United States, possible sources are public libraries, land-grant university or other large research libraries within a state. Individuals in other countries should submit requests through major university, national, or provincial institutions.

If the needed publications are not available from these sources, the requests may be submitted to NAL with a statement indicating their non-availability. Submit one request per page following the instructions for libraries below.

NAL's DOCUMENT DELIVERY SERVICE INFORMATION FOR THE LIBRARY

The following information is provided to assist your librarian in obtaining the required materials.

LOAN SERVICE – Materials in NAL's collection are loaned only to other U.S. libraries. Request for loans should be made through local public, academic, or special libraries.

The following materials are not available for loan: serials (except USDA serials); rare, reference, and reserve books; microforms; and proceedings of conferences or symposia. Photocopy or microform of non-circulating publications may be purchased as described below.

DOCUMENT DELIVERY SERVICE – Photocopies of articles are available for a fee. Requests must be made through local public, academic, or special libraries. The library should submit a separate interlibrary loan form for each article or item requested. If the citation is from an NAL database (CAIN/AGRICOLA, *Bibliography of Agriculture*, or the NAL Catalog) and the call number is given, that call number should be listed in the proper block on the request form. Willingness to pay charges must be indicated on the form. Compliance with copyright law or a statement that the article is for "research purposes only" must be included on the interlibrary loan form. Requests cannot be processed without these statements.

Charges:

- Photocopy, hard copy of microfilm and microfiche – \$5.00 for the first 10 pages or fraction copied from a single article or publication. \$3.00 for each additional 10 pages or fraction.
- Duplication of NAL-owned microfilm – \$10.00 per reel.
- Duplication of NAL-owned microfiche – \$ 5.00 for the first fiche and \$.50 for each additional fiche per title

Billing – Charges include postage and handling, and are subject to change. Invoices are issued quarterly by the National Technical Information Service (NTIS), 5285 Port Royal Road, Springfield, VA 22161. Requesters are encouraged to establish deposit accounts with NTIS. **DO NOT SEND PREPAYMENT.**

Send Requests to: USDA, National Agricultural Library
Lending Branch, ILL, 6th Floor
Beltsville, Maryland 20705-2351

Questions concerning these services may be made by correspondence to Head, Lending Branch or by telephone (301) 344-3755.

AGRICOLA

Citations in this bibliography were
entered in the AGRICOLA
database between January 1979
and the present.

CHILDHOOD OBESITY AND CARDIOVASCULAR DISEASE

SEARCH STRATEGY

S1	10679	CHILD?/TI,DE
S2	4984	INFAN?/TI,DE
S3	2789	ADOLESCEN?/TI,DE
S4	582	TEEN?/TI,DE
S5	16501	(CHILD?/TI,DE OR INFAN?/TI,DE OR ADOLESCEN?/TI,DE OR TEEN?/TI,DE)
S6	3377	OBES?/TI,DE
S7	11512	WEIGHT/TI,DE
S8	75523	CONTROL/TI,DE
S9	687	WEIGHT/TI,DE(W) CONTROL/TI,DE
S10	11512	WEIGHT/TI,DE
S11	4320	LOSS/TI,DE
S12	990	WEIGHT/TI,DE(W) LOSS/TI,DE
S13	224	OVERWEIGHT/TI,DE
S14	4388	OBES?/TI,DE OR WEIGHT() CONTROL/TI,DE OR WEIGHT() LOSS/TI,DE OR OVERWEIGHT/TI,DE
S15	383766	PY=1985 : PY=1990
S16	36776	SF=FNC (FOOD AND NUTRITION)
S17	135	S5 AND S14 AND PY=1985:1990 AND SF=FNC
S18	1666	CARDIOVASCULAR/TI,DE
S19	2994	CHOLESTEROL/TI,DE
S20	2966	HEART/TI,DE
S21	6940	FAT/TI,DE
S22	13049	CARDIOVASCULAR/TI,DE OR CHOLESTEROL/TI,DE OR HEART/TI,DE OR FAT/TI,DE
S23	873	HYPERTENSION/TI,DE
S24	16202	HIGH/TI,DE
S25	14186	BLOOD/TI,DE
S26	4262	PRESSURE/TI,DE
S27	56	HIGH/TI,DE(W) BLOOD/TI,DE(W) PRESSURE/TI,DE
S28	544	CORONARY/TI,DE
S29	471	ATHEROSCLEROSIS/TI,DE
S30	1023	LIPOPROTEIN/TI,DE
S31	2746	HYPERTENSION/TI,DE OR HIGH() BLOOD() PRESSURE/TI,DE OR CORONARY/TI,DE OR ATHEROSCLEROSIS/TI,DE OR LIPOPROTEIN/TI,DE
S32	14537	S31 OR S22
S33	36776	SF=FNC (FOOD AND NUTRITION)
S34	383766	PY=1985 : PY=1990
S35	1542	S32 AND SF=FNC AND PY=1985:1990
S36	1651	S35 OR S17
S37	434	DT=INSTRUCTIONAL
S38	129	DT=KIT
S39	1199	DT=MEDIA
S40	562	DT=MOTION
S41	21	DT=POSTER
S42	1671	DT=INSTRUCTIONAL OR DT=KIT OR DT=MEDIA OR DT=MOTION OR DT=POSTER

S43	71	DT=OTHER ACCOMPANYING MATERIAL
S44	8	DT=POSTERS
S45	243	DT=RECORDING
S46	452	DT=SLIDE
S47	765	DT=OTHER ACCOMPANYING MATERIAL OR DT=POSTERS OR DT=RECORDING OR DT=SLIDE
S48	1	DT=SLIDES
S49	23	DT=TRANSPARENCY
S50	1	DT=VIDEO
S51	41	DT=VIDEOCASSETTE
S52	66	DT=SLIDES OR DT=TRANSPARENCY OR DT=VIDEO OR DT=VIDEOCASSETTE
S53	648	DT=VIDEORECORDING
S54	1	DT=VIDEOTAPE
S55	1216	DT=AV
S56	30	DT=CHART
S57	380	DT=FILMSTRIP
S58	1513	DT=VIDEORECORDING OR DT=VIDEOTAPE OR DT=AV OR DT=CHART OR DT=FILMSTRIP
S59	2	DT=FLIP CHART
S60	23	DT=GAME
S61	25	DT=FLIP CHART OR DT=GAME
S62	2559	S42 OR S47 OR S52 OR S58 OR S61
S63	36776	SF=FNC (FOOD AND NUTRITION)
S64	1695	S62 AND SF=FNC
S65	36	S64 AND S36
S66	1615	S36 NOT S65
S67	156	S35 AND S5
S68	265	S67 OR S17
S69	7	S68 AND S64
S70	258	S68 NOT S69

0001

Accuracy of self-reports of food intake in obese and normal-weight individuals: effects of parental obesity on reports of children's dietary intake.

AJCN Klesges, R.C., Hanson, C.L., Eck, L.H., Durff, A.C. Baltimore, Md. American Society for Clinical Nutrition. American Journal of Clinical Nutrition. Nov 1988, v. 48 (5) p. 1252-1256. charts. Includes 33 references. (NAL Call No. DNAL FNC 389.8 J824).
Abstract: The purpose of this investigation was to determine the accuracy of dietary-intake information of normal-weight vs overweight parents in their reports of their children's food intake. Subjects were 36 families with either normal-weight (n = 20) or overweight (n = 16) children aged 4-9 y. Unobtrusive observers recorded the types and amounts of food eaten by the children at one meal. The next day families were visited in their homes and were asked to provide dietary-intake information from the previous day on their child. Results indicated that parental reports of the dietary intake of their children correlated significantly with the observers' measures of food intake. Neither the father's, the mother's, nor the family's obesity status had an effect on the accuracy of recalled information. The results indicated that the lack of differences consistently observed in dietary intake between obese and normal-weight children could not be explained by differential accuracy of recalled dietary information.

0002

Adipogenic activity in sera from obese children before and after weight reduction

AJCN Hauner, H., Wabitsch, M., Zwiauer, K., Widnaim, K., Pfeiffer, E.F. Baltimore, Md. American Society for Clinical Nutrition. American Journal of Clinical Nutrition. July 1989, v. 50 (1) p. 63-67. charts. Includes 24 references. (NAL Call No. DNAL FNC 389.8 J824).
Abstract: The adipogenic and mitogenic potentials of sera obtained from obese children before and after weight reduction and from lean control subjects were studied in clonal 3T3-L1 fibroblasts. The sera from the lean (n = 14) and obese (n = 12) children under habitual diet contained similar adipogenic activity. However, when the obese children underwent a weight-reduction program for 3 wk (600 kcal/d), the potential of their sera to stimulate glycerol-3-phosphate dehydrogenase, an index of adipogenic activity, was significantly reduced by 32% (p less than 0.01). Similarly, the mitogenic activity of these sera decreased significantly (202 +/- 15 vs 231 +/- 27 micrograms per dish, p less than 0.01). Testing pooled sera from the different groups in cultured rat adipocyte precursor cells gave similar results. This study suggests that human childhood-onset obesity is not mediated by increased circulatory

factors involved in the formation of new fat cells. The adipogenic and mitogenic activity of sera from obese children may be influenced by long-term dietary restriction.

0003

Adolescent nutrition issues and challenges.

DACDA Rosemont, Ill. National Dairy Council Dairy Council digest Literature Review July/Aug 1987 v. 58 (4) p. 19-24. Includes 104 references. (NAL Call No. DNAL FNC 389.8 D14).
Abstract: A detailed review summarizes and discusses various key aspects concerning adolescent nutrition. Specific topics include nutrient requirements of adolescents, the nutritional status of adolescents and principal nutrient deficiencies; the food habits of teenagers; and several nutrition-related issues. These issues cover weight control and eating disorders, special nutrient requirements of adolescents involved in high school athletics, and the nutrient needs of pregnant adolescents and the effect of nutritional status on pregnancy outcome. (wz)

0004

Adolescent obesity.

Bray, G.A. Rockville, Md. Aspen Systems Corp., 1986, c1985. Frontiers in clinical nutrition / edited by Norman Kretzmer. Literature review. p. 153-183. ill., charts. Includes 70 references. (NAL Call No. DNAL RM217.F76 1986).
Abstract: A clinical review details 3 classifications of obesity, viz: (1) an anatomical classification based on adipocyte number and their body distribution, (2) an etiological classification based on specific identifiable factors; and (3) a developmental classification based on obesity onset age and the natural history of obesity in individuals (considering features in childhood, adulthood, and pregnancy). Factors detailed under classification (2) include genetic, nutritional, restricted physical activity, endocrine alterations, hypothalamic, pharmacological, socioeconomic, and psychological factors. Data gleaned from both animal and human studies are discussed. (wz)

0005

Adolescent obesity intervention: validation of the SHAPEDOWN program.

JADA Mellin, L.M.; Sinkard, L.A.; Irwin, C.E., Jr. Chicago, Ill.: The Association. Journal of the American Dietetic Association. Mar 1987, v. 87 (3), p. 333-338. charts. Includes 35 references. (NAL Call No.: DNAL FNC 389.8 AM34).
Abstract: Effectiveness and transferability of the adolescent obesity intervention SHAPEDOWN were evaluated through a randomized experimental design study of 66 adolescents at four diverse clinical sites in California. Participation in

the program was associated with significant improvement in relative weight, weight-related behavior, depression, and knowledge of weight management concepts at post-treatment and at 1-year follow-up (author)

0006

Adoption study of obesity.

NEJMAG Bonds, D.R. ; Crosby, L.D. Boston, Mass. Massachusetts Medical Society. New England Journal of Medicine. July 10, 1986. v. 315 (2) p. 128-130. charts. Includes 11 references (NAL Call No. DNAL FNC 448.8 N442) Abstract Five letters to the editor critique a recently published paper that concluded that obesity in offspring was more strongly associated with maternal than paternal size, and that a significant trend existed between the body-mass index of adopted children and that of their biological parents but not of their adoptive parents. A sixth letter to the editor by the authors of the paper provides a defense and commentary in response to these 5 letters (wz)

0007

Adult heart disease prevention in childhood: a national survey of pediatricians' practices and attitudes.

PEDIAU. Nader, P.R. ; Taras, H.L.-Sallis, J.F.-Patterson, T.L. Elk Grove Village, Ill. ; American Academy of Pediatrics. Pediatrics. June 1987. v. 79 (6). p. 843-850. ill., charts. Includes 47 references. (NAL Call No. DNAL FNC RJ1.P42) Extract: There is controversy about the role of pediatric care givers in reducing presumed risk factors for cardiovascular diseases in children. A national mail survey of 2,000 pediatricians was designed to determine the attitudes, current practices, and knowledge among primary care pediatricians regarding these risks. The response rate was 60% (779 primary care pediatricians). Responses indicated that a majority of pediatricians take a family history of cardiovascular diseases, assess BP, recommend exercise to school-aged children, and advise patients and parents against smoking. Few pediatricians felt confident in their ability to affect change in patient life-styles. There was a relatively low level of provision of dietary advice, and most pediatricians do not measure serum cholesterol levels, except in high-risk older children. A substantial minority do not discuss smoking, even with adolescents. Practices and attitudes varied with the age of the patient. Obesity was the topic most frequently chosen for continuing medical education. Older pediatricians were most likely to advocate and practice risk reduction in children. Attitudes and reported practices also varied by gender and geographic location of the pediatrician as well as the ethnic composition of the practice. Implications for continuing medical education are discussed.(author)

0008

Aerobic capacity, obesity, and atherosclerotic risk factors in male adolescents.

PEDIAU Fripp, Raymond R. , Hodgson, James L.-Kwiterovich, Peter O.-Werner, John C -Schuler, H Gregg-Whitman, Victor Elk Grove Village, Ill American Academy of Pediatrics. Pediatrics. May 1985. v. 75 (5). p. 812-818. charts. Includes 19 references. (NAL Call No. DNAL FNC RJ1.P42) Extract Correlations between aerobic capacity, obesity, and atherosclerotic risks factors were evaluated in adolescents with low-to-moderate levels of physical fitness. Subjects with higher levels of fitness had a more favorable risk profile with decreased body mass index, lower systolic and diastolic blood pressure and plasma triglyceride levels, and higher plasma high-density lipoprotein-cholesterol levels. Simple linear regression analysis revealed an association between body mass index and blood pressure, plasma triglyceride and plasma high-density lipoprotein-cholesterol. The level of aerobic fitness as determined by exercise duration was also associated with the same atherosclerotic risk factors. However, multiple linear regression analysis demonstrated that body mass index provided the largest explanation, by those variables examined, of the interindividual variance in blood pressure, plasma triglyceride, and high-density lipoprotein-cholesterol. Aerobic fitness contributed only minimally to the variation in these risk factors. These findings suggest that if aerobic conditioning is used to modify atherosclerotic risk factors, it should be accompanied by a reduction in weight in adolescents with low-to-moderate levels of physical fitness. (Author)

0009

Anthropometric comparisons among low-income Mexican-American children aged 10 through 14.

JADAA. Guinn, B. ; Semper, T.-Crofts, A. Chicago, Ill. ; The Association. Journal of the American Dietetic Association. Jan 1986. v. 86 (1). p. 57-60. ill., charts. Includes 13 references. (NAL Call No. : DNAL FNC 389.8 AM34). Extract: Skinfold measurements of 795 Mexican-American children 10 through 14 years of age were compared with National Center for Health Statistics data. Analysis revealed significantly greater levels of subcutaneous fat in the children in the sample. The results indicated a high degree of overnutrition among the Mexican-American children studied.(author)

0010

Anthropometric measurements and dietary intakes of Cherokee Indian teenagers in North Carolina.

JADAA. Story, M. ; Tompkins, R.A.-Bass, M.A.-Wakefield, L.M. Chicago, Ill. ; The Association. Journal of the American Dietetic Association. Nov 1986. v. 86 (11). p. 1555-1560. ill., charts.

includes 33 references (NAL Call No DNAL FNC 389.8 AM34)
Extract Anthropometric measurements, rates of obesity, and food intake practices were investigated among 277 Cherokee Indian youths in North Carolina. Mean body weights and triceps skinfold measurements of the Cherokees were significantly higher than national reference data. However, there were no significant differences between height data for the Cherokees and national survey data. Obesity rates were found to be high, but mean energy intakes were not significantly different for lean and fat individuals. It does not appear that the obesity is caused by overeating (author)

0011

Apolipoproteins A-I and B in obese children.

JPGND6 Sveger, T., Flodmark, C.E., Fex, G., Henningsen, N.C. New York, N.Y. Raven Press. Journal of pediatric gastroenterology and nutrition. Nov 1989, v. 9 (4), p. 497-501. charts. Includes 18 references (NAL Call No DNAL RJ446 J68)

Abstract The purpose of this study was to determine whether obese children have abnormal apolipoprotein A-I or B concentrations and if these levels are related to obesity, physical fitness or birthweight. The results were discussed in relationship to developing coronary heart disease as adults.

0012

Assessing knowledge of cardiovascular health-related diet and exercise behaviors in Anglo- and Mexican-Americans.

Vega, W.A.; Sallis, J.F.; Patterson, T.; Rupp, J.; Atkins, C.; Nader, P.R. Duluth, Minn. Academic Press. Preventive medicine. Sept 1987, v. 16 (5), p. 696-709. ill., charts. Includes 16 references. (NAL Call No. DNAL RA421.P684).

Abstract The objective of this work was to develop scales to measure knowledge of health behaviors related to cardiovascular disease among children and adults and to evaluate the psychometric (reliability scale for assessing knowledge) scale; and to investigate Anglo and Mexican American differences in health behavior knowledge on dietary sodium, dietary fat, and exercise and to identify determinants, such as sex and education, self confidence, and cultural orientation, of such knowledge. Test differences in knowledge by ethnicity and sex indicate strong effects for ethnicity among both adults and children; sex did not appear consistently related to knowledge, except males tend to be more knowledgeable about exercise. Education appeared to be the strongest determinant for Anglo-Americans and cultural orientation strongest for Mexican-American adults. It was concluded that the Health Behavior Knowledge Scales were found to be useful in assessing levels of cardiovascular knowledge in Anglo and Mexican

Americans

0013

Assessing needs for youth health promotion

Perry, C.L., Griffin, G., Murray, D.M. Orlando, Fla. Academic Press. Preventive medicine. May 1985, v. 14 (3), p. 379-393. charts. Includes 11 references (NAL Call No DNAL RA421.P684)

Abstract The activities and design of an on-going project in youth heart health promotion are described, based on educational interventions. Objectives are aimed at reductions in cigarette smoking, improved nutrition and exercise patterns, and hypertension prevention. A behavioral needs assessment, the results from 2 communities and their association with gender and age, and the implication of the results for developing intervention activities, are discussed. The limitations of this approach also are described. (wz)

0014

Assessment of body fitness in childhood obesity: evaluation of laboratory and anthropometric techniques.

JADAA, Bandini, L.G.; Dietz, W.H., Jr. Chicago, Ill. The Association. Journal of the American Dietetic Association. Oct 1987, v. 87 (10), p. 1344-1350. charts, forms. Includes 39 references. (NAL Call No. DNAL FNC 389.8 AM34).

Extract The identification of obesity as a pathological diagnosis depends on an accurate assessment of body fatness and a correlation of fatness with pathological consequences. Because total body fat varies with body weight; the proportion of body weight that is fat is probably a more reliable indicator of risk. Among obese children and adolescents, several problems have hindered the development of accurate clinical measures of percent body fat and total body fat. First, the use of direct methods to measure body composition is limited by expense and labor. Second, the relationship between anthropometric indexes and body composition in obese children and adolescents has not been intensively studied. Third, sample sizes of normal weight children have been too small to permit the development of diagnostic criteria. Fourth, the triceps skinfold is less reproducible in overweight subjects. Increases in lean body mass in obese adolescents may confound the use of body mass index as a measure of adiposity. Current laboratory methods for the measurement of body composition include: (1) underwater weighing, (2) 40K counting, (3) isotopic dilution measures, (4) neutron activation, and (5) electrical impedance. This article examines relationships between those methods and anthropometry in the measurement of fatness in children and adolescents, as well as the difficulties in measuring body fatness and the importance of body fat distribution and its relationship to morbidity in children. Current evidence suggests an association of morbidity and upper

segment obesity in adults. Corresponding studies in children and adolescents are yet to be carried out (author)

0015

Association of overweight with serum uric acid concentration in Japanese schoolchildren

NURIS, Iton, R., Oka, J., Echizen, H., Murakami, K., Aeba, H. Stoneham, Mass. Butterworth Nutrition reports international Oct 1988 v. 38 (4) p. 773-781. charts. Includes 11 references (NAL Call No. DNAL RC620.A1N8). Abstract: The association of overweight with serum uric acid concentration was explored in healthy 725 schoolchildren aged 9-18 in Tsumakoi, Japan. Rohrer's index, a measure of overweight, was significantly and positively correlated with serum uric acid concentration in 13-18-yr-old male subjects and 9-18-yr-old female subjects, even after adjusting for age. The mean serum uric acid level of the subjects with the values of Rohrer's index above the sex-age-specific 90th percentile for Japanese children was significantly higher than the other subjects, with the exception of the males of 9-12-yr-old age group. These results suggest that overweight relates with serum uric acid level in Japanese male children aged 13 and over and female children aged 9 and over.

0016

Attitudes, perceptions, and practices of Canadian school children towards obesity.

LeBow, M.J. New York, N.Y. Human Sciences Press Journal of obesity and weight regulation Spring 1988, v. 7 (1) p. 43-55. Includes 41 references (NAL Call No. DNAL RC628.O222). Abstract: The authors questioned over 300 school children about their attitudes toward overweight peers. The overweight children's beliefs were included. The possible reasons for obesity in children were also discussed.

0017

Basal metabolism of obese adolescents: age, gender and body composition effects.

IJOBOP, Katch, V., Rocchini, A., Becque, D., Marks, C., Moorehead, K. London John Libbey & Company. International journal of obesity, 1985, v. 9 (1), p. 69-76. charts. Includes 26 references. (NAL Call No.: DNAL RC628.A102). Abstract: Basal metabolic rate and body density measurements on 67 obese adolescents (37 girls, 30 boys; ages 10.4-16.2) were examined to assess their possible association with sex and age differences. Sex differences did not correlate with body composition or metabolic variables, and correlations between age, body composition, and basal energy expenditure (BEE) were not significant. Lean body mass did not correlate with BEE, suggesting that a "metabolic condition" may exist in obese adolescents that is modulated by excess fat or an unstable body cell mass. Statistical analysis indicated that

basal daily kcal may be best predicted by body surface area and per cent fat for obese adolescent boys, and by body surface area and body weight for obese adolescent girls (wz)

0018

Behavioral and psychological traits of weight-conscious teenagers: a comparison of eating-disordered patients and high- and low-risk groups.

Mallick, M.J., Whipple, T.W., Huerta, E. San Diego, Calif.: Libra Publishers Adolescence, Spring 1987, v. 22 (85), p. 157-166. ill., charts. Includes 19 references. (NAL Call No. DNAL H0793.A44).

Abstract: A study compared the behavioral and psychological traits of adolescent girls having eating disorders (ED) with those of adolescent female athletes and high school students. While the athletes had the highest self-concepts, the ED girls had the lowest self-image scores and scored very low on emotional tone and social relationships scores, suggesting an association between the presence of eating disorders and low self-image. (wz)

0019

Benefits of school-based, family-oriented cardiovascular risk screening and intervention.

Santanello, N.C.; Meyerson, D.A. St. Louis, Mo.: C.V. Mosby Company. Clinical nutrition, Nov/Dec 1988, v. 7 (6), p. 242-245. Includes 13 references. (NAL Call No. DNAL RM216.M342). Abstract: How to attract high-risk young and middle-aged adults into cardiovascular risk screening and intervention programs, and who in the pediatric population to screen, are important and only partially answered questions. This study demonstrates that a school-based, family-oriented program can attract into the health care system a significant number of high-risk adults who otherwise might not be identified. Additionally, screening only those children with a positive family history would have failed to identify almost two thirds of the higher-risk children in our study. A more widespread recognition among the general medical community of the familial nature of cardiovascular disease and of the possibilities for risk modification of entire family units is urged.

0020

Beyond baby fat weight-loss plans for children and teenagers /Frances Sheridan Goulart; foreword by Platon J. Collipp.

---. Goulart, Frances Sheridan. New York: McGraw-Hill, c1985. Includes indexes. xvii, 188 p.: ill.; 22 cm. Bibliography, p. 174-176. (NAL Call No. DNAL RJ399.C6G6).

Abstract: Obese children do not necessarily outgrow their baby fat. Strategies for combatting overweight, from infancy through adolescence with diet and exercise programs are offered. The many possible causes of overweight are discussed: heredity, stress, excess

sugar, metabolic disorders, food allergies, nutritional deficiencies, and illness. Five diet plans to counter these causes are presented, including menus, recipes, junk food alternatives, and eating out advice. Also included are a list of weight control classes, clinics and camps, mail-order sources of foods, and a reference list. (as)

0021

Blood lipids as related to food intake, body composition and cardiorespiratory efficiency in preschool children.
JEGND6 Parizkova, J. Mackova, J. Mackova, J. Stopkova, M. New York, Raven Press. Journal of pediatric gastroenterology and nutrition, Mar/Apr 1987, v. 21, p. 295-298. charts. Includes 13 references. (NAL Call No. DNAL RJ446 J68).
Extract: A group of 3-5-year-old children ($n = 22$) with a level of somatic development and physical fitness (modified step test) that corresponded to a previous measured representative sample was studied. The depot fat composition was 10% ± 4% and obesity prevalence (lean body mass (LBM) was 6.5 ± 0.6 kg. As in previous studies the fat intake was higher as compared to recommended allowances. The blood cholesterol level was 4.9 ± 0.8, high density lipoproteins (HDL) 1.2 ± 0.2, low density lipoproteins (LDL) 3.6 ± 0.8, triglycerides 0.6 ± 0.2 mmol/L and creatine kinase (CK) 42.2 ± 14.4 U/L. The step test index (STI) was 92 ± 9 and the cardiac efficiency index (CEI) was 0.575 ± 0.096. The sex differences were not significant, except for body weight. Marked variability was found in 2 characteristics measured. There were no significant relationships among somatic development, body composition, food intake, STI, CEI, cholesterol, and CK. The LBM/100 height index correlated significantly with CEI, CK, and fat intake. (author)

0022

Blood pressure, body fat, and dehydroepiandrosterone sulfate variation in adolescence.
Katz, S.H., Hediger, M.L., Zemel, B.S., Parks, C.S. Dallas. American Heart Association Hypertension, Apr 1986, v. 5, 14, p. 277-284. charts. Includes 52 references. (NAL Call No. DNAL KCB:5.H8H9).
Abstract: A cross-sectional study of black male and female adolescents (ages 12-16) assessed whether serum dehydroepiandrosterone sulfate (DHEAS) levels were associated with blood pressure variation independent of the established association between increased DHEAS levels and measures of excess body weight and obesity. High DHEAS levels were associated with higher blood pressure in the girls, even after adjusting for the effect of body fat on DHEAS level. In the boys, high serum DHEAS was associated with a low body mass index and higher blood pressure. Related findings also are discussed. The data are statistically summarized and evaluated. (wz)

0023

Body composition, adipocyte size, free fatty acid concentration, and glucose tolerance in children of diabetic pregnancies.
DIAEA Abbott, W.G.H., Tnuillez, P., Howard, B.V., Bennett, P.H., Salans, L.B., Cushman, S.K., Reaven, G.M., Foley, J.E. Alexandria, Va. American Diabetes Association, Diabetes, Oct 1986, v. 35, (10), p. 1077-1080. charts. Includes 33 references. (NAL Call No. DNAL FNC RC659.A1D5).
Extract: Previous studies show that children of women who are diabetic during pregnancy are more obese and have a higher prevalence of non-insulin-dependent diabetes mellitus (NIDDM) than children of women who first developed NIDDM more than 1 yr after the pregnancy (prediabetic mothers) and children of women who have never developed diabetes (nondiabetic mothers). To determine whether lean and obese children of glucose-intolerant pregnancies can be distinguished from similar children of glucose-tolerant pregnancies, we measured body composition, abdominal and gluteal adipocyte size, fasting free fatty acid (FFA), and fasting and stimulated glucose and insulin concentrations during an oral glucose tolerance test in prepubertal children of glucose-intolerant and prediabetic mothers. Each group ranged in adiposity from 6 to 40 per cent body fat. Age, weight, height, and percentage of body fat were similar in the two groups. There were no significant differences in adipocyte size or in glucose, FFA, C-peptide and insulin concentrations between the groups. The correlation between abdominal adipocyte size and fasting insulin concentration ($r = .91$ and 18 , $t = 2.8$, $P = .01$) was stronger in children from glucose-intolerant than from glucose-tolerant pregnancies, respectively. In terms of the parameters we measured, there are no major differences between children of glucose-intolerant and glucose-tolerant pregnancies. (author)

0024

Body composition in children: proposal for a method for calculating body fat percentage from total body density or skinfold-thickness measurements.
AJCNA, Weststrate, J.A., Deurenberg, P. Baltimore, Md. American Society for Clinical Nutrition, American journal of clinical nutrition, Nov 1989, v. 50, (5), p. 1104-1115. charts. Includes 60 references. (NAL Call No. DNAL FNC 389.8 JB24).
Abstract: A method is presented for assessing childhood obesity in a more objective way than most other routine methods used for diagnosing childhood obesity. The sum of bicipital, tricipital, subscapular, and suprailliacal skinfold thicknesses is related to total body density by use of theoretically defined prediction equations. Total body density is used to estimate total body fat percentage by use of age- and sex-dependent equations.

on the relation between body fat percentage and body density. These equations are constructed on the basis of published data on changes in the density of fat-free mass with age in children. With the proposed method childhood obesity can be assessed routinely in a more consistent way than with most other routine methods used to diagnose obesity in children. A preliminary validation study indicated that in children aged 7-10 y predicted body density differed on average less than 1% from measured body density. In addition, predicted body density was highly correlated (r greater than 0.7) with measured body density.

0025

Body composition of Peruvian children with short stature and high weight-for-height II Implications for the interpretation for weight-for-height as an indicator of nutritional status.
AJCNA Trowbridge, F.L. ; Marks, J.S. ; de Pomana, G.L. ; Madrid, S. ; Boutton, T.W. ; Klein, P.D. ; Betnesoa, M.
American Society for Clinical Nutrition
American journal of clinical nutrition
Sept 1987, v. 46 (3) p. 411-418, ill., charts. Includes 21 references. (NAL Call No.: DNAL FNC 389.8 J824)
Extract: In some child populations, low height-for-age, suggesting chronic undernutrition, may paradoxically be accompanied by relatively high weight-for-height, suggesting obesity. This growth pattern was investigated with anthropometric assessment and body composition studies using H₂(18)O stable isotope dilution in 139 preschool-age Peruvian children. Results suggested low height-for-age (15th percentile National Center for Health Statistics NCHS) and high weight-for-height (60th percentile NCHS). Skinfold thicknesses were lower whereas arm muscle areas were more similar to NCHS reference values. Total body water (as percent body weight) was greater than reference values, consistent with lower body fat. Differences in body proportions did not account adequately for the high weight-for-height. The data suggest that the high weight-for-height in these children is not obesity but is associated with lower body fat and greater lean tissue or lean tissue hydration that may reflect dietary, environmental, or genetic influences. Weight-for-height cutoffs for wasting or obesity may require different interpretations for different populations. (author)

0026

Body fat patterning and blood pressure in children and young adults--the Bogalusa heart study.
Shear, C.L. ; Freedman, D.S. ; Burke, G.L. ; Harsha, D.W. ; Berenson, G.S. Dallas : American Heart Association.
Hypertension, Mar 1987, v. 9 (3), p. 236-244, ill., charts. Includes 47 references. (NAL Call No.: DNAL RC685.H8H9).
Abstract: A clinical study assessed the relationship of body fat deposition to

blood pressure in 3784 children and young adults (ages 5-24) residing in a biracial community (Bogalusa, Louisiana). Central body fat (CBF) and peripheral body fat (PBF) correlated with systolic and diastolic (4th phase) blood pressure, respectively, after controlling for differences in age, sex, race, and height, however, PBF did not correlate with blood pressure after adjusting for CBF. On the other hand, CBF correlated with blood pressure after controlling for PBF. The CBF-blood pressure correlation was highest in young children and decreased with age in females to nonsignificance in adulthood, but remained high in both black and white men (wz).

0027

Body size, fatness, and leanness of Mexican American children in Brownsville, Texas: changes between 1972 and 1983.
Malina, R.M. ; Zavaleta, A.N. ; Little, B.B. Washington, D.C. : American Public Health Association, American journal of public health, May 1987, v. 77 (5), p. 573-577, ill., charts. Includes 18 references. (NAL Call No.: DNAL 449.9 AM30).
Extract: Changes in the height, weight, body mass index, triceps skinfold, and arm and estimated midarm muscle circumferences in lower socioeconomic Mexican American children, 6 through 17 years of age, from Brownsville, Texas, were documented on the basis of surveys done in 1972 and 1983. With the exception of height in youths ages 14-17, all parameters show gains at most ages, and in particular an increase in fatness. Brownsville Mexican American youth are similar in height, weight, and the body mass index to Mexican American youth in other areas of Texas. These trends confirm the large proportion of relatively short but heavy children among Mexican Americans. (author)

0028

Caffeine intakes of children from a biracial population: the Bogalusa Heart Study.
JADAA, Arbeit, M.L. ; Nicklas, T.A. ; Frank, G.C. ; Webber, L.S. ; Miner, M.H. ; Berenson, G.S. Chicago, Ill. : The Association, Journal of the American Dietetic Association, Apr 1988, v. 88 (4), p. 466-471, ill., charts. Includes 36 references. (NAL Call No.: DNAL FNC 389.8 AM34).
Abstract: To investigate caffeine intake patterns in children, dietary intakes were examined for a biracial sample of 1,284 infants and children. Twenty-four-hour dietary recalls were completed by parents of children aged 6 months and repeated at ages 1, 2, 3, and 4 years; children 10 years old served as their own respondents and were surveyed at ages 13, 15, and 17 years. The sample was 60% white and 40% black. Additional cohorts of 10-year-olds (no. = 686) were studied for temporal trend. Whites consumed significantly more caffeine than blacks as early as 1 year and persisted at a higher intake level from

2 to 17 years. This trend continued whether intake was measured in total milligrams, milligrams per 1,000 kcal, or milligrams per kilogram body weight. Significant sex differences in caffeine intakes per 1,000 kcal occurred among 15- and 17-year-olds (girls greater than boys). Peak periods of consumption occurred at ages 2, 3, 13, and 17. Snacks contributed large quantities of caffeine, particularly for 10-year-olds. Most frequent sources of caffeine were regular carbonated beverages, chocolate-containing foods, and tea. Mean intakes of caffeine for 10-year-olds were consistent from 1973 to 1982. Those observations document caffeine intakes beginning early in life (author)

0029

Changes in biomedical and physical correlates in behavioral weight loss with retarded youths.

Rotatori, A.F. ; Fox, R.A.; Matson, J.; Mehta, S.; Baker, A.; Lopuch, W.R. New York: Human Sciences Press, Journal of obesity and weight regulation, Spring 1986, v. 5 (1), p. 17-27, 111, charts. Includes 40 references. (NAL Call No. DNAL RC628.0222).

Abstract: A 21-week study of 17 trainable, mentally retarded adolescents (11 overweight; 6 normal weight) assessed the beneficial effects of behavioral therapy (BT) for weight loss. The BT adolescents had greater body weight losses than non-BT adolescents who gained weight. The BT group had significant decreases in diastolic blood pressure, abdominal circumference, and tricep skinfolds. The benefits of BT for achieving weight loss in the mentally retarded are discussed (wz)

0030

Changes in body composition in adolescent boys.

Riumallo, J. ; Durnin, J.V.G.A. London: J. Libbey, Human nutrition: Food sciences and nutrition, Feb 1988, v. 42 (2), p. 107-112, charts. Includes 22 references. (NAL Call No.: DNAL TX341.H85).

Abstract: A longitudinal 3-year study assesses changes in body composition in 47 healthy Scottish children at 10 and 13 years of age. The results indicate that a rapid fat deposit period is followed by an almost 100% lean mass period corresponding to rapid growth. Environmental moderators of these typical changes also are discussed. (wz)

0031

Changes in height velocity of obese preadolescents during weight reduction.

AJDC. Dietz, W.H. Jr. ; Hartung, R. Chicago: American Medical Association, American Journal of Diseases of Children, July 1985, v. 139 (7), p. 705-707, 111., charts. Includes 15 references. (NAL Call No.: DNAL FNC 448.8 AM38).

Abstract: A retrospective study evaluated height velocity before and during weight reduction in a cohort of 19 obese preadolescents (14 girls, 5

boys) who experienced marked reductions in weight/height ratio under a balanced calorie-deficient diet (1.5-2.0 g protein/g body weight) over a period of about 10 months. The results indicated that even mildly energy restrictive diets may effect a reduction in linear growth velocity, emphasizing the importance of careful monitoring in obese children on mildly energy restrictive diets. (wz)

0032

Changes in skinfold thicknesses and body mass index in 171 children, initially 1 to 5 years of age: a 5 1/2-year follow-up study.

ADBEDS. Weststrate, J.A. ; Klaveren, H van-Deurenberg, P. Elmsford, N.Y.: Pergamon Press, Addictive behaviors, 1986, v. 10 (4), p. 313-321, charts. Includes 26 references. (NAL Call No. DNAL RC563.A33).

Abstract: A longitudinal study with a 5.5-year follow-up assessed changes in skinfold thickness (ST) and body mass index (BMI) in 171 children who were initially 1-5 years of age. Body fatness increase with age was associated with initially fatter girls, with children who had fatter mothers, and with children having a lower social class status, but parental fatness and social status were not additively related to changes in ST. Differences in changes in BMI were not found between sexes of the children. (wz)

0033

Characteristics of abnormal food-intake patterns in children with Prader-Willi syndrome and study of effects of naloxone.

AJCN. Zipf, W.B. ; Berntson, G.G. Bethesda, Md.: American Society for Clinical Nutrition, American Journal of Clinical Nutrition, Aug 1987, v. 46 (2), p. 277-281, 111., charts. Includes 36 references. (NAL Call No. DNAL FNC 389.8 J824).

Extract: Prader-Willi syndrome (PWS) is characterized by morbid obesity and abnormal appetite. It has been suggested that appetite is reduced by the administration of the opioid antagonist, naloxone. This has led to the hypothesis that appetite disturbance is a consequence of an abnormal hypothalamic response to appetite effects of endogenous opiates and opiate antagonist may be a useful treatment. To characterize the feeding patterns of PWS children and test this hypothesis, we administered an appetite test to 10 PWS children and 9 obese control children. We also examined the effects of naloxone on eating behavior of the children with PWS. While initial rate of eating did not differ, the PWS group showed a much delayed satiety resulting in a longer period of food intake. No difference in food intake was observed with naloxone (1.6 mg im, 30 min before the feeding test) treatment as compared with saline treatment. (author)

0034

The checkerboard cardiovascular curriculum: a culturally oriented program

Harris, M B Davis, S M -Ford, V L -Isc, H Ken* Ohio American School Health Association Journal of school health Mar 1988 v. 58 (3), p. 104-107. 111. Includes 11 references (NAL Call No. DNAL LB3401 J6)

Abstract: A technical report describes the results of a pilot study of a culturally oriented cardiovascular curriculum designed for 5th-grade children in rural New Mexico. Comprised mostly of Pueblo and Navajo US Indians and Hispanics. The curriculum was tested with 218 such students, incorporating educational materials, examples, and exercises of relevance to the cultures of these children. The study results found increases in knowledge concerning the cardiovascular system, the benefits of exercise, the relation of nutritional habits to obesity, the risks of tobacco usage, and the importance of behavioral habit changes. It is concluded that such programs may enhance promotion of a healthy lifestyle in children in minority ethnic groups (wz)

0035

Childhood obesity

Wolbers, M. New York, N.Y. Vegetarian Life & Times. Vegetarian times, Sept 1985 (97), p. 41-42, 44. 111 (NAL Call No. DNAL TX392 A1V44)

Abstract: Childhood obesity can have a devastating effect on the psyche, especially in a society that is oriented toward thinness. Experiences as an overweight child, even if thin in later years, can permanently color one's self-perception. To help parents avoid trapping their children into this pattern, the following tips are given: select food wisely, do not use food as a reward, do not forbid sweets, eat only at the table, do not mix eating with other distractions, do not insist on a clean plate, limit television watching. For the overweight child, stressing support rather than competition will help the child work through the problem (jd-b)

0036

Childhood obesity.

JOPDA, Korsch, B. St. Louis C.V. Mosby. The Journal of pediatrics, Aug 1986 v. 109 (2), p. 299-300. Includes 4 references. (NAL Call No. DNAL FNC RJ1 A453)

Abstract: Childhood obesity is a multifaceted problem with multiple causes and programs for treatment based on each possible cause or on varying combinations of causes. Factors related in varying degrees to obesity in childhood include overeating, reduced physical activity, genetic predisposition, cultural and socioeconomic background, emotional factors, and metabolic and endocrine conditions. There are important long-term public health implications for obese children, since many will remain obese during their adult lives.

Significant psychosocial effects also have been demonstrated in obese children. Results of research studies examining the social stigma of obesity and its effects on children are summarized. Cultural eating patterns and images of desirable body weight and shape have been shown to vary greatly among cultural groups. Personality assessments and self-esteem measurements of obese children have shown measurable personality disturbances in these children but no evidence of clinical psychosocial problems. Additional research is needed to increase our knowledge of childhood obesity, and to help dispel judgmental, prejudicial reactions and feelings toward obese individuals (aje)

0037

Childhood obesity /edited by Platon J Collipp

Collipp, Platon J., 1932- New York, NY: Warner Books, 1986. c1980. xii, 276 p. 111. 17 cm. Includes

Bibliographical references and index (NAL Call No. DNAL RJ399 C6C5)

Abstract: For children, obesity is a real handicap, undermining health and happiness and frequently continuing into adulthood. In this book, a wide-angle view of current knowledge, theory, prevention, diet, and psychological therapy for infant, child, and adolescent obesity by highly competent authorities in the field is presented. The role of heredity in obesity, fat storage, fat metabolism, programs for obese children, and the psychological causes and effects of obesity represent some of the topics addressed. Hypertension, drug therapy, and surgical treatment for adolescents are other topics reviewed. It is generally recognized that the earlier an obese child enters a program of diet and exercise, the better.

0038

Childhood obesity: a NICHD workshop report.

Blessing, P. Washington, D.C. Office of Human Development Services, Department of Health and Human Services. Children today, Sept/Oct 1986 v. 15 (5), p. 26-29. 111. (NAL Call No.: DNAL HV701.C51).

Abstract: How and why children become obese, the subject of a recent National Institute of Child Health and Human Development workshop, has been attributed to a range of things - from genes to watching television. The fact that obese children face greater risks of developing psychological/social problems, high blood pressure, heart disease, diabetes or even sleep apnea has prompted more intensive investigation into prevention/treatment programs. Overfeeding in infancy, hypnotic effects of watching television, environmental influences, and genetic predispositions may all play a role in childhood obesity. Discussions of prevention/treatment point to the fact that parental involvement is most important, as well as a high quality

weight reduction program. Information on obesity measurement is included (jd-b)

0039

Childhood obesity and self-esteem.

JDPDA Kaplan, K.M. ; Wadden, T.A. St. Louis C.V. Mosby, The Journal of pediatrics Aug 1986, v 109 (2), p 367-370 ill., charts. Includes 15 references (NAL Call No. DNAL FNC RJ1 A452)

Extract To explore the relationship between obesity and self-esteem, the Piers-Harris Self-Esteem Inventory was administered to black inner-city children (grades 4 through 12). Those with chronic illnesses or in special education were excluded, yielding 851 for the study. Body mass index (BMI) served to estimate adiposity, and Ten State Nutrition Survey (TSNS) data yielded reference growth curves BMI and BMI relative to TSNS deals were calculated for each child. Using analysis of variance to compare groups based on relative BMI, small differences in mean self-esteem scores were detected (3 to 4 units or one-third standard deviation). These statistically significant differences are unlikely to be clinically significant. The correlation between BMI and self-esteem was small BMI accounted for only 1% of the variance in self-esteem score. Moreover, all self-esteem scores fell within the normal range. Neither age nor sex affected the relationship. The consequences of childhood obesity may be less harmful than formerly assumed. Our findings may explain why the promise of enhanced self-esteem fails to motivate weight loss in many children (author)

0040

Childhood obesity: Family perspective.

Loader, Peter J. New York Van Nostrand Reinhold International journal of eating disorders, May 1985, v 4 (2) p 211-225. Includes 67 references (NAL Call No. DNAL FNC RA784.A115 F&N).

Extract Childhood obesity remains a poorly understood condition which is difficult to modify. Most investigations and explanations of obesity adopt a biological, psychological, or sociological perspective, with little attention being paid to the family. The work that has been done in this area tends to be based on clinical impressions rather than systematic enquiry, and to focus on the mother-child relationship rather than on the family as a whole. This latter approach, where the family is the basic unit of study, has proved useful in our understanding and treatment of several disorders that can be compared to obesity. This paper argues that the relative absence of family studies constitute a significant gap in our investigation of childhood obesity. Such studies need to be carried out, but they should also be part of a more holistic approach to the problem, which seeks to make links between the observations made at several different levels of enquiry. (author)

0041

Children's cognitive concepts of obesity: a developmental study.

IJDBDP. Wolfle, U.A. ; Farnier, S.C -Rogers, J.S. London : John Libbey & Company, International journal of obesity 1987, v. 11 (1), p 73-83. ill., charts. Includes 27 references. (NAL Call No. DNAL RC628.A102)

Abstract. A study, using a Piagetian theoretical framework involving 96 children divided into 3 age groups (4-5, 8-9, and 12-13 years old), assessed the development of children's concepts of causal factors in the development and reversibility of obesity, and compared the results to the development of physical causality. Only the youngest group understood body identity at higher levels than physical conservation, but the levels of understanding fatness causes and causes of other natural physical events were not different in the youngest group. On the other hand, physical causality was both understood and understood at a higher level than obesity causality among the 2 higher age groups. Differences in the understanding between body identity and obesity causality only were noted for the highest age group.

0042

Children's television: its effect on nutrition and cognitive development.

Palumbo, F.M. ; Dietz, W.H. Jr. Thorofare, NJ. Slack, Pediatric annals, Dec 1985, v. 14 (12), p. 793, 796-798, 800-801. ill., charts. Includes 15 references. (NAL Call No. DNAL FNC RJ1 P35).

Abstract Television's impact on the cognitive development and growth of the maturing child, combined with the pediatrician's role in moderating the effects and influencing television in general, comprise the topics of this article. Growth effects of television stem from effects upon nutrition, eating habits and subsequent obesity. Television advertising, use of food in programming for activities other than hunger satisfaction, and the scarcity of obese children/adults in food-related commercials convey messages affecting children's eating habits and nutritional status. Studies are cited that explored these relationships. Cognitive development as it relates sex, IQ, and specific areas of achievement is discussed, as well as political actions being taken by pediatricians and other child advocates in the legislative arena. Groups such as the Action for Children's Television (ACT) are working hard to improve children's broadcasting in a climate of federal deregulation.(jd-b)

0043

Cholesterol & children a parent's guide to giving children a future free of heart disease /Robert E. Kowalski ; foreword by Dennis M. Davidson ; preface by Paul Y. Qaqundah.

Kowalski, Robert E. New York : Harper & Row, c1988. Includes index. xvii, 302 p. ; 25 cm. Bibliography: p. 292-294. (NAL

Call No DNAL RJ426 AB2K6 1988)

Abstract This text offers a practical program for modifying family diets and eating habits in order to reduce blood cholesterol levels and thereby reduce the risk of heart disease. Intended to provide information to parents concerning methods of lowering cholesterol levels in children, this work includes data on cholesterol and heart disease, detailed analysis of the composition of many foods found in the typical American diet, practical methods for lowering fat and cholesterol intake, and recipes for low cholesterol foods.

0044

Cholesterol in youth unbending the twig.

Boston, Mass Dept of Continuing Education, Harvard Medical School The Harvard Medical School health letter May 1989 v 14 (7) p 2-4 charts Includes 2 references (NAL Call No DNAL R11 H3)

Abstract This article discusses cholesterol testing for children. Questions discussed include what age should testing begin, how often, what are normal values for LDL, HDL and total cholesterol for different age groups and how aggressive should treatment be. Guidelines and recommendations for managing cholesterol intake are included.

0045

Cholesterol levels in childhood as determinants of adult cholesterol levels.

NUREA New York, N.Y. Springer-Verlag New York Inc Nutrition reviews Feb 1989 v. 47 (2) p. 46-47 charts Includes 6 references. (NAL Call No DNAL FNC 389.8 N953)

Abstract Data indicating that plasma cholesterol levels in children are of value in assessing the risk of having high cholesterol levels in adulthood are discussed.

0046

Consumption patterns of critical fat sources among adolescents in 1977-1985.

NURHA, Prattala, R.; Rauhonen, O.-Rimpela, M. Elmsford, N.Y. Pergamon Press, Nutrition research, May 1986, v. 6 (5), p. 485-498, 111., charts. Includes 32 references. (NAL Call No. DNAL QP141.A1N88).

Extract: Five samples (N=2900-4300) of Finnish 12 to 18-year-old boys and girls participated in a postal survey on health habits in 1977, 1979, 1981, 1983 and 1985. The proportions of high-fat milk and butter users decreased during the follow-up time, but the socio-economic and regional differences in consumption patterns remained unchanged. Both in 1977 and 1985, children in eastern and northern provinces, and the children of farmers and lower socio-economic and educational groups, more frequently used butter and high-fat milk than children of higher social and educational groups living in urbanized areas. (author)

0047

Correlates and predictors of serum total cholesterol in adolescents aged 12-17 years: the National Health Examination Survey.

PHRPA Gillum, R.F. Washington, D.C. Public Health Service Public health reports May/June 1989, v 104 (3) p 256-265 charts Includes 36 references (NAL Call No DNAL 151 65 P96)

Abstract To examine correlates and childhood predictors of serum total cholesterol in adolescence, measures of growth, development, and obesity were related to serum total cholesterol levels of youths aged 12-17 years in the National Health Examination Survey. In this sample, drawn from the U.S. population, serum total cholesterol levels were negatively correlated with indicators of growth and maturation in males aged 12-14 years and positively correlated with overweight or obesity at all ages. All measured variables could account for less than 15 percent of cholesterol variation in males and less than 6 percent in females. In white males, indicators of levels of maturation, growth, and changes in body fatness measured 28-53 months earlier were significant predictors of serum total cholesterol in adolescence, explaining 13 percent of its variation. Despite significant associations, indicators of growth, sexual maturation, and obesity explained only a small fraction of serum cholesterol variation in adolescents.

0048

Culture versus biology: children's attitudes toward thinness and fatness.

PEDIAU Feldman, W.; Feldman, E.-Goodman, J.T. Elk Grove Village, Ill. American Academy of Pediatrics, Pediatrics, Feb 1988, v. 81 (2), p 190-194. Includes 20 references. (NAL Call No. DNAL FNC RJ1.P42).

Extract Many of the studies regarding children's acquisition of prevailing cultural concepts of physical attractiveness are flawed by small and unrepresentative samples, measurement instruments of questionable reliability and validity, and experimental designs that do not protect against bias. Additional studies in which these methodologic flaws are overcome must be carried out if we are to understand truly when and how cultural concepts of beauty are acquired. Nevertheless, the majority of the studies already done find that children acquire prevailing cultural values of beauty before adolescence and that thinness is desirable to girls considerably before puberty. It is suggested that the etiology of eating disorders and the reasons for their increasing prevalence will not be discovered by studying only clinical cases. We propose that those interested in this important health problem study children before adolescence in an attempt to learn how preoccupation with weight begins and why thinness is believed to be attractive. Once these are understood, a greater challenge will be the development and

testing of interventions--be they in the schools or using the media--which can effectively prevent this public health problem. When culture and biology clash, people may suffer. (author)

0049

Decreased prolactin secretion in childhood obesity

JOPDA AVRuskin, Theodore W., Pillai, Shashikala-Juan, Christina-Kleinberg, David L. St. Louis C.V. Mosby. The Journal of pediatrics March 1985 v 106 (3), p. 373-378 ill., charts Includes 22 references. (NAL Call No DNAL FNC RJ1.A453).
Extract: Twelve obese patients and 7 control subjects, age and sex matched, whose weights were greater than 200% of ideal weight and 100% of ideal body weight, respectively, underwent intravenous insulin and thyroid releasing hormone (TRH) tests. Serial prolactin growth hormone, insulin, blood sugar, cortisol, glucagon, thyrotropin stimulating hormone, thyroxine, and triiodothyronine were obtained by RIA. Obese patients showed no significant differences from controls in basal and nadir glucose, basal and peak glucagon, cortisol, and thyroid responses to both tests. Basal insulin levels were higher and peak growth hormone responses after insulin were lower in the obese group than in controls. Where as all control subjects had prolactin response to both tests, 5 of 12 obese patients had no responses to insulin. Obese patients had lower prolactin responses at 30 minutes after insulin, and lower prolactin responses at 60 minutes after TRH. Maximum prolactin responses after TRH were lower in obese patients. Maximum prolactin responses after insulin were lower in obese patients. Thus prolactin secretion in childhood obesity is decreased after both stimuli, but more so after IV insulin than TRH and suggests that, as in adult hypothalamic obesity, neuroendocrine regulation of prolactin release in obese children is impaired. (Author)

0050

Design and participation.

PEDIAU. Webber, L.S.; Frank, G.C.-Smock, C.G.-Freedman, D.S.-Berenson, G.S. Elk Grove Village, Ill. American Academy of Pediatrics. Pediatrics. Cardiovascular Risk Factors from Birth to 7 Years of Age: The Bogalusa Heart Study. Nov 1987. v. 80 (5,pt.2), p. 767-779, charts. Includes 33 references. (NAL Call No.: DNAL FNC RJ1.P42).
Extract: Cardiovascular risk factor variables were examined in a cohort of 440 infants from birth through 7 years of age. Anthropometric measures, BP, serum lipid and lipoprotein values, and dietary intake data were obtained according to detailed protocols. Various quality controls to ensure the collection of valid and reliable data were instituted. Participation remained high throughout the study with 80% of the children examined at 6 months and 60% at 7 years of age. Rates were

slightly higher for black than for white children. Children born in the private hospital were more likely to continue in the study than children born in the charity hospital. Children of parents examined when the child was 2 years of age were more likely to be examined during the preschool phases, but parental examination was not related to child examination when the child was 7 years of age. As in studies of school-aged children, measurement errors were lowest for height, weight, and serum total cholesterol. Measurement errors for BP were highest at the younger ages, particularly for diastolic BP. Examination of a newborn cohort throughout time affords the opportunity to study early development of relationships and tracking of cardiovascular risk factors. (author)

0051

Desire for thinness among high school cheerleaders: relationship to disordered eating and weight control behaviors.

Lundholm, J.K.; Littrell, J.M. San Diego, Calif. Libra Publishers. Adolescence. Fall 1986. v. 21 (63) p 573-579, charts. Includes 24 references. (NAL Call No.: DNAL HQ793.A44).
Extract: The eating disorders of anorexia nervosa and bulimia are increasing in frequency among adolescent females. These increases have been linked to the cultural ideal in American society of thinness. Attempting to control weight is one behavioral manifestation of the desire for thinness. One particular group of adolescents, female cheerleaders, often experience pressure to attain and maintain weight that is lower than other adolescents of the same height. This study examined cheerleaders' desire for thinness in relationship to disordered eating and weight control behaviors. A Desire for Thinness Scale and selected scales from three eating disorders instruments were administered to 751 high school cheerleaders from the Midwest. Cheerleaders who scored in the upper third on the Desire for Thinness Scale were compared with those who scored in the lower third. Cheerleaders who expressed a strong desire for thinness had significantly higher scores (p less than .0001) on 7 of 8 eating disorders scales. The greater the desire for thinness, the more likely the tendency to report disordered eating and weight control behaviors associated with bulimia. Implications from this study include an awareness of how a cultural ideal of thinness may indirectly increase disordered eating and weight control behaviors by making weight loss a salient goal. A proactive approach to modifying negative aspects of the cultural emphasis on thinness is proposed. (author)

0052

Detection and treatment of lipid and lipoprotein disorders of childhood
proceedings of the Third International Atherosclerosis Conference, held in Vienna, Austria, April 4-9, 1983

/editors Kurt Widhalm, Herbert K Naito

Widhalm, Kurt -Naito, Herbert K. New York: Liss, c1985. xi, 229 p. Includes bibliographies and index. (NAL Call No. DNAL RC692 I46-1933)

Abstract: A reference text for research workers and clinicians involved in the association of lipid metabolism disorders with cardiovascular disease mortality risk contains 16 authoritative technical papers presented at a 1983 international conference on atherosclerosis. These papers cover clinical screening methods, the effects of reducing serum cholesterol levels, and problems encountered in the long-term treatment of children having lipid and lipoprotein metabolism disorders. Detailed methods are covered for enhancing the reliability of laboratory data, and the latest methods for detecting lipid abnormalities are discussed. The current state of knowledge concerning new therapeutic strategies for treating these abnormalities is reviewed. Practical and applied information of use to clinicians in patient care is included. Data tabulations are presented throughout the text, and literature citations are appended to each page.

0053

Determinants of weight and adiposity in the first year of life

JOPDA Kramer, Michael S., Barr, Ronald G., Leduc, Denis G., Boisjoly, Christiane-McVey-White, Lynne-Pless I, Barry, St. Louis, C.V., Mosby, The Journal of Pediatrics. "Presented in part at the Annual Meeting of the Ambulatory Pediatric Association, Washington, D.C., May 5, 1983." Jan 1985, v. 106 (1), p. 10-14. charts. Includes 22 references. (NAL Call No. DNAL FNC RJ1 A453).

Extract: To overcome methodologic defects (failure to control for confounding factors, univariate statistical analyses) in previous studies of etiologic determinants of childhood adiposity, we carried out a prospective cohort study of 462 healthy, full-term infants observed from birth to 12 months. Postpartum, we obtained sociodemographic data and administered 2 recently validated scales of maternal attitudes toward feeding and infant body habitus. Parental heights and weights and infant feeding variables were determined by interview, and at 6 and 12 months we measured height and weight and triceps, subscapular, and suprailliac skinfolds. Multiple regression analysis was used to determine independently predictive factors for weight, body mass index (BMI equals weight/height squared), and the sum of the 3 skinfold measurements. Birth weight, sex, age at introduction of solids, and duration of breast-feeding were all significant predictors of weight at 12 months. Significant determinants for BMI included birth weight, duration of breast-feeding, sex, and IQ; those for total skinfold were age at introduction

of solid foods and birth weight. Similar results were obtained at 6 months, although slightly less of the variance was explained. We conclude that the ability to predict which babies will be heavy or obese during the first year is limited. Breast-feeding and delayed introduction of solid foods do offer some protective effect, however, and thus efforts to encourage these practices may be reaping some benefit. (author)

0054

Diet of young children and cardiovascular disease

BMJDAE Taitz, L.S. London: British Medical Association. British Medical Journal, Apr 11, 1987, v. 294 (6577), p. 920-921. Includes 16 references. (NAL Call No. DNAL 448.8 B77)

Abstract: A brief report addresses the question of whether the diet of young children should be altered to reduce cardiovascular disease risks. The validity of results of longitudinal studies supporting such a change is discussed. It is concluded that, until there is clearer evidence on the importance of early diet on later health, dietary advice for children under 5 years of age should not be radically changed. (wz)

0055

Dietary calcium intake in hypertension.

Morris, C.; McCarron, D.A. Dallas: American Heart Association. Hypertension, Sept 1987, v. 10 (3), p. 350-353. charts. Includes 40 references. (NAL Call No. DNAL RC685.H8H9)

Abstract: A letter to the editor discusses and critiques the results reported in a recent publication (using data from National Health and Nutrition Examination Surveys (NHANES) I and II) that found that low dietary calcium intake and hypertension were not related, and presents arguments that the evidence is remarkably consistent in demonstrating a weak, but plausible relationship between dietary calcium and blood pressure. A reply letter from authors of the recent publication provides a rebuttal of the arguments of the first letter, rejecting the conclusion of such a relationship, and stating that the data do not support the hypothesis of a negative association between dietary calcium intake and blood pressure and that this issue remains to be resolved in further needed studies. (mjs)

0056

Dietary habits in relation to tracking of cholesterol level in young adolescents: a nine-year follow-up.

NUMEB, Vobecky, J.S.; David, P.; Vobecky, J. Basel; S. Karger. Annals of Nutrition and Metabolism, Sept/Dec 1988, v. 32 (5-6), p. 312-323. charts. Includes 48 references. (NAL Call No.: DNAL RM214.N8)

Abstract: Because the origin of atherosclerosis is often in childhood, an early identification of future adults at risk can contribute to the prevention

of atherosclerosis. In a prospective study on nutrition and health, a cohort of 556 infants was followed from birth to 2 years. Among them, 116 were found at risk for hypercholesterolemia with a cholesterol level greater than or equal to 200 mg/dl. Nine years later, these children at risk and their controls were invited to a follow-up examination of blood lipids, of nutritional status, by anthropometric measurements and personal and family history. From 32.7 to 48.7% of children at risk remained with a high cholesterol level (greater than or equal to 190 mg/dl) in comparison with 15% of controls. The nutritional profile was evidently different, especially the food frequency. The results strongly suggest that the early identification of children at risk and the screening of other family members, siblings and parents for hypercholesterolemia can serve for a sound intervention on nutritional habits.

0057

Dietary sources of energy, protein, fat and fibre in 375 English adolescents
Mackett, A.F., Rugg-Gunn,

A.U., Appleton, D.R., Coombs, A. London, Eng. J. Libbey. Human nutrition applied nutrition, June 1986, v. 40A (3) p. 176-184. charts. Includes 21 references. (NAL Call No. DNAL TX341 H82).

Abstract. A 2-year dietary survey gathered data on the sources of energy, protein, fat, and fiber in the diet of 375 English adolescents (ages 11-14) for development of health education programs for this target population. The principal food source for energy was potatoes (chips, crisps), while that for protein and fat was "meat" (although other important sources of fat also were identified). Dietary fiber was found to be contributed by chips, white bread, crisps, and baked beans. Lower socioeconomic status adolescents appeared to have more non-nutritional food preferences. Differences between sex and social status are discussed. (wz)

0058

Dieting practices and attitudes of overweight, underweight, and ideal weight school girls.

Balentine, M.B.; Stitt, K.; Bonner, J.; Clark, L. Denver, Colo.; American School Food Service Association. School foodservice research review, Fall 1986, v. 10 (2) p. 98-103. ill., charts. Includes 17 references. (NAL Call No. DNAL TX945.S344).

Extract. The study was designed to determine the dieting practices and attitudes of overweight, underweight, and ideal weight adolescents in two school systems. Of the 560 girls, 74 and 26 percent were white and black, respectively. Although almost half were underweight according to height-weight tables, most had tried to diet to lose weight. The overweight adolescent appeared to be preoccupied with food and dieting. The underweight girl is more likely to have anorexia; whereas, the overweight student is more likely to

practice bulimia (author).

0059

Distributions of total cholesterol, triglycerides, and high-density lipoprotein cholesterol in Israeli Jewish children of different geographic-ethnic origins, ages 9-17 years.

Zahavi, I., Goldbourt,

U., Cohen-Mandelzweig, L., Katz, M., Appel, S., Harel, G., Sperling, Z., Lazarovici, M., Hart, J., Neufeld, H.N. Duluth, Minn.

Academic Press. Preventive medicine Jan 1987, v. 16 (1) p. 25-51. ill., charts. Includes 40 references. (NAL Call No. DNAL RA421 F684). Abstract. In this study, 1,153 schoolchildren from the area of Petah-Tikva, Israel, were analyzed for levels of plasma total cholesterol (TC), high density lipoprotein cholesterol (HDL-C), and triglycerides (TG). TC levels decreased in boys as age increased 10-11 to 14-15, and in girls became progressively lower until 12-13 and then gradually increased. TC levels were only partly accounted for by lower HDL-C levels at these ages. In boys, TG levels were inversely related to HDL-C levels. Overall, TC and HDL-C were higher among girls than boys beginning at ages 12-13, with little or no sex differences in TG. (lzp)

0060

Do we fatten our children at the television set? Obesity and television viewing in children and adolescents.

Pfiffner, Dietz, Jr.; William, H.; Gortmaker, Steven L. Elk Grove Village, Ill. American Academy of Pediatrics. Pediatrics, May 1985, v. 75 (5), p. 807-812. ill., charts. Includes 26 references. (NAL Call No. DNAL FNC RJ1.P42).

Extract. The association of television viewing and obesity in data collected during cycles II and III of the National Health Examination Survey was examined. Cycle II examined 6,965 children aged 6 to 11 years and cycle III examined 6,671 children aged 12 to 17 years. Includes in the cycle III sample were 2,153 subjects previously studied during cycle II. These surveys, therefore, provided two cross-sectional samples and one prospective sample. In all three samples, significant associations of the time spent watching television and the prevalence of obesity were observed. In 12- to 17-year-old adolescents, the prevalence of obesity increased by 2% for each additional hour of television viewed. The associations persisted when controlled for prior obesity, region, season, population density, race, socioeconomic class, and a variety of other family variables. The consistency, temporal sequence, strength, and specificity of these associations suggest that television viewing may cause obesity in at least some children and adolescents. The potential effects of obesity on activity and the consumption of calorically dense foods are consistent with this hypothesis. (Author)

0061

Does a vigorous feeding style influence early development of adiposity?

JOPDA Agras, W.S. ; Kraemer, M.C. -Berkowitz, R.I. -Korner, A.F. -Hammer, L.D. St. Louis, C.V. Mosby, The Journal of pediatrics May 1987 v. 110 (5) p. 799-804. 111 charts. Includes 31 references. (NAL Call No. DNAL FNC RJ1 A453)

Abstract: A prospective study of a cohort of healthy infants observed from birth to 2 years of age was carried out to investigate factors influencing the development of early adiposity. Infant suckling was measured in the laboratory twice during the first month of life. Multiple regression analyses revealed that parental educational level and a measure of feeding behavior, the interval between bursts of suckling, accounted for 18% of the variance in triceps skinfold measures at 1 year of age. A lower level of education and shorter interburst interval were associated with increased adiposity. Two feeding variables, pressure of suckling and the number of reported feeds per day, accounted for 21% of the variance in skinfold thickness at 2 years of age. Fewer, but larger, feeds and a higher sucking pressure were associated with a greater degree of adiposity. It seems that a vigorous infant feeding style, consisting of sucking more rapidly, at higher pressure, with a longer suck and burst duration, and a shorter interval between bursts of sucking, is associated with higher caloric intake and greater adiposity. The early development of this feeding style suggests that it may be a genetically endowed behavior. Breast-feeding protected against early adiposity only to the age of 6 months in this cohort of infants. (author)

0062

Does infant nutrition affect adiposity and cholesterol levels in the adult?

JPGND6. Hamosh, M. New York : Raven Press, Journal of pediatric gastroenterology and nutrition, Literature review, Jan/Feb 1988, v. 7 (1), p. 10-16. Includes 94 references. (NAL Call No. DNAL RJ446.J68).

Abstract: The effects of breast feeding during infancy on incidence of obesity and atherosclerosis in adult life are reviewed. The physiological differences between breast fed and bottle fed infants are discussed, particularly with respect to gastric hormones, cholesterol metabolism, and adipose tissue development and obesity. Implications are that genetics as well as postweaning nutrition may need to be closely examined to assess the relationship between breast feeding and incidence of obesity or atherosclerosis.

0063

A dynamic family approach for the prevention of cardiovascular disease.

JADAA, Nicklas, T.A. ; Johnson, C.C. -Arbeit, M.L. -Franklin, F.A. -Berenson, G.S. Chicago, Ill. The Association, Journal of the American Dietetic Association, Nov 1988, v. 88

(11) p. 1438-1440 charts. Includes 16 references. (NAL Call No. DNAL FNC 389.8 AM34)

Abstract: The Family Health Promotion Program draws on the strength of the family and the school to promote healthy life-styles among children with elevated cardiovascular risks.

0064

Early childhood diet, recommendations of pediatric health care providers.

JADAA, Taras, H.L. ; Nader, P.R. -Sallis, J.F. -Patterson, T.L. -Rupp, J.W. Chicago, Ill. The Association, Journal of the American Dietetic Association, Nov 1988 v. 88 (11) p. 1417-1421 charts. Includes 26 references. (NAL Call No. DNAL FNC 389.8 AM34)

Abstract: With the abundance of literature on cardiovascular disease (CVD) prevention during childhood, recommendations for restricted dietary sodium and fat intakes during infancy and childhood are both advocated for preventive health care and criticized because the safety is undetermined. Dietitians, nurse practitioners, and pediatricians were surveyed to determine what dietary recommendations they give to parents and what source of information most influenced their decisions. A fourth group, pediatricians with particular expertise in nutrition, were surveyed as well. The overall response rate was 76%, with a total usable sample of 252. In all professional groups, 54% had no preference for any one commercially prepared formula. More importance was given to sodium content than to fat composition of formulas. On the choice of whole, low-fat, or non-fat milk for both 1- and 6-year-old children, professional groups differed significantly. Dietitians and the subgroup of pediatricians with nutrition expertise were more likely to recommend milk with higher fat content than other professional groups. Recommendations for both sodium- and fat-modified diets for children depended on CVD risk, and opinions varied between groups. Pediatricians and nurse practitioners were more likely to recommend dietary modifications for children with higher CVD risk. The variation in dietary recommendations within and between professional groups strongly indicates the need for research on the safety and efficacy of dietary restrictions in childhood.

0065

Early weight control concerns affect entire life.

Forester, D. Denver, Colo. : American School Food Service Association, School foodservice journal, Mar 1987, v. 41 (3), p. 97-98. 111. (NAL Call No. : DNAL 389.8 SCH6).

Abstract: The facts that one in four elementary school children are obese and that the incidence of heart disease and diabetes are increased with obesity substantiate the need for weight control early in life. However, also of concern are the psychological problems relating

to obesity in children (i.e. peer abuse and discrimination). Heredity, environmental and behavioral factors influencing obesity are reviewed. Recommendations are provided for promoting physical activity and appropriate food choices to children. Techniques for applying behavior modification to control weight are pointed out.

0066

Eating disorders and diabetes.

Fairburn, C. ; Peveler, R. Alexandria, Va. American Diabetes Association. Diabetes forecast. May 1989. v. 42 (5). p. 33-35, 38. (NAL Call No. DNAL RC660.A1D5)

Abstract: This article discusses the problems associated with diabetic patients who also have eating disorders. The discussion includes detection of the eating disorder, medical complications and treatment.

0067

Eating disorders in youth becoming major concern.

Washington, D.C. Community Nutrition Institute. Nutrition week. Oct 22, 1987. v. 17 (42). p. 6. (NAL Call No. DNAL TX341.C6).

Abstract. A technical commentary discusses the increasing prevalence of childhood obesity and the growing epidemic nature of eating disorders in the US. Specific attention is given to findings of obesity prevalence, the relationship of eating disorders to mental depression, and the observed limited success in the management of eating disorders. Research recommendations to determine the nature of eating disorders and the need for strong education and prevention programs also are discussed (wz)

0068

Eating habits food, physiology and learned behaviour /edited by Robert A. Boakes, David A. Popplewell, Michael J. Burton.

; Boakes, Robert A -Popplewell, David A.-Burton, Michael J. Chichester ; New York : Wiley, c1987. xii, 225 p. ill. ; 24 cm. Includes bibliographies and indexes. (NAL Call No.: DNAL TX357.E23).

Abstract: A reference text addressing various aspects of eating behavior presents 8 authoritative reviews by experts from 5 countries (US, England, France, Australia, Scotland) for clinical psychologists and psychiatrists dealing with patients having adverse eating behaviors. Attention is focused on: the nature, epidemiology, clinical aspects, and treatment of bulimia and anorexia nervosa; behavior modification approaches in treating obese patients; hunger, satiety, and eating behaviors and preferences in early infancy and childhood; the concept and underlying mechanisms of palatability; mental signals affecting meal size; and the cognitive experimental psychology of appetite. Eating behavior descriptions, diagnoses, behavioral therapies, and theoretical models are presented

throughout the text

0069

Echocardiographic and electrocardiographic measures in obese children after an exercise program.

IJDBDP. Hayashi, T. ; Fujino, M.-Shindo, M -Hiroki, T.-Arakawa, K. London : John Libbey & Company. International journal of obesity. Oct 1987. v. 11 (5). p. 465-472. ill., charts. Includes 14 references (NAL Call No. DNAL RC628.A1D2).

Abstract. Exercise training and its influence on anthropometric, hemodynamic, and anatomic adaptive changes in the cardiovascular system were studied in 8 obese children, ages 10 to 11. Children participated in a jogging program that met 5 days/week for 1 or 2 years. Weight decreased after 1 year of exercise training. There were no changes in left ventricular wall thickness. Total voltage in SV1 + RV5 decreased after 3 months of exercise training but returned to pre-training voltage after 1 year of training. Implications are that a program of jogging 5 days/week may produce a weight reduction, a decrease of the resting heart rate and an increase in left ventricular end-diastolic dimension. The effects of exercise on cardiac structure may differ among normal weight children, obese children and adults.

0070

Education helps teenagers alter their food habits.

Carroll, A. Denver, Colo. : American School Food Service Association. School foodservice journal. Dec 1986. v. 40 (10). p. 52-53. ill. (NAL Call No. DNAL 389.8 SCH6).

Abstract. In an effort to help students commit themselves to a healthier lifestyle, a high school in Louisiana has taken an innovative approach to the topic of fitness. Concerned about low female participation in the school lunch program, questionable salad bar selections, excessive snacking and misconceptions about sports nutrition, a meaningful nutrition education unit was developed. The program begins with students completing a 24-hour dietary recall based on the food groups. Following sessions feature dietary guidelines, weight management, salad building, snacking, and sports nutrition. Topics are presented by involving students in various activities; sessions last one week. The program was developed by the Louisiana Cooperative Extension Service and the Nutrition Education and Training Program of the state.(jd-b)

0071

Effect of diet and controlled exercise on weight loss in obese children.

JDPDA. Epstein, Leonard H. ; Wing, Rena R.-Penner, Barbara-Kress, Mary Jeanne. St. Louis : C.V. Mosby. The Journal of pediatrics. Sept 1985. v. 107 (3). p. 358-361. charts. Includes 11 references. (NAL Call No.: DNAL FNC RJ1.A453).

Extract: The effects of adding exercise

to diet for weight control in obese children were evaluated by randomizing obese girls to one of two groups: diet and diet plus exercise. During the first 6 weeks of the treatment, children exercised in a supervised three times a week exercise program, in which they walked or ran 3 miles. Significant decreases from baseline weight and in percent overweight were observed for both groups during the year of treatment. Significant decreases in percent overweight were observed at 0 to 2 months and then at 2 to 6 months for the children who were exercising, whereas percent overweight in children in the diet-alone group decreased only from 0 to 2 months. In addition, a significant improvement in fitness was observed only for children in the diet plus exercise group. (author)

0072

The effect of weight loss on the sensitivity of blood pressure to sodium in obese adolescents.

NEJMAG. Rocchini, A.P., Key, J., Bondie, D., Cnico, R., Moorehead, C., Katch, V., Martin, M. Boston, Mass. Massachusetts Medical Society. New England Journal of Medicine. Aug 31, 1989, v. 321 (9), p. 580-585. charts. Includes 45 references. (NAL Call No. DNAL FNC 448.8 N442).

Abstract. Obese and non-obese adolescents were studied to determine the effect of weight loss and the effect of sodium intake on blood pressure. The subjects were put on a weight loss, low salt diet. The results showed that blood pressure is sensitive to dietary sodium intake in obese adolescents. Details of the study and tables of results are included.

0073

Elementary school principals' perceptions of childhood obesity.

Price, J.H.; Desmond, S.M.; Stelzer, C.M. Kent, Ohio. American School Health Association. Journal of School Health. Nov 1987, v. 57 (9), p. 367-370. charts. Includes 5 references. (NAL Call No.: DNAL LB3401.J6).

Abstract. A random survey of 277 elementary school principals assessed their perceptions of childhood obesity and the school's role in dealing with this. The results indicated that half of the principals believed normal weight important to child health, and, while 35% thought schools were not doing enough to combat childhood obesity, a number felt that the school should/could not be involved. Nonetheless, they favored providing low-calorie lunches, eliminating junk food machines, and the role of school nurses in treating childhood obesity. Survey results are statistically summarized.

0074

Emotions and obesity among Mexican-American children.

JSHEA. Guinn, Bobby. Kent: American School Health Association. The Journal of School Health. Mar 1985, v. 55 (3), p. 113-115. charts. Includes 20

references. (NAL Call No. DNAL FNC LB3401.J6 F&N).

Abstract. A survey assessed possible relationships between high body fat levels and emotional motivations for eating in 17 Mexican-American children, ages 10-14. The results indicated that high body fat correlated with emotional motivation of stimulation in the male children, and that high body fat correlated with handling and craving motivators in the female children. The statistical summaries of the survey data are presented and discussed. (wz)

0075

Energy and macronutrient intake of New Zealand adolescents.

ECFNB. Poh Tan, S.; Wells, J.E.; Beaven, D.W.; Hornblow, A.R. London: Gordon & Breach Science Publishers. Ecology of food and nutrition. Nov 1989, v. 23 (3), p. 225-236. charts. Includes 55 references. (NAL Call No. DNAL TX341.E3)

Abstract. During the months of April and May (autumn season in the Southern hemisphere) 1985, a 3-day 24-hour dietary recall was collected from 255 female and 246 male third form students (12-14 years old) from secondary schools of an urban region in the South Island of New Zealand. Energy and macronutrient intake information was derived from this recall data. The median intake of energy for girls was 7.5 MJ and boys 10.1 MJ. Although these intakes were only 83% and 92% of the daily recommended energy levels for girls and boys respectively, the anthropometric measurements taken did not indicate that they were underweight. Their protein intake was more than adequate; the median intake value was 55g for girls and 74g for boys. Sixty-five percent of protein source was of animal origin and this also largely accounted for the high intake of fat (79g for girls and 104g for boys) and in particular saturated fat (37g for girls and 49g for boys). Intake of complex carbohydrates and dietary fibre (17g for girls and 23g for boys) was low but sucrose intake (50g for girls and 66g for boys) was relatively high. Sucrose was derived mainly from common white cane sugar and manufactured foods. The cholesterol intake was moderately high with about 25% of girls and 40% of boys having an intake greater than 300mg per day. The main dietary sources of cholesterol came from the meat group as well as milk and dairy products, but not eggs.

0076

Energy expenditure and intake in infants born to lean and overweight mothers.

NEJMAG. Roberts, S.B.; Savage, J.; Coward, W.A.; Chew, B.; Lucas, A. Boston, Mass.: Massachusetts Medical Society. New England Journal of Medicine. Feb 25, 1988, v. 318 (8), p. 461-466. 111., charts. Includes 36 references. (NAL Call No.: DNAL FNC 448.8 N442).

Abstract. A postnatal study assessed the contributions of low energy expenditure (EE) and high energy intake to excessive

weight gain in infants born to 6 lean and 12 overweight mothers. No significant difference in weight, length, skinfold-thickness, metabolic rate at 0.1 and 3 months after birth, and metabolizable energy intake at 3 months after birth in infants who became overweight at 1 year of age (50% of those born to overweight mothers) and infants who did not, but total EE 3 months after birth was ca. 20% lower in those who became overweight. The results indicated that reduced EE influenced excessive weight gain in infants from overweight mothers. (wz)

0077

Energy expenditure of young men from obese and non-obese families.

Avons, P. ; James, W.P.T. London, Eng. J. Libbey. Human nutrition clinical nutrition. July 1986. v. 40C (4). p. 259-270. charts. Includes 26 references (NAL Call No. DNAL TX341.H8)

Abstract This study reassesses a previous study of children of obese and non-obese parents. Metabolic rates were observed during sleep, at rest, in response to food and during a variety of exercises for the whole body calorimetric effects. Conclusions were that consistent demonstration of metabolic abnormalities in individuals who are normally considered prone to obesity is difficult. Only some of this adult group confirmed the observation of low energy requirements in children from obese families. (ran)

0078

Energy intake in children at high and low risk of obesity.

Griffiths, M. ; Rivers, J.P.W -Payne, P.R. London J Libbey Human nutrition clinical nutrition Nov 1987. v. 41C (6) p 425-430. charts Includes 12 references (NAL Call No. DNAL TX341 H8).

Abstract A study of energy intakes of 37 young children at low or high risk of subsequently developing obesity (degree of risk judged from parenteral body weight status) found higher intakes for the high-risk group. The results supported those of an earlier study, suggesting that, since basal metabolism and body weight are correlated, studies of child obesity risks be directed at studies of differences in energy expenditure.

0079

Energy need for weight maintenance in human beings: effect of body size and composition.

JADAA. Forbes, G.B. ; Brown, M.R. Chicago, Ill. : The Association. Journal of the American Dietetic Association. A continuing education self-assessment test is appended to the article. Apr 1989. v. 89 (4). p. 499-504. charts, forms. Includes 21 references. (NAL Call No.: DNAL FNC 389.8 AM34).

Abstract: We estimated energy requirement by determining the amount of food needed to maintain body weight in a controlled environment. In a study of 28 adolescents and adults of widely varying

weight and body fat content, we found that the energy required for weight maintenance was directly proportional to body weight ($r=0.92$). The increased energy requirement of the obese is due in part to their larger lean weight and in part to their greater burden of body fat; together these account for 87% of the variance. For these subjects, who were engaged in light physical activity, the ratio of total energy to basal metabolic rate was 1.52 ± 0.16 .

0080

An epidemiologic study of maladaptive eating attitudes in a Canadian school age population.

INDIDJ. Leichner, P. ; Arnett, J.-Rallo, J.S.-Srikameswaran, S.-Vulcano, B. New York John Wiley & Sons. The International Journal of eating disorders. Sept 1986. v. 5 (6). p. 969-982. charts Includes 24 references (NAL Call No. DNAL RA784.A15).

Abstract. Although cases of anorexia nervosa and bulimia are being seen increasingly by health care professionals, little data is available on the prevalence of these disorders in the general population of school age children. Using a validated eating attitude test (EAT), a total of 5150 students, aged 12-20, from public schools and one university in the Province of Manitoba were surveyed. Overall, 5 per cent of males and 22 per cent of females scored 30 or above on the scale, suggesting significant concerns and attitudes regarding eating. These concerns were somewhat higher in urban versus rural settings and seemed to increase between the ages of 12 and 13 and remain high thereafter. Many of the students who score high on the EAT were overweight, suggesting that these attitudes or concerns are not specific to anorexia nervosa and/or bulimia. (author)

0081

Epidemiological studies on cardiovascular risk factors during childhood: total and HDL cholesterol in relation to diet.

Knuiman, J.T. ; West, C.E. New York : Liss, c1985. Detection and treatment of lipid and lipoprotein disorders of childhood : proceedings of the Third International Atherosclerosis Conference, held in Vienna, Austria, April 4-9, 1983 / editors, Kurt Widhalm, H.K. Naito. p. 139-144. Includes 18 references. (NAL Call No.: DNAL RC692.I467 1983).

Abstract: A brief summary is presented on the results of 2 recent international epidemiological studies that were directed toward finding answers to questions posed by the World Health Organization in 1977 concerning relationships between dietary and circulatory levels of cholesterol and risk factors for cardiovascular disease in children. The first study was designed to obtain data on serum total and high-density-lipoprotein (HDL) cholesterol levels in 7-8 year-old boys from 16 countries with different

coronary heart disease (CHD) mortality rates. The second study examined whether a high saturated fat, low complex carbohydrate diet ("westernized diet") is associated with relatively high serum total and HDL cholesterol levels in boys from Finland, the Netherlands, Italy, the Philippines and Ghana. The results show that higher serum total and HDL cholesterol levels are associated with westernized diets, representing higher CHD risks. The results also show that there are large differences in the level of CHD risk among different groups of countries. (wz)

0082

Erythrocyte Na⁺-K⁺ ATPase activity in childhood: regulation by genetic factors independent of body weight.

IJDBDP. Mazelis, A.G.; Larson, S.-Ginsberg-Fellner, F. London: John Libbey & Company International journal of obesity, Dec 1987, v. 11 (6) p. 561-570. 111 charts. Includes 34 references. (NAL Call No. DNAL RC628.A102).

Abstract: The significance of sodium-potassium ATPase activity in the development of childhood obesity was studied. The author specifically wanted to determine if (1) the activity of ATPase is correlated with the development of obesity and (2) if enzyme levels are genetically determined and possibly unrelated to the development of obesity. It was demonstrated that there was no significant relationship between obesity and erythrocyte sodium-potassium ATPase activity in children of varied ages, body weight, and ethnic origin. In addition, the enzyme's activity showed no relationship to the diabetic state of the mother. It was concluded that erythrocyte sodium-potassium ATPase in children is primarily regulated by genetic factors which are generally independent of body weight.

0083

Estimated overweight and obesity in Mexican American school children.

IJDBDP. Malina, R.M.; Zavaleta, A.N.-Little, B.B. London: John Libbey & Company, International journal of obesity, 1986, v. 10 (6), p. 483-491. charts. Includes 24 references. (NAL Call No. DNAL RC628.A102).

Abstract: A 11-year longitudinal study (1972 and 1983) examined overweight prevalence in a cohort of 6-17 year-old Mexican-American children. The results indicate that body mass index (BMI) and triceps skinfold (TSF) measurements may vary in sensitivity as predictors of overweight and obesity. Children classified as obese by BMI had larger estimated midarm muscle circumferences than those classified as obese by TSF alone or by both TSF and BMI. (wz)

0084

The evolution of serum lipoproteins in infancy.

NUREA. Washington, D.C. Nutrition Foundation. Nutrition reviews, Oct 1986, v. 44 (10), p. 324-326. charts. Includes 13 references. (NAL Call No. DNAL FNC

389.8 N953).

Abstract: Recent studies concerning developmental changes in serum lipoprotein subfractions (chylomicrons, and very-low-, low-, and high-density lipoproteins) during infancy are summarized and discussed. The results demonstrate that the concentration of these subfractions and the proportions of their components (protein, triglyceride, free and esterified cholesterol, phospholipid) differ markedly from adult lipoproteins. The possible influence of feeding on these differences is discussed. (wz)

0085

Exercise adaptation responses for gastric inhibitory polypeptide (GIP) and insulin in obese children--possible extra-pancreatic effects.

DIAEA. Kahle, E.B.; D'Donizio, T.M.-Walker, R.B.-Eisenman, P.A.-Reiser, S.-Cataland, S.-Zipf, W.B. Alexandria, Va.: American Diabetes Association. Diabetes, May 1986, v. 35 (5), p. 579-582. 111 charts. Includes 27 references. (NAL Call No. DNAL FNC RC658.A105).

Extract: Thirteen obese children and matched controls were fed a mixed meal, and responses were evaluated at fixed intervals for glucose, insulin and gastric inhibitory polypeptide (GIP). The obese children were evaluated before and within 48 h after completion of a 5-mo exercise training program (ETP). The ETP included three aerobic exercise sessions per week and modest diet restrictions. Caloric expenditure was increased by approximately 300 kcal/exercise session. Weight gain was minimal over the 5 mo. An unexpected increase in GIP response and improved insulin tolerance were recorded for the obese children post-ETP. GIP values were higher (P less than 0.05) at 30 and 60 min and led to a highly significant elevation (P less than 0.01) of the integrated GIP response for post-ETP obese versus both pre-ETP and normal-weight controls. Insulin values were lower (P less than 0.05) at 30 and 60 min and led to a lower integrated insulin response (P less than 0.0585) for post-ETP obese children. However, the obese children continued to secrete more insulin (P less than 0.05) than normal-weight controls. Glucose tolerance, similar for pre-ETP obese subjects and controls, did not change in post-ETP children. Exercise-induced improvement in glucose utilization in these obese children was associated with an increase in GIP secretion. This contrasts with reports that calorie restriction will improve glucose utilization with decreased insulin and GIP secretion. The study demonstrates a previously unreported uncoupling of GIP and insulin secretion and suggests shifts in peripheral tissue sensitivity to insulin-induced glucose uptake. These shifts may, in part, be influenced by GIP. (author)

0086

External responsiveness to food and non-food cues among obese and non-obese children.

IJOBDB. Sobhany, M.S. ; Rogers, C.S
London John Libbey & Company
International journal of obesity 1985
v 9 (2), p 99-106 11 charts.
Includes 53 references (NAL Call No.
DNAL RC62B.A102).

Abstract A study assessed the hypothesis that obese (relative to non-obese) children were less able to delay an immediate small gratification for a larger, delayed reward when exposed to food items (but not non-food items). The ability of preschool children to delay gratification was significantly less than that of school children. Obese preschool and school children had lower gratification delay scores for food items (candy, cupcakes) than comparable non-obese children. Delay scores for non-food items (toys, balloons, comic books) were similar for obese and non-obese children. Hence, the findings of this study support the hypothesis (wz)

0087

Factors related to obesity in preschool children.

JADAA. Patterson, R.E. ; Typpo, J.T.-Typpo, M.H.-Krause, G.F. Chicago, Ill. The Association. Journal of the American Dietetic Association, Oct 1986, v. 86 (10), p. 1376-1381, charts. Includes 70 references (NAL Call No DNAL FNC 389.8 AM34)

Extract The purpose of this research was to examine the following variables for their relationship to the prevalence of preschool obesity: familial aggregation of obesity, infant feeding practices, socioeconomic status, and parents' attitudes toward the use of food for non-nutritive purposes. Parents completed a biographical data form. Height, weight, and skinfold measurements were obtained from 94 preschool children and their biological parents. Both parents answered a Child Feeding Opinion Questionnaire. Anthropometric measurements were evaluated using percentile rankings from NHANES. On the assumption that subjects over the 75th percentile for triceps skinfold were overweight and those above the 90th percentile were obese, 23.4 per cent and 7.5 per cent of the children, 9.6 per cent and 5.3 per cent of the mothers, and 29.8 per cent and 10.6 per cent of the fathers were overweight or obese, respectively. Most parent-child anthropometric correlations were statistically significant. No statistically significant relationships were found between infant feeding practices and childhood obesity. Mothers' educational level varied inversely with the children's weight for height. Mothers and fathers opposed the use of food for reward, punishment, soothing, or affection. The parents' child feeding attitudes had no obvious relationship with the children's anthropometric measurements.(author)

0088

Family-based behavioral weight control in obese young children.

JADAA. Epstein, L.H. ; Valoski, A.-Koeske, R.-Wing, R.R. Chicago, Ill. The Association. Journal of the American Dietetic Association, Apr 1986, v. 86 (4), p. 481-484, charts. Includes 20 references. (NAL Call No DNAL FNC 389.8 AM34).

Extract. The effects of a behavioral weight control program for children aged 1 to 6 were documented. Significant weight change associated with no decrease in child height percentile was observed. Nutrient analysis showed significant decreases in caloric and fat intake. Significant decreases in percent RDA were observed only for iron. Increases in nutrient density were observed for all nutrients except fat.(author)

0089

Fat-free mass in children and young adults predicted from bioelectric impedance and anthropometric variables.

AJCNA. Guo, S. ; Roche, A F.-Houtkooper, L. Baltimore, Md. American Society for Clinical Nutrition. American journal of clinical nutrition, Sept 1989, v. 50 (3), p. 435-443, charts. Includes 37 references. (NAL Call No. DNAL FNC 389.8 J824).

Abstract Fat-free mass (FFM) values calculated from densitometry by using a multicomponent model were significantly (p less than 0.05) larger than those from the Siri two-component model (465 males, 441 females), especially in children and females. The multicomponent model assumes the density of FFM varies by age and sex because of differences in its constituents. With FFM values from the multicomponent model as the dependent variable and impedance and anthropometry as the independent variables, prediction equations were derived by all possible subsets of regression (140 males, 110 females). These equations had RMSEs of 2.2-2.3 kg and CVs of 5.0-5.8%. Cross-validation results were excellent for young adults but less satisfactory for children for whom a different calf skinfold had been measured. The selected equations are applicable to healthy white individuals aged 7-25 y.

0090

Fat in the diets of adolescent girls with emphasis on isomeric fatty acids.

AJCNA. Van den Reek, M.M. ; Craig-Schmidt, M.C.-Weete, J.D.-Clark, A.J. Bethesda, Md. : American Society for Clinical Nutrition. American journal of clinical nutrition, Apr 1986, v. 43 (4), p. 530-537, charts. Includes 36 references. (NAL Call No.: DNAL FNC 389.8 J824).

Extract: To determine the amount of isomeric fatty acids in the diets of a segment of the American population, daily food intake was collected, using the duplicate portion method, from eight healthy white adolescent girls for 7 days. The fifty-six diets were analyzed for fatty acids by gas-liquid

chromatography. The amount of trans isomers of octadecenoic acid (18:1) in the diets of the eight girls ranged from 3.5 to 8.2 per cent of total fatty acids with an average of 5.3 per cent. Other trans fatty acids included trans isomers of 14:1 and 16:1, and cis,trans and trans,cis isomers of 18:2. No measurable amounts of trans trans octadecadienoic acid (18:2tt) were found in the diets of the girls. The total trans fatty acid content of the diets averaged 6.5 per cent of total fatty acids. The daily consumption of total trans fatty acids by the eight girls over a 1 wk period averaged 3.1 g, with 2.6 g of this being 18:1. (author)

0091

Fat nutrition in RDS infants

Van Nuys, Calif. Nutrition & the M.D. Nutrition & the M.D. Oct 1986 v. 12 (10) p. 1 (NAL Call No.: DNAL TX341 N836)

Abstract: A brief review highlights the importance of dietary fat intake for infants with respiratory distress syndrome (RDS) noting that critically ill, unresuscitated RDS infants require intravenous therapy with a fat emulsion to correct low linoleic acid status. It also is noted that such therapy should include carnitine to ensure the absence of a carnitine deficiency. It is argued that this therapy is appropriate for low birth weight infants as well as those with RDS. (wz)

0092

Fatness and obesity among the parents of lean probands.

ECFNE. Garn, S.M., Sullivan, T.V., Hawthorne, V.M. London: Gordon & Breach Science Publishers. Ecology of food and nutrition, 1989, v. 22 (4), p. 277-283. charts. Includes 17 references. (NAL Call No.: DNAL TX341 E3)

Abstract: As shown in 1368 proband-parent pairings involving lean probands and their parents, the fathers and mothers of lean probands tend to be of reduced fatness level themselves (-0.25 Z scores for the sum of two skinfolds) and infrequently obese (10.2% of cases as compared with 16% expectancy). However, these descriptive values are curvilinearly related to the age of the proband, maximizing when the proband is an adolescent. For adolescents, fathers and mothers of lean probands average 0.36 Z scores below the total sample and are least often obese (less than 8%). It may be concluded that adolescents comprise the prototypical sample for family-line studies of fatness and leanness, either because "familial" obesity is best expressed in this age group or because parent-child resemblances in fatness level are a temporal function of the living-together (or cohabitational) effect.

0093

Fatness and obesity of the parents of obese individuals

JFSTA. Garn, S.M., Sullivan, T.V., Hawthorne, V.M. Mysore: Association of Food Scientists and Technologists. Journal of food science and technology, Dec 1989, v. 50 (6), p. 1308-1313. charts. Includes 20 references. (NAL Call No.: DNAL 389.8 J823).

Abstract: As shown in 1419 pairings of obese probands with their parents drawn from a larger series of greater than 9000 proband-parent pairings, the fathers and mothers of obese probands are of increased fatness level (+0.27 Z scores) and more often obese than expected (odds ratio 1.50 overall). However, the tendency towards increased fatness and a greater prevalence of obesity among the parents of obese probands bears a curvilinear relationship to the age of the proband, being least when the probands are young, peaking when the sons and daughters are teen-agers, and declining thereafter. Parents of lean probands in turn tend to be lean themselves (averaging -0.25 Z scores) and least often obese when their progeny are teen-aged. As shown in a two-generational context, familial obesity is best demonstrated in adolescents and their parents, either reflecting years spent in common or a specific etiology for adolescent-onset obesity.

0094

Fatty acid composition of serum cholesteryl esters in 3- to 18-year-old Finnish children and its relation to diet.

AJCNA. Moilanen, T.; Rasanen, L.; Viikari, J.; Akerblom, H.K.; Ahola, M.; Uhari, M.; Pasanen, M.; Nikkari, T. Bethesda, Md.: American Society for Clinical Nutrition. American journal of clinical nutrition, Oct 1985, v. 42 (4), p. 708-713. ill., charts. Includes 24 references. (NAL Call No.: DNAL FNC 389.8 J824).

Extract: The composition of fatty acids in serum cholesteryl esters (CE) was analyzed gas chromatography in 1348 boys and girls aged from 3 to 18 yr. A dietary survey was carried out simultaneously using the 48-h recall method. The dietary P/S ratio had highly significant correlations with CE fatty acids: positive with linoleate and total omega 6 fatty acids and negative with saturated, monounsaturated, and omega 3 polyunsaturated fatty acids. The highest mean percentage of CE-linoleate was found in 15-yr-old girls and lowest in 3-yr-old girls. Age, sex, and the degree of puberty had no independent effect on CE-linoleate after it had been adjusted for the effect of dietary P/S ratio by analysis of covariance. These results indicate that the fatty acid composition of serum CE depends on the quality of dietary fat and that CE-linoleate is a useful reflector of the dietary P/S ratio. The negative correlation between CE omega 3 fatty acids and dietary P/S ratio may be due to displacement of the

omega 3 acids in serum CE by the much higher proportion of dietary linoleate

0095

Fear of obesity among adolescent girls.

PEDIAU Moses, N. ; Baniliv, M.M.-Lifshitz, F. Elk Grove Village, Ill American Academy of Pediatrics Pediatrics Mar 1989 v 83 (3) p 393-398. Includes 43 references (NAL Call No. DNAL FNC RJ1.P42)
Abstract The perceptions concerning weight, dieting practices, and nutrition of 326 adolescent girls attending an upper middle-class parochial high school were studied in relation to their body weight. Underweight or overweight students were those with greater than 10% body weight differential for height. The high school students reported an exaggerated concern with obesity regardless of their body weight or nutrition knowledge. Underweight, normal weight, and overweight girls were dieting to lose weight and reported frequent self-weighing practices. As many as 51% (n = 60) of the underweight adolescents described themselves as extremely fearful of being overweight and 36% (n = 43) were preoccupied with body fat. A distorted perception of ideal body weight was documented, particularly among the underweight students; the greater the underestimation of perceived ideal body weight, the greater the actual deficit in ideal body weight for height of the students (r = .73, P less than .001). Normal weight and overweight girls had better concordance between their actual and perceived ideal body weight for height. The frequency of bingeing and vomiting behaviors was similar among the three weight categories. The data suggest that fear of obesity and inappropriate eating behaviors are pervasive among adolescent girls regardless of body weight or nutrition knowledge

0096

Food intakes of children, the DHSS, and the prevention of heart disease.

NUHEB Crawford, M.A. ; Doyle, W.-Drury, P.J.-Meadows, N. Berkhamstead A B Academic Publishers. Nutrition and health. 1987. v. 5 (1/2). p. 65-77. charts. Includes 21 references. (NAL Call No. DNAL RC620.A1N84).
Abstract A reanalysis was conducted on data developed in a British Department of Health and Social Security (DHSS) survey of the food intakes of school children to assess the effect of such intakes on heart disease prevention. The reanalyzed data are presented, showing the consumption patterns of energy, nutrients, cholesterol, junk food, fat quality, lean meats, unsaturated and saturated fatty acids, and dietary changes. It was concluded that the children's diets were unsatisfactory for preventing cardiovascular disease in later life.(wz)

0097

Food patterns in youth and their effect on later life

Boulton, T.U.C. ; -Nichols, J -Magarey, A. London Libbey, 1987. Food and health issues and directions / edited by Mark L Wahlqvist ... et al. . p 6-11. charts. Includes 33 references (NAL Call No. DNAL RA784.F6)
Abstract Data are summarized and discussed concerning food and energy intakes of Australian boys and girls and the intakes of micronutrients at different meals, relative to their effect on subsequent heart disease risks in later life. The data indicate that children in Australia have lower total fat intakes and a lower polyunsaturated/saturated fatty acid ratio than children in Finland, a country having one of the highest rates of ischemic heart disease. Mean daily nutrient intake patterns also are discussed (wz)

0098

Foods, fads, and fats in under fives

NUHEE Taitz, L.S. Berkhamstead A B Academic Publishers. Nutrition and health 1987. v 5 (3/4). p. 203-209. ill., charts. Includes 26 references. (NAL Call No. DNAL RC620.A1N84).
Abstract The relationship between diet and atherosclerosis of children, 5 years old or younger, is reviewed. Childhood atherosclerosis is discussed as follows: cardiovascular risk factors such as blood lipid levels, high blood pressure and glucose tolerance; and screening and dietary intervention. Findings of the Bogalusa heart study which associates obesity and low density lipoprotein cholesterol to cardiovascular disease, are compared with the Committee on Medical Aspects of Food Policy's report on diet and cardiovascular disease

0099

Frontiers in clinical nutrition /edited by Norman Kretchmer.

. Kretchmer, Norman.; 1923-. Rockville, Md Aspen Systems Corp . 1986. c1985. xii, 258 p. : ill. ; 24 cm. Includes bibliographies and index. (NAL Call No. DNAL RM217.F76).
Abstract: A reference text for clinical nutritionists and health professionals focuses on various clinical nutrition aspects as related to mothers and their children. The 14 papers comprising the text were presented by experts in their respective fields of investigation at a recent conference and are grouped among 3 principal themes: (1) clinical nutrition during pregnancy and lactation (including maternal nutrition, pregnancy-induced hypertension, and diabetic pregnancy); (2) clinical nutrition concerning the fetus and the infant (including prenatal nutrition and premature birth, growth-retarded and high-growth infants, inborn metabolic errors, diarrhea and malnutrition in children, and hypocalcemia and rickets in newborns); and (3) clinical nutrition in children and adolescents (including adolescent obesity, anorexia, and bulimia; and nutritional requirements

for iron) - Illustrations and data are presented throughout the text and literature citations follow each of the text chapters

0100

Genetic-environmental considerations

AJCN. Falkner, F. Bethesda, Md.

American Society for Clinical Nutrition

American journal of clinical nutrition.

Feb 1985, v. 41 (Suppl.2), p. 436-439

111. Includes 5 references (NAL Call

No. DNAL FNC 389.8 J824).

Abstract Interactions of genetic and environmental factors which may affect growth during infancy are discussed, including considerations of prenatal growth and the maternal-fetal nutritional conduit. Particular attention is focused on indices of early maturity (e.g., birth weight) and other measures of growth and maturity (e.g., head circumference, bone and muscle growth, body fat pattern). Maternal genetic factors appear to be more important than paternal genetic factors, contributing to about 25% of the variance in fetal growth. Differences in female and male growth patterns and in demography are discussed. (WZ)

0101

Genetic-environmental considerations.

AJCN. Falkner, Frank Bethesda, Md.

American Society for Clinical Nutrition

American journal of clinical nutrition.

Feb 1985, v. 41 (2, Suppl), p. 436-439

111. Includes 5 references (NAL Call

No. DNAL FNC 389.8 J824).

Abstract Aspects of the interaction of genetic and environmental factors upon growth in infancy and an assessment of their significance in infant development are reviewed and discussed. Specific attention is given to anthropometric indices of early maturity (e.g., body weight, head circumference) and the other measures of growth and maturity (e.g., body fat, skeletal maturity).

(wz)Workshop on Introduction of Food to Infants

0102

Good nutrition for your growing child.

FDACB. Hale, E. Rockville, Md. Food

and Drug Administration, Department of

Health & Human Services, F.D.A.

consumer, Apr 1987, v. 21 (3), p. 20-27.

111., charts. (NAL Call No., DNAL FNC

HD9000.9.U5A1).

Abstract: While the adults of America seem to be shaping up in terms of nutrition/health practices, many experts feel that this lifestyle improvement has not yet hit the general ranks of younger Americans. In efforts to change this trend, research institutions and medical organizations are turning their attention to the special dietary needs of children, with an emphasis on prevention of such diseases as obesity, coronary heart disease, cancer and osteoporosis. Although disagreement exists on how to best accomplish this, there is agreement that children have an increased need for all nutrients and calories, particularly iron, calcium, folic acid, vitamin B12 and copper.

Treatment of childhood obesity, what children should eat as opposed to what many actually do eat, and specific nutrient needs of young people are addressed. Tables of Recommended Dietary Allowances and USDA Daily Food Guide for children are included (jd-b)

0103

Growth and adiposity of term infants fed whey-predominant or casein-predominant formulas or human milk.

JPGND6 Harrison, G.G.; Graver,

E.J.; Vargas, M.; Churella, H.R.; Paule,

C.L. New York: Raven Press, Journal of

pediatric gastroenterology and

nutrition, Sept/Oct 1987, v. 6 (5) p.

739-747, 111., charts. Includes 24

references. (NAL Call No. DNAL

RJ446.J68).

Extract. Growth, estimated composition of weight gained, and stool patterns of term infants who were fed either a whey-predominant formula or casein-predominant formula in a random design and of breast-fed infants were compared. All infants (N = 111) were healthy, singleton products of uncomplicated pregnancies. Birth weights and other anthropometric measures in the first few days of life were not different among the three feeding groups. Formula or breast milk was the infants' principal source of energy from birth to age 16 weeks. Average energy intakes of formula-fed infants and change of intakes with age were similar in both groups at all ages. Feeding groups were not significantly different at any age in weight, length, weight or length gain, head circumference, skinfold measurements, upper arm fat area and muscle area, or estimated total body fat. Stools of infants on the whey-predominant formula differed from both the breast-fed and casein-predominant formula groups. (author)

0104

Habitual daily energy expenditure and activity levels of lean and adult-onset and child-onset obese women.

AJCN. Blair, D.; Buskirk, E.R.

Bethesda, Md.; American Society for

Clinical Nutrition, American journal of

clinical nutrition, Mar 1987, v. 45 (3),

p. 540-550, charts. Includes 51

references. (NAL Call No.: DNAL FNC

389.8 J824).

Extract. The energy expenditure (EE) of eight lean, eight adult-onset obese (A00), and eight child-onset obese (C00) women was determined over three 24-h periods by the factorial method, modified by subject-keypunched and mechanically recorded activity diaries. Mean daily EE was significantly higher in the pooled obese women (2472 +/- 488 kcal) than in lean women (1979 +/- 302 kcal) due to higher energy costs of sedentary and light activity in the obese. EE during moderate-to-strenuous activity was similar between groups because lean women performed these activities more vigorously. A00 and C00 differed significantly in neither mean EE nor habitual activity. Fat-free mass

(FFM) was a better predictor than body weight of both mean daily EE and the energy cost of activity. These data indicate that EE is positively related to obesity. Obese women tend to limit possible EE by reducing the vigorousness of weight-supported activity. (author)

O105

Health screening in schools. Part II. JOPDA Cross, Alan W. St. Louis C.V. Mosby. The Journal of pediatrics Nov 1985 v. 107 (5) p. 653-661. Includes 32 references. (NAL Call No. DNAL FNC RJ1.A453).

Abstract: The second part of a 2-part review of health screening in schools discusses the incidence, detection, treatment, and recommendations of 4 health-related themes (scoliosis, dental problems, growth and nutrition, and hypertension), and describes program decisions that are needed at the community level. It is argued that, once a screening program is adopted, it needs to be efficiently administered and regularly evaluated, and that all school districts, regardless of the current level of health services, should form a school health committee to study the health needs of the students and the available local health sources. (wz)

O106

Help your teen trim down.

Koral, A New York, N Y W/W Twentyfirst Corp. Weight watchers Oct 1986 v. 19 (9), p. 46-49. ill. (NAL Call No. DNAL RC628.W4).

Abstract: Studies show that 4.5 million teenagers are overweight, and most of these are female. The primary factor in adolescent obesity is lack of exercise. Poor eating habits also contribute to overweight in this population group. More teenagers are home alone today than 10 years ago, and snacking has become a major contributor to adolescent diets. Parents play an important role in helping their overweight children change habits and lose the extra weight. Parents must be understanding and reassuring rather than ridiculing and reprimanding. Because of the wide range of physical and emotional changes occurring in the teenager, weight loss can be a challenge to both the adolescent and the parent. (lsp)

O107

Helping obese youth.

CHEJA, LeBow, Michael D. Ottawa : Canadian Home Economics Association. Canadian home economics journal = Revue canadienne d'économie familiale. Spring 1985, v. 35 (2), p. 72-75. Includes 14 references. (NAL Call No.: DNAL 321.B C162).

Extract: Obesity among juveniles is difficult to explain and treat yet easy to describe. This article points out contradictions to be found among studies of prevalence, potential contributors, and possible concomitants. It then addresses the topic of intervention, emphasizing behavior modification methods as adjuncts to such other accepted therapies as dieting and

exercising. Following this analysis, the article examines four obese youths. Where possible, guidance to minimize the occurrence of a pitfall is provided (author)

O108

Hepatic storage of iron and ferritin in different ethnic groups in Singapore.

NUMEB, Wong, C.T. ; Saha, N. Basel, Switzerland S. Karger. Annals of nutrition and metabolism. Sept/Oct 1985, v. 29 (5), p. 267-273. charts. Includes 16 references. (NAL Call No.: DNAL RM214.N8).

Extract: The concentrations of non-haem iron, ferritin and ferritin-iron were measured in the livers of 137 adults and children collected at necropsy. The concentrations of non-haem and ferritin iron were found to be 146.6 plus or minus 95.2 micrograms per gram and 61.6 plus or minus 32.4 micrograms per gram, respectively, in males and 108.0 plus or minus 61.7 micrograms per gram and 60.6 plus or minus 26.4 micrograms per gram, respectively, in females. The values for males in Singapore were lower than those reported in developed Western countries. No correlation was observed between storage iron and age, of ferritin concentration and age. Concentrations of non-haem iron and ferritin were similar for persons dying from accident and coronary heart disease. The non-haem iron concentration in Chinese (187.9 plus or minus 101.0 micrograms per gram) was significantly greater than that in Indians (103.1 plus or minus 65.8 micrograms per gram), while the ferritin concentration in Chinese (6.18 plus or minus 2.37 mg/g) was significantly greater than either Malays (3.81 plus or minus 1.8 mg/g) or Indians (3.52 plus or minus 1.6 mg/g). A significant positive correlation was observed between the non-haem iron and ferritin and also ferritin-iron in Chinese males (r values of 0.678 and 0.598, respectively) and Indian males (r values of 0.576 and 0.612, respectively). However, the correlation between these indices was not significant in the case of Malay males. In premenopausal women the non-haem iron correlated well with ferritin (r=0.737) and ferritin iron (r=0.826) while the correlation was lacking in postmenopausal women. (author)

O109

High-density lipoprotein-cholesterol subfractions in adolescent twins.

PEDIAU, Bodurtha, J.N. ; Schieken, R.-Segrest, J.-Nance, W.E. Elk Grove Village, Ill. : American Academy of Pediatrics. Pediatrics, Feb 1987, v. 79 (2), p. 181-189. ill., charts. Includes 59 references. (NAL Call No.: DNAL FNC RJ1.P42).

Extract: Data on the levels of high-density lipoprotein-cholesterol (HDL-C) and subfractions in 102 adolescent twin pairs and their parents are presented. Children with a family history of premature cardiovascular death had lower levels of HDL2-C than did those without such a history. White

girls reporting a high level of physical activity had higher levels of HDL-C and HDL2-C than did their more sedentary peers. In general, children of mothers who smoked had lower HDL2-C than did children of nonsmoking mothers. These findings suggest that low levels of HDL2-C in children may identify families in which there is an increased risk of coronary heart disease and that parental smoking may contribute to changes in this risk factor in the children of smokers as well as in the smokers themselves. (author)

O110

HLA system, body fat and fat distribution in children and adults
IJOBDP. Bouchard, C. ; Perusse, L. ; Rivest, J. ; Poy, R. ; Morissette, J. ; Allard, C. ; Theriault, G. ; Leblanc, C. ; Tremblay, A. London : John Libbey & Company. International journal of obesity 1985, v. 9 (6), p. 411-422. charts. Includes 10 references. (NAL Call No. DNAL RC628.A102).
Abstract A genetic and anthropometric study tested the presence of associations between the alleles or the genotypes at the A, B, and C loci of the HLA system and percent body fat, subcutaneous fat, and fat distribution in 1578 volunteers 705 adults (mean age, 43.2) and 873 children and adolescents (mean age, 14.6) of both sexes who lacked metabolic disorders and who were not grossly obese. The results did not support earlier reports of an association between high fat content and antigens B18, Bw35, or Cw4, no consistent pattern of association was present between alleles or genotypes of the HLA system and percent body fat, subcutaneous fat, or fat distribution. (wz)

O111

Implementation of a pilot school-site cholesterol reduction intervention.
Resnicow, K. ; Orlandi, M.A. ; Vaccaro, D. ; Wynder, E. Kent, Ohio : American School Health Association. Journal of school health. Feb 1989, v. 59 (2), p. 74-78. charts. Includes 31 references. (NAL Call No.: DNAL LB3401.J6).
Abstract School children, ages 5-11, were given cholesterol screening tests as part of a school health program. Students identified as being at risk were eligible for a cholesterol reduction intervention workshop.

O112

Implementation of "Heart Smart:" a cardiovascular school health promotion program.
Downey, A.M. ; Frank, G.C. ; Webber, L.S. ; Harsha, D.W. ; Virgilio, S.J. ; Franklin, F.A. ; Berenson, G.S. Kent, Ohio : American School Health Association. Journal of school health. Mar 1987, v. 57 (3), p. 98-104. charts. Includes 36 references. (NAL Call No.: DNAL LB3401.J6).
Abstract: Based on data developed during a 12-year study of 8000 children to identify cardiovascular (CV) risk factors (Bogalusa, Louisiana Heart

Study). a comprehensive, research-based CV school health promotion program, "Heart Smart," was developed. This report describes the development, implementation, and evaluation of this program vs. a "high-risk" approach in 4 elementary schools. The aims, objectives, and goals of this program are specified, together with a description of its theoretical framework, design, evaluation methods, and intervention strategies.

O113

Implications and treatment of adolescent obesity.

Dietz, W.H. Jr. St Louis, Mo. C. V. Mosby Company. Clinical nutrition May/June 1985, v. 4 (3), p. 103-108. charts. Includes 43 references. (NAL Call No. DNAL RM216.M342).
Abstract An overview of adolescent obesity addresses its incidence, natural history, its effect on adult morbidity, and its causes, diagnosis, treatment, and prevention. While adolescent obesity is the most prevalent morbid nutritional disease in the US, therapeutic effects have produced a low rate of short-term remission. It is argued that epidemiologic and clinical data indicate that family interactions appear to be the best vehicle for prevention and that television viewing may be the most logical behavior to select for adolescent behavior modification. The results of various weight reduction regimens for adolescents also are discussed. (wz)

O114

An inborn error of cholesterol biosynthesis.

NUREA. Washington, D.C. Nutrition Foundation. Nutrition reviews. Oct 1986, v. 44 (10), p. 334-336. Includes 6 references. (NAL Call No. DNAL FNC 389.8 N953)
Abstract. A case study is presented of a 2-year-old boy with inherited mevalonic aciduria (mevalonic kinase deficiency), complicated with an interruption of isoprenoid compound biosynthesis, including hypocholesterolemia. After diagnosis, treatment, and hospital discharge on palliative therapy, the subject was rehospitalized with diarrhea, dehydration, hyponatremia, and hypokalemia several weeks later, and died from infection and his metabolic disorder. It is suggested that such patients be fed the unavailable, essential isoprene biosynthesis products. (wz)

O115

Increased basal glucose production and utilization in children with recent obesity versus adults with long-term obesity.

DIAEA. Bougueres, P.F. ; Artavia-Loria, E. ; Henry, S. ; Basdevant, A. ; Castano, L. Alexandria, Va. : American Diabetes Association. Diabetes. Apr 1989, v. 38 (4), p. 477-483. charts. Includes 38 references. (NAL Call No.: DNAL FNC RC658.A1D5).
Abstract: To characterize the

abnormalities of glucose homeostasis and insulin action early in the course of human obesity, we studied in vivo glucose kinetics in seven children who were recently massively overweight. At time of study they were gaining weight at a rate of 13.5 +/- 1.4 kg/yr. They were compared with six age-matched control subjects. Six adults with long-term obesity and five normal adults were studied in parallel. The obese children and adults were normoglycemic and hyperinsulinemic. We found that glucose production and utilization were remarkably higher in obese children (295 +/- 18 mg/min; 7.6 mg.kg⁻¹ lean body mass.min⁻¹) than in control children (129 +/- 13 mg/min; 4.4 mg.kg⁻¹ lean body mass.min⁻¹, P less than .01) and obese adults (151 +/- 8 mg/min; 3.1 +/- 0.3 mg.kg⁻¹ lean body mass.min⁻¹, P less than .01). Obese adults had normal rates of glucose production and utilization. Insulin- and non-insulin-mediated glucose uptake, estimated with somatostatin-induced suppression of endogenous insulin secretion, contributed almost equally to the excess glucose utilization observed in the obese children. When studied with the euglycemic-hyperinsulinemic clamp, obese children could not increase glucose disposal to the same extent as normal children and were not able to adequately suppress their endogenous glucose production. Recently obese children are therefore characterized by an increased basal glucose turnover rate and an already established insulin resistance of the liver and probably the skeletal muscles

O116

Increasing pediatric obesity in the United States.

AJDC. Gortmaker, S.L.; Dietz, W.H. Jr.; Sobol, A.M.; Wehler, C.A. Chicago American Medical Association. American Journal of Diseases of Children. May 1987. v. 141 (5). p. 535-540, 111, charts. Includes 45 references. (NAL Call No. DNAL FNC 448.8 AM38).
Abstract. Triceps skin-fold measurements of children obtained in 4 US population surveys were used to estimate trends in pediatric obesity between 1963-80. The results indicated a marked increase in US childhood obesity prevalence, suggesting a 50% increase in obesity prevalence in children of ages 6-11 and a doubling in the prevalence of "superobesity" (at or above the 95th percentile) between 1976-80. The 1976-80 increase in both of these obesity categories was greater than their 1966-70 increase, suggesting an acceleration in obesity prevalence in the US. This trend is attributed to environmental factors. (wz)

O117

Infant growth and obesity in Samoa.

NUREA. Washington, D.C.: Nutrition Foundation. Nutrition reviews. Aug 1986. v. 44 (8). p. 265-267. Includes 11 references. (NAL Call No.: DNAL FNC 389.8 N953).
Abstract: A discussion of recent reports

on infant growth and obesity incidence in American Samoa concludes that the birth weight and growth of Samoan infants are greater than for infants of other countries, regardless of feeding regimen or rural-urban differences. This relatively rapid prenatal and postnatal growth in weight is consistent with the reported incidence of overweight in Samoan childhood and adult life. Genetic influences in these observations are postulated. (wz)

O118

Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales.

Barker, D.J.P., Osmond, C. Boston. Mass. Little, Brown and Company. The Lancet. May 10, 1986. v. 1 (8489). p. 1077-1081, 111, charts. Includes 32 references. (NAL Call No. DNAL 448.8 L22)

Abstract. A study explored the possible association between poor living standards and ischemic heart disease (IHD) by a detailed geographical comparison in England and Wales of infant mortality between 1921-25 and adult mortality from IHD and other leading causes between 1968-78. When the division of the country into 212 local authority areas was considered, a strong geographical association was found between IHD mortality in 1968-78 and infant mortality in 1921-25. Only 3 of 24 other leading causes of mortality in 1968-78 (bronchitis, rheumatic heart disease, stomach cancer) were related as well to infant mortality in 1921-25. It is argued that the results indicate that poor nutrition in early life increases subsequent health risks in adult life (wz)

O119

Infant nutrition and adult health.

Dugdale, A.E. Boston, Mass.: Little, Brown and Company. The Lancet. Apr 23, 1988. v. 1 (8591). p. 948, charts. Includes 11 references. (NAL Call No.: DNAL 448.8 L22).

Abstract. A letter to the editor cites an editorial published in a previous issue of the same journal that suggested that risk factors during infancy result in a high risk for ischemic heart disease (IHD) in later life without providing firm evidence to support this hypothesis in humans. This letter provides anthropometric data among children in Australian Aboriginal communities that indicate that a lower birth weight or weight gain was associated with a higher subscapular/triceps skinfold ratio, indicating a tendency towards central obesity. It is concluded that, while direct evidence is lacking that these childhood fat distribution patterns continue into adult life, the high level of central obesity among Aboriginal adults suggest that they do. It further is noted that central obesity is only an indicator and not, of itself, one of the underlying processes for increased IHD risk in adults, although the data indicate that early malnutrition can

Cause a high-risk distribution of subcutaneous fat. (wz)

O120

Insulin and blood pressure during weight loss in obese adolescents.

Rocchini, A.P.; Katch, V.-Schork, A.-Kelch, R.P. Dallas: American Heart Association Hypertension Sept 1987 v. 10 (3), p. 267-273, 111, charts. Includes 26 references. (NAL Call No. DNAL RC685.H8H9).

Abstract: A 20-week weight loss study on 50 obese adolescents assessed whether a relationship exists between fasting serum insulin and blood pressure in obese adolescents, and what effect weight loss might have on such a relationship. Weight loss was correlated with significant declines in insulin and blood pressure, with the decrease in blood pressure significantly correlating with changes in insulin and body weight. However, when the effect that weight loss had on blood pressure was corrected for, blood pressure and insulin remained correlated only for those subjects involved in both diet and exercise, suggesting that exercise was critical in determining the role of insulin in blood pressure regulation during weight loss (mjs)

O121

Interrelationships of glucose and protein metabolism in obese adolescents during short-term hypocaloric dietary therapy.

AJCN. Dietz, W.H. Jr.; Wolfe, R.R. Bethesda, Md.: American Society for Clinical Nutrition. American Journal of Clinical Nutrition. Sept 1985, v. 42 (3), p. 380-390, 111, charts. Includes references. (NAL Call No. DNAL FNC 389.8 J824).

Extract: We studied the interrelationship of nitrogen balance (N-bal) and rates of glucose appearance (Ra), determined isotopically using U-13C-glucose, in 14 obese adolescents consuming either (.5 g protein and 1.0 glucose)/kg ideal body weight/day or an isonitrogenous diet made isocaloric with fat. Nitrogen balance was significantly more positive with added glucose. Changes in plasma insulin, free fatty acids, or beta-hydroxybutyrate did not reliably predict N-bal. The Ra of glucose decreased significantly on both diets, but was significantly lower after the addition of fat. A significant correlation of N-bal with Ra was observed only in the absence of dietary glucose. Insulin levels correlated with N-bal only in the presence of dietary glucose. Nitrogen balance in the absence of dietary carbohydrate may be a consequence of net peripheral protein catabolism that is not directly mediated by the need for gluconeogenic precursors. (author)

O122

Is there a transient, obesity-related hypertension of adolescence?

CPEDA. Stickler, G.B. Hagerstown, Md.; J.B. Lippincott Company. Clinical Pediatrics. Nov 1986, v. 25 (11), p.

573-574. Includes 9 references. (NAL Call No. DNAL FNC RJ1 C55).

Abstract: A technical commentary raises the question of appropriateness for diagnosing overweight children with diastolic blood pressure above the 95th percentile as having hypertension, noting literature reports recommending that such children be considered as having an obesity-related elevated blood pressure. It is argued that, while it has been recommended that caution be exercised in labeling children as hypertensive because of psychosocial and economic implications and that the term "high normal blood pressure" be used, it is not clear whether this approach will avoid the development of symptoms that appear in association with being labeled "hypertensive." It is concluded that the difficulty in the appropriate choice of terminology and of treatment requires further definition and evaluation. (wz)

O123

Is there an excess of saturated fat in infant formula?

JAMA. Chicago: American Medical Association JAMA Journal of the American Medical Association. Dec 6, 1985, v. 254 (21), p. 3037-3038. Includes 3 references. (NAL Call No. DNAL FNC 448.9 AM37).

Abstract: A "letter to the editor" from 5 medical doctors at the UCLA School of Medicine raises the concern that recent increases in the saturated fat content of commercial infant formulas may present an increased cardiovascular (CV) disease risk to infants in latter life because high saturated fat intakes in later years have been associated with increased CV risk and because the American Heart Association has cautioned that CV disease has its beginnings in childhood. The authors also cite expert opinion that saturated fat intake should not be reduced for infants to less than 10% of total calories, as recommended for older children and adults, since infants usually have very low cholesterol levels. It also is noted that infant formula producers claim that the increased saturated fat levels (from 20% up to 55-80%) in the "improved" formulas more closely represent the saturated fat content of breast milk. (wz)

O124

Issues in childhood nutrition.

Beaudette, T. (ed.). Littleton, Colo.: Seminars in Nutrition. Seminars in Nutrition. Mar/Apr 1987, v. 6 (4), p. 1-11, 1-20, charts. Includes 64 references. (NAL Call No.: DNAL TX341.N828).

Abstract: This review provides coverage of the advances in pediatric nutrition for the dietetic practitioner. Issues on childhood nutrition: recommended dietary allowances; obesity in childhood and adolescence; etiology of obesity; obesity and therapeutic interventions; implications for dietitians; childhood origin of atherosclerosis; cardiovascular risk factors (blood pressure, hypercholesterolemia);

carbohydrates; osteoporosis prevention, and implications for dietetic practice are discussed. A pretest, posttest with their answers, and references are included. (rah)

O125

Kids & diet.

Liebman, B. Washington, D.C. Center for Science in the Public Interest. Nutrition action health letter. Apr 1986, v. 13 (4) p. 1, 4-7 charts. Includes 23 references. (NAL Call No. DNAL TX341.N98)

Abstract Discussing the controversies, surrounding children's diets and heart disease, this article presents differing views of such organizations as the American Heart Association and American Academy of Pediatrics. The former recommends a lower fat diet for the young child, while the latter feels there is no clear evidence that fatty streaks found in the arteries of this age group progress to atherosclerosis. A point of debate appears to be at what age cholesterol levels of 200 or more becoming damaging to arteries. Other discussions center around diet and anemia and breast cancer. It is concluded that all will benefit from developing good eating habits at a young age, and that the schools can be instrumental in helping to achieve this. (jd-b)

O126

Lipid and lipoprotein levels of Newfoundland school children.

NUMEB. Fodor, J.G. ; Balram, B.C.-Hovden, S. Basel, Switzerland S Karger. Annals of nutrition and metabolism. July/Aug 1987 v. 31 (4). p. 231-236 charts. Includes 9 references (NAL Call No. DNAL RM214.N8).
Extract Serum concentrations of total cholesterol, HDL, LDL, and VLDL cholesterol were measured in 1,033 boys and girls age 8-10 years and 14-16 years who were living in two geographically distinct areas on the East and West coast of Newfoundland. The respondents enrolled to this study were school children attending four schools selected at random from all schools in the area. Ninety percent of the invited students participated in the study. The Newfoundland children had higher total cholesterol levels than those reported for children matched for age, sex and race living in the United States. However, their LDL cholesterol levels were found to be similar to those observed in US samples. It was concluded, therefore, that the higher total cholesterol levels of the Newfoundland children were due to their higher HDL cholesterol levels. (author)

O127

Lipids, lipoproteins and alpha-tocopherol: Relationship and changes during adolescence.

NUMEB. Widhalm, K. ; Holzl, Monika-Brubacher, G. Basel, Switzerland ; S. Karger. Annals of nutrition and metabolism. Jan 1985, v. 29 (1). p. 12-18. ill., charts. Includes 26

references. (NAL Call No. DNAL RM214.N8)

Extract From May 1976 until June 1982 a longitudinal study in 54 apparently healthy Austrian schoolchildren with a mean age of 11.2 years at their first visit was performed. The aim of this study was to determine if there are any age-related changes in serum lipids, lipoproteins and alpha-tocopherol concentrations during adolescence and whether a permanent relationship between lipoproteins and alpha-tocopherol can be observed. Total cholesterol showed a significant decrease from age 11 to 14 years in boys as well as in girls; thereafter, a slight increase could be shown. Similar changes could be observed for LDL cholesterol. No significant sex differences were found either in total or in LDL cholesterol, whereas in HDL cholesterol concentrations, a decrease in boys between 12 and 14 years and an increase in girls from 13 years onwards led to significantly lower values in boys than in girls from the age of 16 years onwards. No consistent changes could be shown for alpha-tocopherol blood levels. Nevertheless, a close relationship between total cholesterol and alpha-tocopherol could be observed during all our investigations and, to a lesser degree, between LDL cholesterol and alpha-tocopherol. Significant correlations between alpha-tocopherol and HDL cholesterol and between alpha-tocopherol and triglycerides occurred only occasionally. (author)

O128

Lipoproteins in the progeny of young men with coronary artery disease: children with increased risk.

PEDIAU Lee, J. ; Lauer, R.M. -Clarke, W.R. Elk Grove Village, Ill American Academy of Pediatrics. Pediatrics. Aug 1986, v. 78 (2) p. 330-337. ill., charts. Includes 45 references. (NAL Call No.: DNAL FNC RJ1.P42).
Extract: The authors studied 173 progeny from 63 families in which the father had angiographically diagnosed coronary artery disease by 50 years of age. To assess the nature of the coronary risk factors in these families, we measured their height and weight to calculate Quetelet index (wt/ht²), BP, fasting plasma cholesterol, triglyceride, low-density lipoprotein cholesterol, and high-density lipoprotein cholesterol were measured in the affected fathers, their wives, and their progeny. These values were compared with age- and sex-specific values from the Lipid Research Clinic data. Sixty-five percent of the affected fathers and 51% of the progeny had elevated triglyceride, elevated low-density lipoprotein cholesterol, diminished high-density lipoprotein cholesterol, or combinations thereof. The distribution of the lipids and lipoproteins in the children bore a close resemblance to those observed in the affected fathers. A significant number of the mothers had diminished high-density lipoprotein cholesterol, which was attributed to their obesity. Screening the progeny of young coronary

artery disease patients is therefore highly productive in identifying young people at excessive risk for future coronary artery disease. Early identification of this young high-risk population offers an opportunity for early initiation of preventive measures. (author)

O129

A long-term aerobic exercise program decreases the obesity index and increases the high density lipoprotein cholesterol concentration in obese children.

IJOBOP. Sasaki, J. ; Shindo, M.-Tanaka, H -Ando, M.-Arakawa, K. London : John Libbey & Company International journal of obesity. 1987 v. 11 (4). p. 339-345. 111. charts. Includes 28 references. (NAL Call No. DNAL RC628.A102).

Abstract A 2-year study of the effect of aerobic exercise on body weight and serum lipids in 41 obese Japanese children (20 girls, 21 boys, initial age, 11) revealed a significant decline in obesity index and a significant increase in high-density-lipoprotein cholesterol in both sexes. Total serum cholesterol, however, was unaltered. The overall health benefits of the exercise program are discussed (wz)

O130

Magnesium and other nutrient deficiencies as possible causes of hypertension and low birthweight

NUHEB Wynn, A ; Wynn, M. Berkhamstead A B Academic Publishers. Nutrition and health. 1988 v. 6 (2) p 69-88 charts Includes 92 references (NAL Call No DNAL RC620 A1N84).

Abstract 7.2 percent of babies born in England and Wales in 1986 had birthweights below 2,500 g. Low birthweight and hypertension are associated. European trials have reported that oral supplementation with physiological amounts of magnesium during pregnancy reduces pregnancy hypertension and also miscarriage, preterm birth and fetal growth retardation. Magnesium deficiency causes hypertension and low birthweight in animals. In humans deficiency of thiamin and other B vitamins has also been reported to cause pregnancy hypertension and low birthweight. Magnesium and B vitamins are essential for the same biochemical reactions in energy metabolism. There is evidence that magnesium consumption of substantial numbers of women in Europe and North America is too low to support a healthy pregnancy. Magnesium and thiamin are lost in processing many foods. British trials of magnesium supplementation are advocated. It is suggested that more attention should be given to magnesium in nutritional advice.

O131

Maternal variables related to potentially high-sodium infant-feeding practices.

JADAA. Schaefer, Lynda J. ; Kumanyika, Shiriki K. Chicago, Ill. The Association. Journal of the American

Dietetic Association. Apr 1985 v 85 (4). p 433-438 charts Includes 49 references. (NAL Call No. DNAL FNC 389.8 AM34)

Extract Infant-feeding practices potentially associated with high-sodium intake were assessed for 87 white and 68 black mothers and were analyzed for associations with epidemiological correlates of hypertension. Race and education affected breast-feeding incidence. Education affected breast-feeding duration, introduction of solid foods, salting of infant food, and use of salty snack foods. Salting of infant food was also associated with maternal salt use and with positive family hypertension history; black mothers more often fed salty snacks (Author)

O132

Metabolic evaluation of obese and nonobese siblings.

JOPDA. Elliot, D.L. ; Goldberg, L.-Kuehl, K.S.-Hanna, C St. Louis, Mo. C.V Mosby. The Journal of pediatrics June 1989. v. 114 (6) p 957-962 charts. Includes 43 references (NAL Call No. DNAL FNC RJ1.A453)

Abstract Objective To test the hypotheses that obese adolescents have a lower resting metabolic rate and less aerobic endurance than their nonobese siblings. Design Case-referent study of obese and nonobese siblings from the same kindred. Setting Tertiary referral center Participants Telephone screening of community volunteers resulted in a consecutive sample of 16 kindreds. Obese and nonobese siblings were similar in age, height, and pubertal status. Significantly more female subjects were in the obese group (p less than 0.01). Measurements and main results Body composition studies revealed that the obese siblings had higher body fat (p less than 0.001) but that fat-free mass was similar to that of the lean siblings. Resting metabolic rates determined by indirect calorimetry for the obese and nonobese pairs did not differ. Although the obese siblings appeared less fit when maximal oxygen consumption was measured in relation to total weight, maximal oxygen consumption did not differ when values were standardized for fat-free mass. Conclusions The obese adolescents did not have a reduced resting metabolic rate. As in adults, the relationship between resting metabolic rate and fat-free mass was similar for obese and nonobese children and adolescents. Any decreased sport participation by the obese siblings was not due to inherent reductions in aerobic capacity.

O133

Modification of risk factors for coronary heart disease.

NEUMAG. Walter, H.J. ; Hofman, A.-Vaughan, R.D.-Wynder, E.L Boston, Mass. : Massachusetts Medical Society. New England journal of medicine. Apr 28, 1988. v 318 (17). p. 1093-1100. charts. Includes 46 references. (NAL Call No. DNAL FNC 448.B N442).

Abstract A 5-year study of 3388 school children in 37 New York City schools assessed the effectiveness of an educational intervention program aimed at reducing coronary heart disease risk factors. Significant decreases in plasma total cholesterol and favorable dietary and health knowledge trends were obtained. However, not all targeted risk factors were altered.(wz)

O134

Modifying the eating behavior of young children.

Perry, C.L. ; Mullis, R.M.-Maile, M.C
Kent, Ohio American School Health Association Journal of school health. Dec 1985, v 55 (10) p. 399-402 ill., charts Includes 17 references (NAL Call No. DNAL LB3401.J6)

Abstract: A pilot study carried out in eight 3rd-grade and 4th-grade classrooms assessed the effectiveness of a nutrition education curriculum stressing the importance of a low-salt, low-fat, increased-complex-carbohydrate diet for promoting cardiovascular health. The results indicated that the students had altered their diets to produce a marked decrease in their intake of fat and salt and a concomitant increase in their intake of complex carbohydrates, relative to a control group involving 8 additional 3rd-grade and 4th-grade classrooms. The characteristics of this curriculum, based on social learning theory, are described.(wz)

O135

A multivariate model for assessing eating patterns and their relationship to cardiovascular risk factors: the Bogalusa Heart Study.

AJCN A Nicklas, T A. Webber, L S -Thompson, B.-Berenson, G.S
Baltimore, Md American Society for Clinical Nutrition, American Journal of Clinical Nutrition June 1989, v. 49 (6), p. 1320-1327. Includes 33 references. (NAL Call No.: DNAL FNC 389.8 J824).

Abstract: Eating patterns were studied in 1275 adolescents and young adults (aged 12-24 y). Factor analysis of 64 foods consumed weekly revealed 17 eating-pattern factors, accounting for 57% of the item variance. Factor I (12 food items from either the seafood or meat group) accounted for 8% of the variance, factor II (snacks), 5%, and factors III (fats and pasta) and IV (beef and chicken), 4% each. Remaining factors accounted for from 1.9% to 3.9%. The factors were effective in discriminating eating patterns across race and gender. Significant age effects were also noted for 10 of the 17 factors. Eating patterns for persons in the upper or lower quartiles differed consistently for specific cardiovascular (CV) risk factors. Use of this statistical model to identify differences in eating patterns by race, gender, and CV risk factors during maturation can assist health professionals in targeting food sources for changing eating behavior.

O136

The natural history of serum lipids and lipoproteins during childhood.

Strobl, W. ; Widhalm, K. New York Liss, c1985. Detection and treatment of lipid and lipoprotein disorders of childhood proceedings of the Third International Atherosclerosis Conference, held in Vienna, Austria, April 4-9, 1983 / editors, Kurt Widhalm, H.K. Naito; p. 101-121, ill., charts. Includes 76 references. (NAL Call No. DNAL RC692.I467 1983).

Abstract: A clinical review provides a brief description of and data on changes in serum cholesterol (C), triglycerides (TG), and lipoprotein (LP) and apo-LP levels occurring with age during the fetal period, infancy, and childhood in healthy children, with special emphasis on the neonatal and adolescent periods. These data indicate that the most striking age-dependent changes in serum lipid and LP levels include a dramatic rise in C, TG, low-density-LP-cholesterol (LDLC), and apo-LP levels during the first weeks of life, and a notable decline in total C, TG, LDLC, and high-density-LP-cholesterol (HDL) during adolescence. The data also reveal a considerable degree of tracking (probably from infancy) of serum C, LDLC, HDLC, and apo-LP levels with age during childhood.(wz)

O137

The news about overweight teens.

Adler, V. New York, N.Y. W/W Twentyfirst Corporation, Weight watchers, Mar 1989, v. 22 (2) p. 14, 16 (NAL Call No. DNAL RC628 W4).

Abstract: According to studies, obesity in adolescents has increased substantially over the last 20 years. Many of these teens will become obese adults, predisposing them to a number of health problems. Inheritance, exercise, psychological influences, and dietary habits are discussed as influencing factors in weight gain. Suggestions for proper diet and weight control are included.

O138

No correlation between adiposity and food intake: why are working class children fatter?

AJCN A Rolland-Cachera, M.F. ; Bellisle, F. Bethesda, Md. : American Society for Clinical Nutrition, American Journal of Clinical Nutrition, Dec 1986, v. 44 (6), p. 779-787, ill., charts. Includes 41 references. (NAL Call No.: DNAL FNC 389.8 J824).

Extract: Many studies have failed to show a correlation between individual energy intake and obesity. However, the prevalence of overweight is higher in populations with higher caloric intake. In this study on a population of French children, no correlation was found between energy intake and individual corpulence (wt/ht² index or skinfold thickness), but a higher proportion of overweight children was found in lower social classes where energy intake is traditionally higher. A hypothesis is

Proposed to account for this apparent contradiction the typical lifestyle or diet in a given population challenges individual adaptive capacities. the more caloric the socially accepted diet, the higher the proportion of individuals who are challenged beyond their adaptive threshold. Socially determined factors such as a high calorie diet act in a permissive way in the development of obesity although the caloric intake of obese individuals may be no different from that of nonobese peers (author)

O139

Nutrition and obesity.

Dietz, W.H. Jr. Boston Butterworths. c1987. Pediatric nutrition theory and practice / edited by Richard J. Grand, James L. Sutphen, William H. Dietz. Literature review, p. 525-538, ill., charts. Includes 93 references (NAL Call No. DNAL RJ206.P42).
Abstract: A brief report reviews some of the nutritional aspects of obesity in children and adolescents. Topics include the definition, prevalence, and pathophysiology of child obesity, energy balance; environmental factors, clinical manifestations; evaluation and assessment; and weight reduction and dietary and behavioral modification treatments. Patient compliance and the role of the community in combating child obesity also are discussed (wz)

O140

Nutrition and the heart.

Heymsfield, S.B. Andrews, J.S. Hood, R.-Williams, P.J.-Bagatell, C.J. Boston Butterworths, c1987. Pediatric nutrition theory and practice / edited by Richard J. Grand, James L. Sutphen, William H. Dietz. Literature review p. 597-613, ill., charts. Includes 102 references (NAL Call No. DNAL RJ206.P42).

Abstract: A technical report initially describes the cardiovascular changes that accompany suboptimal nutritional status, including an under- or over-abundance of essential nutrients and pediatric atherosclerosis. This is followed by an examination of somatic and nutritional changes that occur in children whose initial problem is cardiac in nature. Topics include: the heart in suboptimal nutritional states; vitamin, mineral, and trace element imbalances; the characteristics and pathophysiology of atherosclerosis; and the nutritional status of and energy metabolism in children with heart disease. Factors influencing dietary therapy are discussed. (wz)

O141

Nutrition curriculum for families with high blood pressure.

USHEA, Farris, Rosanne P. ; Frank, Gail C.-Webber, Larry S.-Berenson, Gerald S. Kent : American School Health Association. The Journal of school health. Mar 1985. v. 55 (3). p. 110-113. charts. Includes 18 references. (NAL Call No.: DNAL FNC LB3401.J6 F&N).
Abstract: A nutrition education curriculum was designed for 48 students

(ages 8-18) having high blood pressure. The curriculum promoted reduced Na and energy intakes, increased K intake, nutrition education games, taste-tests promoting acceptable snacks, and self-monitoring of intake behavior. While no association was found between curricular compliance and medication use and blood pressure change, children with the highest post-test scores had low Na/creatinine ratios. (wz)

O142

Nutrition for children /Dorothy E.M. Francis ; with a foreword by Barbara E. Clayton.

Francis, Dorothy E. M. 1938-. Oxford Oxfordshire : Boston : Blackwell Scientific. : St Louis, Mo. : Blackwell Mosby Book distributor. 1986. xi, 164 p. ill. : 21 cm. Includes bibliographies and index. (NAL Call No. DNAL RJ206.F73)

Abstract: A reference text for parents and professionals details the nutritional needs of infants and children for promoting optimal growth and provides practical guidelines for infant/child nutritional feeding based on current scientific knowledge. The text addresses 3 specific areas of child nutrition, the role of nutrition in health and disease; the etiology, treatment, and management of overweight and obesity in children; and the role of vitamins and minerals in promoting child nutritional health. Information also is included on catabolic states, enteral feeding in children, infant feeding and weaning, and special therapeutic dietary regimens. Reference information is included in tabular form throughout the text and in 4 appendices; literature citations are appended to each of the 3 principal sections of the text.

O143

Nutrition statements of the National Health and Medical Research Council. Canberra, Australia : Australian Government Publishing Service. Journal of food & nutrition. 1986. v. 42 (3). p. 146-147. charts. (NAL Call No.: DNAL FNC 389.9 AU73).

Abstract: The One-hundredth session of the Australian National Health and Medical Research Council resulted in the incorporation of the recommended dietary intakes for the following nutrients: magnesium, iron, calcium and vitamin A. A table of acceptable weights-for-height was recently adopted and definitions for obesity and overweight were proposed. Infant feeding was considered regarding the marketing of unconventional foods for infants and the discouragement of the use of skim milk and goat milk. Hair analysis techniques marketed by commercial businesses for nutritional assessment were criticized as scientifically unacceptable by the Council. (lsp)

0144

Nutritional status, birth weight and breast feeding of elementary first grade Chilean students.

NURIB. Ivanovic, D. Ivanovic, R -Butron, C. Stoneham, Mass. Butterworth Publishers Nutrition reports international Dec 1987 v 36 (6) p 1347-1361, ill., charts. Includes 31 references (NAL Call No. DNAL RC620.A1N8).

Extract The purpose of this study was to assess the nutritional status of elementary first grade Chilean students from different socioeconomic status (SES) and to determine the influence of birth weight and the duration of breast feeding over it. A random sample of 306 students from public and private schools, of both sexes and from high, medium and low SES, was chosen from the Metropolitan Area of Santiago, Chile. Results showed a high prevalence of overweight (23.9%) and obesity (8.9%) and very low incidence of undernutrition (1%), expressed as percentage of OMS standard weight for height. In relation to Waterlow classification, the greater part of the students presented an adequate nutritional status (85.0%), a smaller proportion 0.3% was moderately or severely wasted and only 0.7% was seriously malnourished. Students from high SES registered a significantly better nutritional status and birth weight than low SES but lesser duration of breast feeding. Moreover, nutritional status was significantly and positively correlated with birth weight and not with duration of breast feeding. Results confirm the nutritional status improvement in Chilean school population (author)

0145

Obesity a family matter: Creating new behavior.

JADAA Frankle, Reva T Chicago, Ill. The Association. Journal of the American Dietetic Association, May 1985, v. 85 (5) p. 597-602, ill., charts. Includes 63 references. (NAL Call No. DNAL FNC 389.8 AM34).

Extract The family experience has a significant effect on overweight/obesity among family members. Success may depend on the extent of family functioning and on finding a support system within and/or outside the family. This aspect of weight control is considered in the nonmedical weight control program described. (author)

0146

Obesity among children: it's growing bigger.

New York, N.Y. The Letter, Tufts University diet & nutrition letter, Nov 1987, v. 5 (9), p. 7. (NAL Call No. DNAL TX341.T83).

Abstract. A health report concerning the increase in obesity prevalence in the US notes that measurements of fat deposits in children 6-11 years of age indicate that obesity prevalence in this age group may have increased up to 54% from 1963 to 1980, with cases of gross obesity rising almost 100%. Major

increases in obesity and gross obesity prevalence in adolescents also are noted. The characteristics of such children and the associated physical and psychological risks they face are discussed. Constructive strategies for reducing body weight in obese children also are described (wz)

0147

Obesity and atherosclerosis as consequences of early weaning.

Hann, P. New York Raven Press Nestle nutrition workshop series. Contributions from 10th Nestle Nutrition Workshop "Weaning Why, What, and When?" December 1987, New Delhi, India 1987, v. 10 p 93-113, ill., charts. Includes 62 references. (NAL Call No. DNAL RC620.A1N47).

Abstract A technical review presents a variety of data and experimental results (principally from animal studies and limited human studies) to indicate that early prenatal and postnatal nutrition and other factors have lasting effects on subsequent infant development. The review focuses on 2 pathological states thought to be influenced by early nutrition, viz. atherosclerosis and obesity. Specific attention is given to the late effects of early nutritional changes. The review includes 24 figures and 2 tables. A brief commentary on this review is appended. (wz)

0148

Obesity and food intake in children: evidence for a role of metabolic and/or behavioral daily rhythms.

Bellisle, F., Rolland-Cachera, M.F -Deheeger, M -Guilloud-Bataille, M London Academic Press, Appetite, Oct 1988 v 2 (2), p 111-118, charts. Includes 27 references. (NAL Call No. DNAL QP141.A1A64)

Abstract The distribution of daily energy intake was studied in 339 French children age 7-12 years. The results were compared among groups of different weight status from lean to obese. Obese and fat children ate less at breakfast and more at dinner than leaner peers.

0149

Obesity and television viewing in children and adolescents.

NUREA, Washington, D.C. : Nutrition Foundation. Nutrition reviews, Jan 1986, v. 44 (1), p. 9-12, charts. Includes 10 references. (NAL Call No. DNAL FNC 389.8 N953).

Abstract. This paper discusses the limitations of a recently published pediatric study that found a significant correlation between television watching and obesity, particularly in 12-17 year-old children. Limitations included the use of data collected in the 1960s (NHANES), the definition of obesity based solely on triceps skinfold, the lack of energy expenditure data, and the development of data from different sources (children or parents). Additional studies are needed that overcome these limitations, use more recent data, and that are designed to determine whether it is television and

consequent inactivity that cause obesity, or whether obese children choose to watch television (wz)

0150

Obesity in boys the distinction between fatness and heaviness

Griffiths M., Rivers J.P.W., Hoinville, E.A. London, Eng. U. Libbey Human nutrition clinical nutrition July 1985 v. 39C (4) p. 259-269 ill., charts. Includes 10 references (NAL Call No. DNAL TX341 H8)

Abstract A longitudinal anthropometric study of 2350 preschool boys revealed no definitive association between obesity (triceps skinfolds) and the development of heaviness (weight-for-height ratios). The results of this study indicate that in childhood obesity studies it is important to identify 3 categories: children who are heavy but not fat, children who are fat but not heavy, and children who are both fat and heavy. The study results are statistically summarized and their implications are discussed (wz)

0151

Obesity in children.

Brooke C.G., Abernethy, London, Eng. U. Libbey Human nutrition applied nutrition Aug 1985 v. 39A (4) p. 304-314 ill. Includes 4 references (NAL Call No. DNAL TX341 H82)

Abstract A literature review examines the assessment: organic causes, genetic causes, consumption pattern, energy expenditure, psychological causes, physiological effects, and treatment, of obesity in childhood. Due to the difficulties in ensuring the reduction of food intake, the performance of regular exercise, the motivation to lose weight, and patient compliance, the authors do not recommend treatment of otherwise normal children with mild-to-moderate obesity who are not motivated to lose weight. Besides reducing food consumption, increasing exercise, and therapy with certain thermogenic drugs, psychiatric referral and hospitalization also are discussed as therapeutic approaches (wz)

0152

Obesity in offspring of diabetic Pima Indian women despite normal birth weight.

DICAD2. Pettitt, D.J.; Knowler, W.C.; Bennett, P.H.; Aleck, K.A.; Baird, H.R. Alexandria, Va. American Diabetes Association Diabetes care, Jan/Feb 1987, v. 10 (1) p. 76-80 ill. Includes 20 references (NAL Call No. DNAL RC660.A1D53)

Abstract A clinical study evaluated the relative influences of birth weight and maternal diabetes on the development of obesity in the offspring of Pima Indian women. The offspring were grouped among 3 age groups (5-9, 10-14, and 15-19 years old) for each of 3 categories of maternal diabetes status (diabetic, non-diabetic, pre-diabetic). Birth weight was not predictive of subsequent obesity for offspring of diabetic and

pre-diabetic mothers, although offspring of diabetic mothers were heavier than those of non- and pre-diabetic mothers, independent of birth weight. (wz)

0153

Obesity in the school age child.

Summer S.K. Denver, Colo. American School Food Service Association School foodservice research review Fall 1986 v. 10 (2) p. 82-86. Includes 32 references (NAL Call No. DNAL TX945 S344)

Extract Factors associated with development of child and adolescent obesity, long-term health implications, and treatments for successful management are reviewed. Comprehensive programs including behavior modification, nutrition education, and physical activity are effective in the treatment of mild to moderate obesity in children. The school setting is appropriate for treatment and prevention programs (author)

0154

Onset of obesity in a 36 year birth cohort study

BMJOAE Braddon, F.E.M.; Rodgers, B.; Wadsworth, M.E.J.; Davies, J.M.C. London. British Medical Association. British medical journal, Aug 2, 1986, v. 293 (6542) p. 299-303, charts. Includes 36 references (NAL Call No. DNAL 448 B 877)

Abstract Data on the change in prevalence of obesity with age and for individual differences in the course and pattern of obesity are presented from a longitudinal study designed to assess the predictability of childhood obesity in adult life for a large cohort of children who were followed from birth to 36 years of age. The study revealed that only 21 per cent of the obese 36-year-olds were obese at age 11. The implications of the observed inaccuracy in the prediction of adult obesity from childhood obesity and the need for developing preventive measures are discussed (wz)

0155

Oxygen uptake and energy output during walking of obese male and female adolescents

AJCN A Katch, V.; Becque, M.D.; Marks, C.; Moorehead, C.; Rocchini, A. Bethesda, Md. American Society for Clinical Nutrition. American journal of clinical nutrition, Jan 1988, v. 47 (1), p. 26-32 ill., charts. Includes 21 references. (NAL Call No.: DNAL FNC 389.8 J824).

Extract Oxygen uptake and steady-rate energy output of 7 obese male and 13 obese female adolescents (greater than 178% ideal body weight) walking at four different speeds (1.167, 1.5667, 1.7833, and 2.125 m/s) were studied. Body composition was measured by hydrostatic weighing, and steady-rate energy output by open circuit spirometry. Energy output was expressed as kJ/min (kcal/min) and indexed to body mass and fat-free mass. A 2-by-4 ANOVA (sex by speed) revealed significant differences

in the energy output between the speed conditions. There was no significant difference between the sexes. A non-linear increase in caloric output with increasing speed indicated a decreasing efficiency with increasing speed of walking. Possible reasons include biomechanical factors such as increased upper-body forward lean needed to maintain balance at faster speeds of movement; increased energy output due to increased inertial extra energy output needed to accelerate the limbs and torso; and increased body fat (author)

0156

Pediatric hypertension

AJCCA. Menta, S.K. Chicago. American Medical Association. American journal of diseases of children. Aug 1987. v. 141. (8) p. 893-894. Includes 10 references. (NAL Call No. DNAL FNC 448 8 AM38)
Abstract: Due to the dissatisfaction of physicians with the high blood pressure values reported in the 1970 report of the National Heart, Lung and Blood Institute's (NHLEI) Task Force on Blood Pressure Control in Children as well as the report's nonconsideration of weight and height, and lack of data for children younger than 2 years, the NHLEI published new recommendations in January 1987. These new recommendations were based on a larger number of children and results were based on age, height and weight. The 95th percentile blood pressure levels were considerably lower than in the previous reports. Blood pressure values of children younger than 2 years were also included. Use of the phase V of Korotkoff sound (muffling of sound) for the diastolic blood pressures could be helpful during the transition from childhood to adolescence. In conclusion, the new NHLEI report offers a unique opportunity to study blood pressure prospectively in children and adolescents.

0157

Perception of eating and exercise in children as a function of child and parent weight status.

Epstein, L.H.; Valoski, A.-Wing, R.R.; Perkins, K.A.; Fernstrom, M.-Marks, E.-McCurley, J. London. Academic Press. Appetite. Apr 1989. v. 12 (2). p. 105-118. charts. Includes 45 references. (NAL Call No. DNAL QP141.A1A64).
Abstract: The present study assessed the effects of child body weight (obese/lean) and familial loading for obesity (two obese parents/two lean parents) on the psychophysics of sweetness, fatness and exercise workloads, as well as subjective ratings of foods varying in sugar and fat and activities varying in energy expenditure. Children were measured in a baseline state and at 6 months after the obese children had participated in a family-based behavioral weight control program. No differences between obese and lean children in perceptual ratings were observed. However, offspring of obese or lean parents differed on intensity rating, food palatability and activity enjoyment ratings. Intensity

ratings for sweetness in offspring of obese parents was increased, with a similar trend for intensity ratings for fatness. Offspring of obese parents rated all foods and activities with lower palatability and enjoyment ratings than offspring of lean parents. After a 6-month family-based behavioral weight loss treatment, obese children had significant decreases in percent overweight while lean children remained stable. Changes in the pattern of food ratings were observed for the obese children, with a reduction in liking for foods high in fat and/or sugar, and an increase in ratings for foods lower in fat and sugar. The effects of parental obesity on food and exercise intensity ratings and hedonic ratings were maintained. Overall, these results suggest parental weight influences behavioral factors related to obesity in children.

0158

Persistence of juvenile-onset obesity over eight years: The Bogalusa Heart Study

Freedman, D.S.; Snear, C.L.; Burke, G.L.; Srinivasan, S.R.; Webber, L.S.; Harsha, D.W.; Berenson, G.S. Washington, D.C. American Public Health Association. American journal of public health. May 1987. v. 77 (5) p. 588-592. ill., charts. Includes 36 references. (NAL Call No. DNAL 449.9 AM3U).
Extract: The persistence of obesity and overweight over eight years was assessed in a biracial (Black-White) cohort of 1,490 two- to 14-year-olds. Initial levels of triceps skinfold thickness (TRSF) and Rohrer index (weight/height³) were moderately predictive of subsequent levels ($r = 0.54$ and 0.67 , respectively). However, TRSF and Rohrer index tended to track most strongly in Black females ($r = 0.64$ and 0.72) and less well in both White females ($r = 0.45$ and 0.57) and preschool children ($r = 0.45$ and 0.54). Based on elevated levels of TRSF or Rohrer index, children were classified as obese or overweight, respectively. Of the 222 children who were initially above the 85th percentile for TRSF, 43 per cent remained obese after eight years. Persistence of overweight was slightly greater at follow-up, with 50 per cent of initially overweight children staying above the 85th percentile for Rohrer index. Severe, initial obesity/overweight (greater than 95th percentile) and consecutively elevated levels increased the probability of remaining obese/overweight. Results indicate that moderate, juvenile-onset obesity is malleable, but that the child who is extremely obese over consecutive examinations is likely to become an obese adult. (author)

0159

Persistently low blood retinol levels during and after parenteral feeding of very low birth weight infants: examination of losses in intravenous administration sets and a method of

prevention by addition to a lipid emulsion

PEDIAU. Greene H L., Phillips, B L., Franck, L., Fillmore, C M., Said H M., Murrell, J E., Moore, M E C., Briggs R. Elk Grove Village, Ill. American Academy of Pediatrics. Pediatrics June 1987. v. 79 (6). p. 894-900. ill., charts. Includes 20 references. (NAL Call No. DNAL FNC RJ1 P42)

Extract: Very low birth weight infants have little storage of hepatic retinol and are, therefore, highly dependent upon an exogenous supply. The recent association between low serum retinol level and bronchopulmonary dysplasia and the persistently low serum levels of retinol during total parenteral nutrition prompted a prospective study to evaluate serial changes in serum retinol levels during 1 month of total parenteral nutrition (retinol dose 455 micrograms/d) and again during 1 month of total enteral feeding (retinol dose 200 to 300 micrograms/d) in the same infants. Infants were divided into two groups. Group 1 consisted of infants weighing less than 1,000 g (n = 24) and group 2 consisted of infants weighing 1,000 to 1,500 g (n = 17). Although initial mean levels of retinol were similar in both groups (14.8 +/- 0.9 and 13.5 +/- 0.7 micrograms/dL), there was wide variation between infants. In group 1 infants, there was a significant (P less than .01) decline in retinol level by the second week of life (to 9.2 +/- 1 microgram/dL), which persisted during total parenteral nutrition, but increased to 13.4 +/- 2 after 1 week of enteral feeding. This level was maintained throughout enteral feeding. In group 2 infants, there was no significant change in serum retinol level throughout the study. During total parenteral nutrition, several infants had retinol levels below 10 micrograms/dL, a level associated with signs of retinol deficiency in older children. Because losses of retinol are known to occur in smaller volume total parenteral nutrition solutions, it was speculated that losses of retinol in our patients were due to retinol losses in the total parenteral nutrition delivery system. Because lipid might prevent these losses, in vitro studies were done to compare retinol losses in total parenteral nutrition v the usual daily dose of Intra-lipid. Results indicated 80% loss of retinol in total parenteral nutrition compared to only 10% loss in Intralipid. Lipid stabil

0160

Physical activity and fitness in obese children.

IJOBOP. Huttunen, N.P. ; Knip, M., Paavilainen, T. London : John Libbey & Company. International journal of obesity. 1986. v. 10 (6). p. 519-525. charts. Includes 27 references. (NAL Call No. : DNAL RC62B.A102).

Abstract: A study assessed whether obese children are less active and have a lower working capacity than normal weight children, and examined the role of physical activity in overweight

reduction and the possible consequences of weight reduction on physical fitness. It was concluded that obese children are less physically fit than normal weight children. It was observed, however, that weight reduction in obesity improves the maximum oxygen consumption towards normal. The study involved 31 obese and 31 normal weight children (ages ca 6-16 years). (wz)

0161

Physiology of fat absorption

Watkins, J B Boston Butterworths. c1987. Pediatric nutrition: theory and practice / edited by Richard J. Grand, James L. Sutphen, William H. Dietz. p. 127-137. ill., charts. Includes 29 references. (NAL Call No. DNAL RJ206 P42)

Abstract: A technical report stresses aspects of fat digestion and uptake relative to clinical conditions commonly encountered in the infant and young child, and highlights new advances in the understanding of the processes of lipid digestion that relate to patho-physiological aspects of fat malabsorption. Following a brief historical review of fat digestion and uptake, attention is focused on the forms of dietary lipids; the sources and activity of principal lipid-digestive enzymes; the physico-chemical states of lipids during fat digestion; fat absorption related to the intraluminal environment in infants; and intracellular events affecting triglyceride and cholesterol absorption. Schematics illustrating various aspects of lipid digestion, metabolism, and transport are included. (wz)

0162

Plasma cholesterol levels of 6585 children in the United States: results of the know your body screening in five states.

PEDIAU. Resnicow, K. ; Morley-Kotchen, J., Wynder, E. Elk Grove Village, Ill. American Academy of Pediatrics. Pediatrics Dec 1989. v. 84 (6). p. 969-976. charts. Includes 39 references. (NAL Call No. : DNAL FNC RJ1.P42).

Abstract: Medical and public health recommendations regarding detection and treatment of hyperlipidemia in children have generally been based on two principal epidemiologic studies: the Lipids Research Clinics Population Study (1972 to 1976) and the Bogalusa Heart Study (1973 to 1974). The present study was initiated to further describe the distribution of plasma cholesterol levels in a multiracial sample of American children. Between 1984 and 1988, the total cholesterol levels of 6585 children from 22 schools were measured as part of the Know Your Body School Health Program. For the entire population, ages 5 to 18 years, the mean total cholesterol concentration was 166.4 mg/dL. Total cholesterol was significantly higher in girls (168 mg/dL) than in boys (165 mg/dL), although sex differences were inconsistent across race/ethnicity. The mean value for blacks, 173 mg/dL, and

Hispanics, 168 mg/dL, was higher than for Asians 165 mg/dL, and whites 163 mg/dL. Across race/ethnicity, values tended to peak between ages 8 and 10 years for girls and approximately 10 years of age for boys. These values are slightly higher than those reported in the Lipids Research Clinics and Bogalusa Studies. Public health implications of these findings are discussed.

O163

Plasma vitamin E values at birth and in 3 and 6 month old infants.

Labadarios, D., Shephard, G.S., Hesseling, P.B., Hutchison, M.E. London. *John Libbey Journal of nutrition, growth and cancer* 1985 v. 2 (1) p. 15-23. ill., charts. Includes 26 references. (NAL Call No. DNAL OP141.A1J685)

Abstract: A clinical study was designed to assess the normal level of plasma vitamin E (alpha-tocopherol (AT)) in newborns, and in infants 3 and 6 months of age in the urban Black population of Cape Town, South Africa. Mean plasma AT levels in newborns was found to be 3.2 mg/L, and for infants 3 and 6 months of age was 10.8 and 9.2 mg/L, respectively. At these 3 respective ages (0, 3, and 6 months), mean plasma cholesterol was 1.6, 2.9, and 3.3 mmol/L, respectively, and mean plasma triglyceride were 0.4, 1.4, and 1.3 mmol/L, respectively. Statistically significant correlations were found between plasma AT and cholesterol, between AT and triglycerides, and between AT and cholesterol plus triglycerides, but no correlation was found between maternal plasma AT and plasma AT in newborns. The implications of these findings are discussed. (WZ)

O164

Potassium, magnesium, and calcium balance in obese adolescents on a protein-sparing modified fast.

AJCN. Stallings, V.A.; Archibald, E.H.; Pencharz, P.B. Bethesda, Md.; American Society for Clinical Nutrition. *American journal of clinical nutrition*. Feb 1988, v. 47 (2), p. 220-225. ill., charts. Includes 33 references. (NAL Call No.: DNAL FNC 389.8 J824).

Extract: Ten obese adolescents consumed the protein-sparing modified fast (PSMF), a high-protein, low-carbohydrate diet, for 92 +/- 19 d and lost 14.7 +/- 5.3 kg. The effect of weight loss using the PSMF on potassium, magnesium, and calcium was evaluated using balance method at days 2-4 and 12-20. RBC-Mg, RBC-K and total body K (TBK). The vitamin- and mineral-supplemented PSMF allowed positive Ca and K balance and improved Mg balance (p less than 0.005). TBK decreased significantly, 118.7 +/- 13.7 to 97.4 +/- 1.0 g (3.04 +/- 0.35 to 2.50 +/- 0.03 mol) from baseline to 90 d (p less than 0.001), with no change in the RBC-K concentration. An unexpected finding was a significant decrease in RBC-Mg, 3.2 +/- 1.1 to 1.8 +/- 0.3 mmol/L (p less than 0.001) after 60-90 d on the diet despite maintenance of normal serum Mg level (author).

O165

Potential history of cardiovascular disease as an indication for screening for lipoprotein abnormalities in children.

JOPDA. Dennison, E.A.; Kikuchi, D.A.; Srinivasan, S.R.; Webber, L.S.; Berenson, G.S. St. Louis, Mo. C.V. Mosby. *The Journal of pediatrics* Aug 1989, v. 115 (2), p. 186-194. charts. Includes 31 references. (NAL Call No. DNAL FNC RJ1.A453).

Abstract: We studied the relationship between parental history of cardiovascular disease and risk for adverse lipid and lipoprotein levels in a total community study of 3313 children (ages 4 to 17 years, 63% white, 37% black). Older white children (11 to 17 years) with a parental history of heart attack or diabetes were 4.3 and 5.6 times, respectively, more likely to have high levels (greater than or equal to 95th percentile) of serum total cholesterol than those without such a history (all p less than 0.05). White children with a parental history of heart attack or diabetes were twice as likely to have an elevated (greater than or equal to 95th percentile) low-density lipoprotein cholesterol (LDL-C) level than those without such a history (both p less than 0.05). In contrast, parental history of cardiovascular disease did not predict elevated levels of total cholesterol or LDL-C in black children. However, older black children with a parental history of heart attack, hypertension, or diabetes were 4 1/2 to 5 times more likely to have low levels (less than or equal to 5th percentile) of high-density lipoprotein cholesterol than those without such a history (all p less than 0.05). Only 40% of white children and 21% of black children with elevated LDL-C levels had a parental history of vascular diseases. These findings raise questions about the current practice of screening only children with a family history of cardiovascular disease to identify those with elevated total cholesterol and LDL-C levels.

O166

Preliminary report from a conference entitled "Prevention of Adult Atherosclerosis During Childhood."

JOPDA. LaRosa, J.; Finberg, L. St. Louis, Mo.; C.V. Mosby. *The Journal of pediatrics*, Feb 1988, v. 112 (2), p. 317-318. (NAL Call No. DNAL FNC RJ1.A453).

Abstract: A preliminary report from the Ross Conference which was held on September 20 to 22, 1987, reviewed and discussed the role of diets in children and adults in producing arterial lesions that cause coronary occlusions. There was widespread agreement, for adult populations, that not only is lowering of low density lipids desirable for the genetically high-risk population, but a general population strategy for diet change is also warranted; attendees agreed that fat intake over 40% of calories is excessive for the general population of children. There was

unanimous agreement on the appropriateness of advising a life-style for children and adolescents that includes not only a prudent diet, but also regular exercise and no smoking. Four positions, broadly agreed to by the participants, are included (nan)

0167

Prevention of adult atherosclerosis during childhood report of the 95th Ross Conference on Pediatric Research. Columbus, Ohio (625 Cleveland Ave Columbus 43216) Ross Laboratories, c1988 "August 1988"--P 4 of cover viii, 130 p ill., 22 cm Includes bibliographies. (NAL Call No. DNAL RC692 R6 1987)

Abstract This conference report discusses dietary and lifestyle changes in childhood that have been proposed as ways to prevent atherosclerosis in later life. Evidence that atherosclerosis begins in childhood and that it can be modified by dietary changes was presented. Topics include cholesterol metabolism, evolution of the atherosclerotic process in childhood diet and lifestyle recommendations, cholesterol screening in children, changing diets to lower fat and cholesterol, and, obesity, physical activity and smoking.

0168

Primary prevention of atherosclerosis in childhood: the role of lipids / edited by Robert I. Levy.

; Levy, Robert I., 1937-. New York, NY: Biomedical Information Corp., c1985. Proceedings from a videoconference. New York City, March 28, 1985. 54 p. ill., 26 cm. Includes bibliographies. (NAL Call No. DNAL RJ426.A82P7).

Abstract This collection of papers is concerned with the role of lipids in the prevention of atherosclerosis in childhood. Paper topics include lipids, lipoproteins and lipid transport; distribution of plasma lipid concentrations in infancy and childhood, cholesterol and atherosclerosis, lessons from clinical trials in humans; and, modification of blood cholesterol levels in a school-based population.

0169

Progeny's lipid and lipoprotein levels by parental mortality.

CIRCAZ, Glueck, C.J., Laskarzewski, P.M.-Suchindran, C.M.-Chambless, L.E.-Barrett-Connor, E.-Stewart, P.-Heiss, G.-Tyroler, H.A. Dallas: American Heart Association, Inc. Circulation, Jan 1986 v. 73 (1,pt.2) p. I-51-I-61. ill., charts. Includes 61 references. (NAL Call No.: DNAL RC681.A1C8).

Abstract. An hypothesis that parental mortality attributed to cancer or heart disease in adults is predictive of lipid and lipoprotein (LP) levels in the progeny of adults was examined using data from over 6000 participants in the Lipid Research Clinics study. Most of the significant correlations were found for parent-son pairs, with sons showing higher plasma cholesterol and

low-density LP cholesterol levels who had fathers die of heart disease before age 60. Maternal heart disease mortality before 60 was associated with lower high-density LP cholesterol. Maternal and paternal mortality from cancer before 60 was associated with higher triglyceride levels in sons (wz)

0170

Protein-sparing diet for severely obese adolescents. Design and use of an equivalency system for menu planning JADAA Bell, Louise ; Chan, Linda-Pencharz, Paul B. Chicago, Ill: The Association Journal of the American Dietetic Association, Apr 1985 v. 85 (4) p. 459-464. charts. Includes 18 references. (NAL Call No. DNAL FNC 389 B AM34)

Extract A protein-sparing, low-energy diet is suitable treatment for severely obese adolescents. The dietary regimen requires adequate protein and fluid, plus nutrient supplements. Meal planning is simplified by the use of a protein equivalency system developed for this diet. Recent experience with 21 adolescents resulted in satisfactory weight control. (Author)

0171

Prudent life-style for children: dietary fat and cholesterol.

PEDIAU Finberg, L. ; Dweck, H.S.-Holmes, F.-Kretchmer, N.-Mauer, A.M.-Reynolds, J.W.-Suskind, R.M.-Benson, J.D.-Miguel, S.G.-Purvis, G.A. Elk Grove Village, Ill: American Academy of Pediatrics, Pediatrics, Sept 1986 v. 78 (3), p. 521-525. Includes 29 references. (NAL Call No.: DNAL FNC RJ1.P42).

Abstract. A brief report by the Committee on Nutrition of the American Academy of Pediatrics discusses ancillary findings of the 1983 Committee's statement, "Toward a Prudent Diet for Children," with respect to dietary fat and cholesterol. Particular attention is given to coronary artery disease risk and hypercholesterolemia, the advantages and disadvantages of decreasing serum cholesterol levels, and to recommendations for changes in the infant diet. It is argued that there is no direct evidence for prospective studies that dietary recommendations proposed by the NIH Consensus Development panel for subjects older than 2 years of age will be effective in decreasing serum cholesterol during the first 2 decades of life, or that they will adequately support growth and development. Recommendations include infant feeding practices, use of a varied diet, screening for obesity and hazardous cholesterol and lipoprotein profiles, and the use of optimal total fat intakes at 30-40% of calories. (wz)

0172

Relation of body fat distribution to hyperinsulinemia in children and adolescents: the Bogalusa Heart Study.

AJCNA. Freedman, D.S. ; Srinivasan, S.R.-Burke, G.L.-Shear, C.L.-Smoak, C.G.-Harsha, D.W.-Webber, L.S.-Berenson,

G S Bethesda, Md American Society for Clinical Nutrition. American journal of clinical nutrition. Sept 1987. v. 46 (3). p. 403-410 ill., charts. Includes 45 references (NAL Call No. DNAL FNC 389.8 J824).

Extract The relation of body fat distribution to plasma levels of glucose and insulin during an oral glucose tolerance test was examined in 355 Black and White school-age children. Both central and peripheral fat were similarly related to fasting, 30-min, and 1-h glucose. Unlike peripheral fat, central body fat was more strongly related to the 1-h insulin response ($r = 0.35$ vs 0.26). This association remained significant for central fat independent of peripheral fat ($r = 0.18$). The strong relation of central fat to insulin response was noted in both races and sexes but not in either sexually immature or relatively thin children. These findings indicate that, even in early life, a central body fat pattern relates positively to insulin response to glucose load. Thus, knowledge of body fat localization may help identify persons most susceptible to hyperinsulinemia in early life. (author)

0173

Relation of body fat patterning to lipid and lipoprotein concentrations in children and adolescents: the Bogalusa Heart Study.

AJCN. Freedman, D.S. ; Srinivasan, S.R.-Harsha, D.W.-Webber, L.S.-Berenson, G S. Baltimore, Md. : American Society for Clinical Nutrition. American journal of clinical nutrition. Nov 1989. v. 50 (5) p 930-939. charts. Includes 48 references. (NAL Call No. DNAL FNC 389.8 J824).

Abstract: Although a truncal distribution of adipose tissue in adults is associated with several metabolic complications, its importance in early life has received little attention. The relation of several anthropometric measures to serum concentrations of lipids, lipoproteins, and apolipoproteins was therefore examined in 361 children who were between ages 6 and 18 y. (Children had been selected previously because of extreme levels of very-low-density- and low-density-lipoprotein cholesterol.) Analyses revealed two groups of anthropometric variables: truncal measures (waist circumference and subscapular, subcostal, and suprailliac skin-fold thicknesses) and thickness of peripheral skinfolds (femoral, triceps, calf, and biceps). After generalized obesity was adjusted for children with high concentrations of both cholesterol fractions had more truncal fat but less peripheral fat than did children with low lipoprotein cholesterol concentrations. A truncal fat pattern was also associated with decreased concentrations of high-density-lipoprotein cholesterol and apolipoprotein A-1. Knowledge of fat patterning may help identify persons prone to hyperlipidemia

0174

The relationship among attitudes, behaviors, and biomedical measures of adolescents "at risk" for cardiovascular disease.

Adeyanju, M. ; Creswell, W.H. Jr Kent, Ohio American School Health Association Journal of school health Oct 1987 v. 57 (8) p 326-331 charts Includes 21 references (NAL Call No DNAL LB3401.J6).

Abstract A 4-year study assessed trends in biomedical measures, health behavior, and attitudes in 93 high school students clinically classified as being at risk for subsequent cardiovascular disease. Most subjects reported negative health behavior changes in smoking, diet, alcohol usage, and stress, although smokeless tobacco use declined with age and the students acknowledged the potential harmful effects of smoking. A positive correlation was found among health status, attitudes, and behavior

0175

Relationship of changes in obesity to serum lipid and lipoprotein changes in childhood and adolescence.

JAMA. Freedman, D.S. ; Burke, G.L.-Harsha, D.W.-Srinivasan, S.R.-Cresanta, J.L.-Webber, L.S.-Berenson, G.S. Chicago : American Medical Association JAMA : Journal of the American Medical Association. July 26, 1985. v. 254 (4). p. 515-520. ill., charts. Includes 47 references. (NAL Call No. DNAL FNC 448.9 AM37).

Abstract A 5-year longitudinal study of the relationship of changes in triceps skin-fold thickness (TSFT) to changes in the levels and profiles of serum lipids (cholesterol, triglycerides, lipoproteins) is reported for a group of 1598 children who were examined initially at 5-12 years of age. Positive age independent correlations were found between TSFT changes and serum level changes of total cholesterol, triglycerides, and low-density and very-low-density lipoprotein cholesterol, while weaker (but significant) inverse relationships were found between TSFT changes and serum changes of high-density lipoprotein cholesterol. The implications of these and related findings are discussed. (WZ)

0176

Relationships between anthropometrical measurements, lipemia and apolipoproteinemia in a sample of Madrid secondary school adolescents.

NURIB. Goni, I. ; Garcia-Diz, L.-Cuesta, C.-Sanchez-Muniz, F.J.-Castro, A. Stoneham, Mass. Butterworth. Nutrition reports international. June 1989. v. 39 (6). p. 1167-1175. charts. Includes 24 references. (NAL Call No.: DNAL RC620.A1N8).

Abstract: Sixty two adolescents, 27 boys and 35 girls, participated in a study to evaluate the relationships between anthropometrical measurements, lipemia and apolipoproteinemia. None of adolescents had a body mass index (BMI) higher than 25. Lipemia and apoproteinemia patterns were similar to

those described as low risk levels for cardiovascular disease in subjects younger than 20. BMI showed a very significant correlation with weight, fat percentage and body fat content in boys and girls. In boys Total Cholesterol (TC) and Triglycerides (Tg) showed a very significant correlation (.47 and .48 respectively) to fat percentage. Tg levels were also highly correlated with body fat content (.54). No consistent correlations were seen between body fat or fat percentage to TC or Tg, however a significant correlation (-.35) was found between fat percentage and apoprotein (apo) AI/apo B ratio in girls. Apo B had a significant and a very significant correlation (.38 and .60) with TC in boys and girls respectively.

O177

Relationships between obesity and nutrient intake of Chilean adolescents.
NURIE Ivanovic, D. Stoneham, Mass
Butterworth Publishers Nutrition
reports international, Feb 1988, v. 37
(2), p. 227-239. charts. Includes
30 references. (NAL Call No. DNAL
RC620 A1N8)

Extract: The relationships between obesity and nutrient intake were investigated in 521 adolescent schoolers graduating from elementary and high school, in the Metropolitan Area of Santiago, Chile. The sample included adolescents of both sexes, from public and private schools and from high, medium and low socioeconomic status (SES). Obesity was defined as a percentage of weight for height (W/H) greater than 120, according to CMS Tables. Standard procedures for 24-hour dietary recall interviews were used to collect data and adequacy of intake was assessed by FAO/OMS pattern. Results indicated that obesity affected mainly older adolescent females, with a slight major incidence in low SES and significantly associated with a low nutrient intake and percentage of height for age. This is the variable with the greatest explanatory power of the present nutritional status (W/H). In males no significant differences were found between nutrient intake of obese and nonobese. Results suggest that further researches are necessary to approach this problem from a multicausal context. (author)

O178

Resting metabolic rate in lean and obese children: relationship to child and parent weight and percent-overweight change.

AJCN. Epstein, L.H.; Wing, R.R.; Cluss, P.; Fernstrom, M.H.; Penner, B.; Perkins, K.A.; Nudelman, S.; Marks, B.; Valoski, A. Baltimore, Md.; American Society for Clinical Nutrition. American journal of clinical nutrition, Feb 1989, v. 49 (2), p. 331-336. charts. Includes 35 references. (NAL Call No. DNAL FNC 389.8 J824).

Abstract: Two studies were conducted to assess differences in metabolic rate as a function of child weight (study I), and the interaction of child and parent

weight (study II). In both studies obese children had higher resting metabolic rates (RMRs) than lean children (p less than 0.05). Child weight accounted for 72 and 78% of the variance in RMR in studies 1 and 2 respectively. Including parental weight did not improve the prediction of RMR. After 6 mo of treatment, obese children decreased percent overweight, whereas lean children showed no change (p less than 0.01). RMR in both groups remained unchanged after 6 mo. These results indicate that the RMR is higher in obese than in lean children, that changes in percent overweight that result from increases in height and no change in weight do not decrease RMR over 6 mo, and parent weight does not improve the prediction of child RMR.

O179

Retinol, beta-carotene and alpha-tocopherol status in a French population of healthy children.
IZVIAK, Malvy, J.M.D., Mourey, M.S.; Carlier, C.; Caces, P.; Dostalova, L.; Montagnon, E.; Amedee-Manesme, O. Bern
Hans Huber International journal for vitamin and nutrition research, 1989, v. 59 (1), p. 29-34. charts. Includes 32 references. (NAL Call No. DNAL 389.8 233)

Abstract: The authors studied a group of healthy children aged 1-16 living in France. Their blood cholesterol and fat-soluble vitamin levels were measured. Charts are included to show the relationship between vitamin levels and age group, and cholesterol levels and vitamin levels. The results of this study are discussed and compared with studies of children in underdeveloped countries.

O180

Role of fat distribution during growth and its relationship to health. Summary.
AJCNA. Bray, G.A.; Bouchard, C. Bethesda, Md.; American Society for Clinical Nutrition. American journal of clinical nutrition, Mar 1988, v. 47 (3), p. 551-522. (NAL Call No. DNAL FNC 389.8 J824).

Abstract: A brief summary report discusses the technical highlights of a 1987 meeting organized to discuss the current state of knowledge concerning the role of body fat distribution during child development on subsequent health. Topics discussed include: the relationship of body fat distribution to health risks; methods for measuring body fat and its distribution; the major patterns of body fat in young boys and adolescents, and hypotheses offered to explain the regional distribution of body fat. The potential for altering body fat distribution also is discussed. (wz)

O181

Routine cholesterol surveillance in childhood.
PEDIAU. Garcia, R.E.; Moodie, D.S. Elk Grove Village, Ill.; American Academy of Pediatrics. Pediatrics, Nov 1989, v. 84 (5), p. 751-755. Includes 26

references (NAL Call No DNAL FNC RJ1 P42)

Abstract Coronary heart disease is the leading cause of death in the United States and there is reason to believe that it begins in childhood. Evidence is accumulating that early diagnosis and treatment of hypercholesterolemia, a major coronary risk factor, can markedly reduce the incidence of atherosclerotic heart disease in later life. A pediatric group practice consisting of six pediatricians and a pediatric nurse practitioner performed a cholesterol surveillance study of 6500 children between 3 and 18 years of age. Parents and patients were counseled regarding other coronary risk factors, and the American Heart Association diet was recommended. According to the results of the study, 1251 children (19%) exceeded the acceptable 90th percentile for cholesterol and that 143 of 299 significantly hypercholesterolemic children (48%) had no family history of premature myocardial infarction or known hypercholesterolemia. The current recommendation is that only those children from high-risk families should be screened for an elevated cholesterol level. The authors conclude, as a result of this study, that all children older than 3 years of age should have a cholesterol test and that advice regarding avoidance of high-risk coronary life-style behaviors should be a routine part of pediatric anticipatory guidance.

0182

Safety and efficacy of long-term diet and diet plus bile acid-binding resin cholesterol-lowering therapy in 73 children heterozygous for familial hypercholesterolemia.

PEDIAU, Glueck, C.J. ; Mellies, M.J.; Dine, M.; Perry, T.; Laskarzewski, P. Elk Grove Village, Ill. American Academy of Pediatrics. Pediatrics. Aug 1986. v. 78 (2). p. 338-348. charts. Includes 46 references. (NAL Call No.: DNAL FNC RJ1.P42).

Extract. Our specific aim was to examine the efficacy and safety of long-term cholesterol-lowering diet and bile acid-binding resin therapy in 73 children heterozygous for familial hypercholesterolemia (FH). We prospectively followed accretion of height and weight in 40 FH children for 5.8 years on diet alone and in 33 FH children for 4.3 years on diet and bile acid-binding resins (8 to 20 g/d). In 67 of these 73 children, sequential data on plasma cholesterol lowering was obtained, including 32 children on diet plus bile acid-binding resins and 35 on diet alone. For all 73 children, median age, sex, and race-specific percentiles for height and weight at entry were 50 and 50, respectively, and, 5.7 years later, were unchanged at 50 and 50. Initial and final percentiles for height and weight were closely correlated. Percentile distributions for height and weight at entry into the study did not differ from those at the end of follow-up, in both the 40 FH children on

diet alone and the 33 on diet plus bile acid-binding resins. Tracking of height and weight did not differ in the 40 children on diet alone v the 33 on diet plus bile acid-binding resins. During 6 years of follow-up there were no significant differences in the percentage of serial, postbaseline measurements for height which were either less than or greater than or equal to baseline percentiles, comparing 40 FH children on diet alone, 33 FH children on diet plus resin, and 39 normal children (on ad libitum diet). FH children on diet or plus resin had a smaller percentage of weight measurements equal to or more than baseline percentiles than normals on follow-up probably reflecting restriction of total fat intake to less than 35% of calories. On diet alone, 32 FH children had total plasma cholesterol of 307 +/- 8 mg/dL (mean +/- SE); bile acid-binding resins were added to diet in these children at an average age of 11.5 years, with this regimen maintained for 4.6 +/- 0.4 years, leading to a mean reduction in

0183

School nurses' perceptions of childhood obesity.

Price, J.H. ; Desmond, S.M.; Ruppert, E.S.; Stelzer, C.M. Kent, Ohio : American School Health Association. Journal of school health. Oct 1987. v. 57 (8). p. 332-336. charts. Includes 31 references. (NAL Call No DNAL LB3401.J6).

Abstract A questionnaire survey of 250 of nurses randomly selected to assess their beliefs concerning childhood obesity found that most of the 220 respondents believed normal weight to be important to childhood health (85%) and that school nurses should set an example by maintaining a normal weight (77%). However, they appeared skeptical of children's ability to lose weight and of their potential influence in aiding children to do such, but they believe that schools should be more active in combating this problem.

0184

School promotion of healthful diet and exercise behavior: An integration of organizational change and social learning theory interventions.

Parcel, G.S. ; Simons-Morton, B.G.; O'Hara, N.M.; Baranowski, T.; Kolbe, L.J.; Bee, D.E. Kent, Ohio : American School Health Association. Journal of school health. Apr 1987. v. 57 (4). p. 150-156. charts. Includes 34 references. (NAL Call No.: DNAL LB3401.J6).

Abstract: A report discusses the rationale for approaching health promotion and cardiovascular disease risk reduction through schools, summarizes baseline findings, and presents the conceptual and theoretical bases for the interventions. The program, called "Go for Health," combines the use of a healthful diet, nutrition and physical education, the evaluation of classroom measures, and examinations of the nutrient content of the school lunch. The design

characteristics of the program are generated and justified (wz)

0185

Secular trends of obesity in early life the Bogalusa Heart Study

Shear, C L., Freedman, D S., Burke, G L., Harsha, D W., Webber, L S., Berenson, G S. Washington, D C. American Public Health Association. American journal of public health Jan 1988 v 78 (1) p 75-77. ill., charts. Includes 20 references. (NAL Call No. DNAL 449 9 AM3J)

Extract. Secular changes in height and weight measurements were examined in five- to 14-year-olds from 1973 to 1984. The age-sex specific 85th percentile was used to classify persons as overweight (based on ponderal index; kg/m³). Secular increases in weight (2.5 kg) and ponderosity (0.5-0.7 kg/m³) were found. Gains in ponderosity over the 11-year period were greater at the 75th percentile than at the 25th percentile, and the prevalence of overweight increased from 15 per cent to 24 per cent. (author)

0186

Self-esteem of adolescents enrolled in a weight reduction program

JADAA. Drake, M.A. Chicago, Ill. The Association. Journal of the American Dietetic Association. Dec 1988 v 88 (12). p 1581-1582. charts. Includes 9 references. (NAL Call No. DNAL FNC 389.8 AM34)

Abstract. The relationship between obesity and self esteem in female adolescents enrolled in a weight reduction program is examined.

0187

Self-selected dietary protein sources and serum lipid patterns of adolescent humans.

NURIB. Potter, S., Kies, C. Stoneham, Mass. Butterworth, Nutrition reports international. Nov 1989 v. 40 (5). p. 863-870. charts. Includes 21 references. (NAL Call No. DNAL RC620.A1NB).

Abstract: The objective of the current study was to investigate possible relationships between self-selected animal to vegetable protein intake ratios and serum lipid levels in adolescent humans. Negative correlations were found between blood serum lipids and the protein intake ratios. Negative correlations were also found between percent total fat and caloric intakes and serum lipids. These results were not consistent with those obtained from adult populations implying that the physiological demands of growth were possibly more influential on serum lipids than dietary components.

0188

Serum cholesterol (total, low-density lipoprotein, and high-density lipoprotein cholesterol), triglyceride levels, and fat consumption among Jerusalem Arab and Jewish schoolchildren.

Tamir, D.; Edelstein, P., Reshef, A., Halfon, S.T., Palti, H. Duluth, Minn.

Academic Press. Preventive medicine Nov 1987 v 16 (6) p 752-760. charts. Includes 29 references. (NAL Call No. DNAL RA421.P684)

Abstract. A clinical survey assessed fat consumption patterns and serum levels of total cholesterol, lipoprotein (LP) cholesterol fractions, and triglycerides (TG) in 643 Jewish and 198 Arab first- and second-grade school children in Jerusalem. While no significant differences were found in saturated fat and cholesterol intakes between the 2 ethnic groups, the Arab children had lower total, high-density LP, and low-density LP cholesterol levels and higher TG levels than the Jewish children. It is concluded that further studies are needed to explain these results (wz)

0189

Serum lipid levels in elementary and junior high school children and their relationship to relative weight.

Yamamoto, A., Horibe, H., Sawada, S., Uyama, M., Matsuzawa, Y., Yamamura, T., Yokoyama, S., Kasagi, F., Kameda, K. Duluth, Minn. Academic Press. Preventive medicine Jan 1988. v 17 (1) p 93-108. ill., charts. Includes 44 references. (NAL Call No. DNAL RA421.P684).

Abstract. A study assessed serum cholesterol, lipoprotein, and triglyceride levels in 2626 school children (ages 7-15) in 3 elementary schools and a junior high school in a suburban area of a city in Japan, historically known to have low coronary heart disease rates. Cholesterol levels increased with age in both sexes, but junior high boys had lower levels than girls of the same age. The effects of physical exercise on serum lipid composition also were observed. (wz)

0190

Serum lipids and lipoproteins.

PEDIAU. Freedman, D.S.; Srinivasan, S.R., Cresanta, J.L., Webber, L.S., Berenson, G.S. Elk Grove Village, Ill. American Academy of Pediatrics. Pediatrics. Cardiovascular Risk Factors from Birth to 7 Years of Age: The Bogalusa Heart Study. Nov 1987. v. 80 (5,pt.2). p. 789-796. ill., charts. Includes 48 references. (NAL Call No. DNAL FNC RJ1.P42).

Extract: Serum lipids and lipoprotein cholesterol fractions were examined in a newborn cohort that was followed from birth to 7 years of age. Although white and female infants had higher cord blood levels of high-density lipoprotein cholesterol (HDL-C) than did black and males infants, respectively, these differences did not persist throughout early childhood. Mean levels of all serum lipids and lipoproteins increased greatly in the first 6 months of life, and by 2 years of age, levels approached those seen in adolescents. Infants consuming cow's milk had higher 6-month levels of serum total cholesterol and low-density lipoprotein cholesterol than did formula-fed infants. However, milk source in infancy did not significantly

influence total cholesterol or low-density lipoprotein cholesterol levels at age 7 years. Serum lipid and lipoprotein levels at age 7 years were associated with previously measured levels as early as 6 months of age and infants with unfavorable levels were likely to have similar adverse levels at 7 years of age. In addition, increases in obesity between 6 months and 7 years of age were positively associated with increases in levels of serum triglycerides. These results suggest that certain persons at increased risk for cardiovascular disease can be identified in infancy. (author)

0191

Serum transport of cholesterol in adolescents in four different socioeconomic levels

NURIE Saitua, M.T., Ivanovic D. Los Altos, Calif. Genor-X, Inc. Nutrition reports international April 1985, v. 3, (4) p. 943-954. charts. Includes 23 references. (NAL Call No. DNA, RC62C A1N8)

Extract: The purpose of this study was to evaluate the serum transport of cholesterol in Chilean students. Total cholesterol (Total-C) and cholesterol in LDL and HDL were measured in 152 Elementary and High School students from Santiago, of both sexes and belonging to a high, medium, and low socioeconomic level (SEL), as determined by the Graffar Modified Scale. Total-C and LDL-C serum concentration were higher in females over sixteen years of age from high socioeconomic level. This same group had one of the lowest values for HDL-C in our sample, so LDL-C/HDL-C ratio is significantly higher in females over sixteen years of age, from high socioeconomic level. Our results confirm that socioeconomic level conditions a certain life style, that reflects in a lipoprotein pattern enhances the possible atherogenic risk. (author)

0192

Skinfold assessment as an intervention technique: one aspect of a nutrition education program.

JADAA Hunt-Pellow, J. Chicago, Ill. The Association, Journal of the American Dietetic Association, Mar 1986, v. 86 (3), p. 369-370. Includes 10 references. (NAL Call No., DNAL FNC 389.8 AM34). Abstract: Various interventions for nutrition education used by a registered dietitian at a small, rural school for grades 1-4 are described, including the monitoring of skinfolds in the 4th grade class over several school years to see whether body fat proportion would be altered over time. Obesity incidence in 4th grade girls and boys was about 54 per cent (7 girls) and 33 per cent (3 boys), respectively. A year later in the 5th grade, obesity incidence had dropped to 5 girls (ca. 38 per cent) and 2 boys (22 per cent). A year later, the number of obese girls and boys remained the same but their body weight gains were reduced. (wz)

0193

Socio-demographic differences in fat and sugar consumption patterns among Finnish adolescents.

ECFNE Prattala, R. London: Gordon & Breach Science Publishers. Ecology of food and nutrition 1988, v. 22 (1) p. 53-64. charts. Includes 35 references. (NAL Call No. DNAL TX341 E3)

Abstract: A representative sample (response rate 83%) of 3734 Finnish adolescents was surveyed on health habits by a mailed questionnaire in 1983. Questions included use of different milk types, butter, margarine, sweet pastry, coffee sugar, coffee, sweetened yogurt, candies and soft drinks. Food consumption patterns were analysed by age, sex, place of residence (rural/urban) and father's occupation and education using logistic regression analysis. Father's socio-economic status (SES) had a strong effect. The use of margarine, butter, high and low fat milks, and sweet pastry; age affected coffee, sweets and sweetened yoghurt, and sex affected coffee sugar and soft drinks. Children of farmers and lower socio-economic groups used high-fat milk, butter and sweet pastry more often than those of white-collar families. SES effect was stronger among the younger age groups. Coffee and soft drink usage increased by age, sweetened yogurt decreased, whereas use of candies was highest around age 15. Girls, especially in the older age groups, used less coffee sugar and soft drinks than boys. The SES effect on the use of foods consumed daily in Finnish homes was strong, whereas those adolescents buy outside home was independent of family background, indicating multidimensionality of food consumption and distinctions between adolescent and adult life styles.

0194

Some questions you may have about overweight children.

Oklahoma City, OK: Chickasaw Nation, WIC Program, 1987. Cover title. 3 p. ; 22 cm. (NAL Call No., DNAL E75.A5 no.A-34).

Abstract: A discussion of overweight children is presented in a question and answer format. Included is information on whether a fat child is healthy; why some children are overweight and others are not; and tips for parents on what to do if their child is overweight.

0195

SomeBODY's companion /by Lynda Corby and Patti Clark.

Corby, Lynda, 1949-; Clark, Patti, 1951-. Saskatoon, Saskatchewan: Fifth House Ltd., 1985. Companion to the authors' You're someBODY: how to be a slim kid. 44 p. ; 28 cm. Includes bibliographical references. (NAL Call No. DNAL jRU399.G6C61).

Abstract: A children's activity book and companion's guide are the major components of this weight management program for children. The book is designed for a parent-child partnership but can also be used by a variety of

professionals, including teachers, dietitians, nutritionists, home economists, and public health nurses. Self-discovery and the development of self-esteem are the two basic principles incorporated throughout the book. Self-discovery begins with a variety of record-keeping methods that children use: 1) to identify eating and exercise behaviors that contribute to overweight and 2) to develop a plan for change. Children's activities are self-explanatory for the most part; additional details and explanations are provided where necessary in the companion's guide. Appendices in the companion's guide include 1) children's food experience books, 2) nutrition and fitness resources for parents and children, 3) activities for children with handicapping conditions, and 4) addresses of federal and local resource groups.

O196

The stoplight diet for children and children / Leonard H. Epstein and Sally Squires, foreword by Jane E. Brody Epstein, Leonard H. Squires, Sally Boston. Little, Brown, c1988. xiii, 232 p. forms, 25 cm. (NAL Call No. DNAL RJ399.CCF6 1988).

Abstract: The stop-light diet is a comprehensive weight reduction diet for children which emphasizes nutrition, physical activity, behavior modification, and family teamwork. The approach to this diet enables parents to work with children to learn good eating habits. In the Stop-Light Diet Program, foods are classified by a color-coded system: green for very low calorie, nutritious, "all-you-can-eat" foods, yellow for foods moderate in calories and rich in nutrients, and red for foods which provide very little nutrition but are very high in calories. The text is divided into two sections, one for parents and one for children. The parent's guide provides tips on following the diet, giving reinforcement, learning nutrition, setting a good example, exercise, etc. The children's section describes the food classification system, provides tips on good eating, exercise, and following the rules in easy-to-read text. Weekly quizzes and charts appear in each section to assist parents and children. The appendix contains a weight-for-height table, the Stop-Light Diet food guide color classifications with serving size suggestions and calorie content, daily food chart, daily weight chart, basic four food group guidelines, menu ideas, exercise chart, and a diet contract to be signed by parents and children.

O197

Storage of medium-chain triglycerides in adipose tissue of orally fed infants. AJCNA. Sarda, P.; Lepage, G.-Roy, C.C.-Chessex, P. Bethesda, Md. American Society for Clinical Nutrition. American journal of clinical nutrition. Feb 1987, v. 45 (2), p. 399-405. ill.,

charts. Includes 34 references. (NAL Call No. DNAL FNC 389.8 J824).
Extract: The effect of the fatty acid content of the diet on that of adipose tissue was studied in 5 newborn infants studied prior to feeding and 30 infants fed ad libitum from birth with either human milk or a commercial formula as the sole nutrient. Significant positive linear correlations of dietary intake on adipose tissue content of fatty acids were found for both long- and medium-chain fatty acids (MCFAs). Infants stored up to 12% of MCFAs in their subcutaneous fat. The technique of direct transesterification improved the recovery of the volatile MCFAs and could explain the finding that medium-chain triglyceride storage in adipose tissue is more extensive than in previous reports. This study documents that MCFAs are not used solely as a source of energy; they can be reesterified or serve for chain elongation, before being deposited in fat stores. (author)

O198

Students learn to grow up, not out. Ganse, R.D. Denver, Colo. American School Food Service Association. School foodservice journal, May 1988, v. 42 (5), p. 44-45. ill. (NAL Call No. DNAL 389.8 SCH6).

Abstract: Overweight students at Hillendale Elementary School in Baltimore, Maryland, are getting an opportunity to learn about healthful lifestyles and how to control their weight. A program developed and taught by the school guidance counselor and nurse allows a group of overweight students to share their problems with each other and to discuss body image, fitness, diet, and their role in weight control. Students learn about healthy eating and exercise habits and how to prepare healthful foods. They are encouraged to follow a specific diet pattern, keep a food diary, and maintain their present weight without gaining more. In addition to developing the skills needed to live a weight-conscious lifestyle, many students have shown improved self-esteem as a result of participating in the program.

O199

Studies of the dietary habits, food consumption and nutrient intakes of adolescents and young adults. WRNDA. Bull., N.L. Basel; S. Karger. World review of nutrition and dietetics. Literature review, 1988, v. 57, p. 24-74. charts. Includes 133 references. (NAL Call No. DNAL 389.1 W892).
Abstract: It is the purpose of this review to bring together the findings of recent dietary studies among 10- to 25-year-olds and to identify any common features or dietary patterns which appear to be characteristic of young people. Tables referring to original studies are included on food, nutrient, and energy intakes for subjects in several different age groups. Factors affecting adolescent diets and recommendations for ways in which adolescents and young adults could be

encouraged toward dietary change are discussed

0200

Sweet Preference and body fatness. Neonatal data

Grinker, J.A. ; Gropman-Rubin, J. -Bose, K. New York : Alan R. Liss Nutrition and behavior, 1986. v. 3 (3) p. 197-209. ill., charts. Includes 49 references. (NAL Call No. DNAL OP141 A1N86)

Extract: An examination of studies relating sensory responsiveness (for sweet detection and recognition threshold, as well as preference) fails to reveal systematic differences in the response of obese compared to nonobese individuals. Also, studies report food consumption data that fail to show a direct linear relationship between sweet or sugar intake and body weight in normal-weight and obese individuals. Recent studies have, however, reported an enhanced fat-carbohydrate preference in the obese as compared with normal-weight individuals. The current study reports sucking responses to a variety of sucrose solutions (0.0612m-0.5m) by neonates from families with or without maternal obesity. No significant differences in any sucking parameters were seen in spite of increased birth weight and fatness in infants of obese mothers. These results from the "preobese" are in agreement with earlier data reporting no enhanced sweet preference in obese individuals. The possibility of different feeding strategies once infants receive solid foods by obese and normal-weight mothers was also examined. (author)

0201

Thin kids the proven, healthy, sensible program for children who want to lose weight ... /Mindy Cohen and Louis Abramson with Ruth Winter. --.

Cohen, Mindy. ; Abramson, Louis.-Winter, Ruth.; 1980-. New York : Beaufort Books, c1985. xvii, 206 p. : ill. ; 23 cm. (NAL Call No. DNAL RJ399.C6C64 1985).

Abstract: A healthy, sensible diet plan designed specifically for children and used successfully in the "Thin Kids" weight loss program is introduced. Practical guidance is provided for beginning a weight loss program on topics such as 1) placing the child in control of his/her own weight, 2) setting reasonable weight loss goals and keeping accurate detailed food and activity records, 3) recognizing common dieting "pitfalls" and targeting behaviors that need to be changed, and 4) differentiating among behavior, response, reinforcement and reward. The comprehensive and balanced diet plan considers all facets of a child's life, including 1) challenges at home and at school-class parties, peer pressure, after-school snacking, 2) temptation times-birthday parties, holidays, vacations, eating out and 3) food "pushers"-well intentional friends and relatives who equate food with love and affection. Complete menus and calorie values are provided for 10 weeks on the

weight loss program, along with recipes, food shopping suggestions, and meal preparation ideas. A complete exercise program that provides an easy-to-follow schedule of activities is presented, with illustrations. Relaxation techniques to help children cope with stress and perhaps respond better to a food program are outlined. A number of first-person success stories and practical advice on making weightloss a family project highlight the "Thin Kids" program's basic principles and components. (aje)

0202

A three-year study of obesity and its relationship to high blood pressure in adolescents.

Adeyanju, M. ; Creswell, W.H. -Stone, D.B.-Macrina, D.M. Kent, Ohio : American School Health Association Journal of school health, Mar 1987. v. 57 (3). p. 109-113. ill., charts. Includes 14 references. (NAL Call No. DNAL LB3401.J6)

Abstract: A 3-year behavioral and clinical study followed 356 cases of adolescents during their high school years to assess the relationship between obesity and high blood pressure. The results indicated that more girls than boys were overfat as seniors; while overweight trends between sexes and between blacks and whites were relatively constant, obesity increased among black girls; and positive correlations were found between percent ideal body weight, body mass index, skinfold thickness, and blood pressure in black and white females. Related findings also are discussed.

0203

Today's children are fatter not fitter say experts.

Smith, S.M. New York, N.Y. : Environmental Nutrition, Inc. Environmental nutrition, Apr 1989. v. 12 (4). p. 1, 6-7. Includes 6 references. (NAL Call No.: DNAL TX341.E5).

Abstract: The incidence of obesity in children now is greater than in past decades. Less structured exercise, snacking, more television watching, promotion of sugar-laden food, poor eating habits and psychological or family problems are contributing factors. The author believes that obese children should lose weight but dieting needs to be monitored by a qualified professional. Obesity prevention tips and weight loss programs for youths are included.

0204

Tracking of body composition variables.

PEDIAU. Harsha, D.W. ; Smoak, C.G.-Nicklas, T.A.-Webber, L.S.-Berenson, G.S. Elk Grove Village, Ill. : American Academy of Pediatrics. Pediatrics. Cardiovascular Risk Factors from Birth to 7 Years of Age: The Bogalusa Heart Study. Nov 1987. v. 80 (5,pt.2). p. 779-783. ill., charts. Includes 14 references. (NAL Call No.: DNAL FNC RJ1.P42).

Extract: Height, weight, and skinfold measurements were obtained on a cohort

of 447 children from birth (weight) or 6 months of age (height and skinfold) and monitored yearly thereafter until 7 years of age. At age 7 years, 250 remained for follow-up screening. A significant degree of tracking was found for all variables from age 1 to age 7 years. Height and weight tracked most strongly. (age 1- to 7-year correlations = .42 and .44 respectively,) whereas skinfold tracked somewhat lower (.28). Earlier levels of each anthropometric variable were the best predictor of later level of that parameter. Implications for early detection and treatment of growth abnormalities are discussed. (author)

0205

Update in maternal and infant nutrition
CHEJA, Johnston, E. M. Ottawa, Canadian Home Economics Association. Canadian Home Economics Journal. Literature review. Summer 1989. v. 39 (3). p. 113-120. Includes 98 references. (NAL Call No. DNAL 321.8 C162)
Abstract: Results from a number of recent studies in maternal and infant nutrition may influence a variety of our nutrition recommendations for pregnancy and infancy. The purpose of this review is to emphasize some of the new research that either confirms or questions established practice. Currently, controversial issues concerning maternal nutrition include weight gain for the obese pregnant woman, the use of vitamin and mineral supplements during pregnancy, and the effects of secondhand smoke on the fetus. Topics in infant nutrition which require clarification are the use of unmodified cow's milk, the appropriate level of fat for the infant, and the prevention of obesity. All health professionals should participate in continuing education activities to ensure that current recommendations are based on the best available information.

0206

The validity of self-reported weight loss and weight gain efforts in adolescents.
INDIDJ, Rosen, J. C.; Poplawski, D. New York, N.Y.: John Wiley & Sons. The International Journal of Eating Disorders. July 1987. v. 6 (4). p. 515-523. 111. Includes 17 references. (NAL Call No. DNAL RA784.A115).
Extract: There has been a proliferation of epidemiological survey studies of weight reducing and eating behaviors in adolescents; however, the validity of these self-report questionnaires has received little attention. The present study was designed to determine whether self-report measures of efforts to lose or gain weight and use of specific weight control methods are consistent with other measures. There were 98 high school volunteers who completed a questionnaire about weight change efforts. Parallel versions of the questionnaire were also returned by a parent and a friend or sibling. There were 165 high school subjects who completed the questionnaire and also

recorded food intake, exercise, and various weight control methods for 7 days. External raters agreed with subjects' reports that they were trying to lose weight and weight losers consumed much less food according to their eating records. Self-reported weight gainers consumed much more food than others, but agreement with external raters was lower. External raters agreed with subjects who reported skipping meals and exercising to lose weight and the subjects exhibited these behaviors more frequently in their eating records. However, there was low consistency between the self-report questionnaire and external measures of drastic weight control behaviors such as vomiting and fasting. With the exception of drastic weight control behaviors, the results of this study are generally positive for the validity of self-report questionnaires. (author)

0207

Vitamin E status of northern Canadian newborns: relation of vitamin E to blood lipids.
AJCNA, Godel, J. C. Baltimore, Md. American Society for Clinical Nutrition. American Journal of Clinical Nutrition. Aug 1989. v. 50 (2). p. 375-380. charts. Includes 22 references. (NAL Call No. DNAL FNC 389.8 J824).
Abstract: Vitamin E status was determined in two groups of Canadian newborns: a northern group, mainly aboriginal (Indian and Inuit), and a southern group, mainly nonnative. Serum vitamin E, cholesterol, and triglyceride levels were measured in cord blood and ratios of vitamin E to both cholesterol (E:chol) and cholesterol plus triglyceride (E:chol+TG) were calculated. For the combined groups the mean serum concentration of vitamin E (8.71 ± 2.45 micromole/L), cholesterol (1.77 ± 0.46 mmol/L), and triglyceride (0.65 ± 0.30 mmol/L) as well as the ratios E:chol ($5.00 \times 10^{-3} \pm 1.26 \times 10^{-3}$) and E:chol+TG ($3.60 \times 10^{-3} \pm 0.77 \times 10^{-3}$) were within normal limits. Significant north-south differences were found only in the mean triglyceride concentration, which was lower ($p = 0.03$), and E:chol+TG, which was higher ($p = 0.002$), in the northern than in the southern group. No differences attributable to differences in race were found. Only one infant, an Inuit in the northern group, was found to be deficient in vitamin E.

0208

Weight management of youth with Prader-Willi syndrome.
INDIDJ, Mullins, J. B.; Vogl-Maier, B. New York, N.Y.: John Wiley & Sons. The International Journal of Eating Disorders. May 1987. v. 6 (3). p. 419-425. Includes 13 references. (NAL Call No. DNAL RA784.A115).
Extract: Prader-Willi (PW) syndrome is a relatively rare disorder in which hypotonia, hypogonadism, obesity, and some degree of cognitive and behavioral difficulties are experienced. A special rehabilitation group program for 9

children with this syndrome was undertaken in a residential setting for 26 days each of 3 years to facilitate weight reduction, nutrition awareness, physical fitness, and social adaptability. Parent education was also undertaken. The nutrition education included a red, yellow, and green diet system. An exercise program and social program with a behavior modification system of rewards was established. Selected measures were taken before and after each program. Over a 3-year period, the 8 children who came for each of 3 summers maintained their weight control instead of vastly increasing their weight as untreated PW children do. It was therefore inferred that the children had gained nutrition awareness. The children improved to some extent on measures of physical fitness. Guidelines are suggested for school personnel and parents who are in contact with PW children. (author)

Q209

Weight reduction techniques in the pediatric patient

Lowenstein, M. K. Gaithersburg, Md. Aspen Publishers. Topics in clinical nutrition. Jan 1987. v. 2 (1) p. 49-54. Includes 4 references. (NAL Call No. DNAL RM214 T66)

Abstract. An authoritative overview by a clinical dietitian outlines practical methods of assessing and achieving weight loss goals with overweight pediatric patients, while establishing a psychological framework for assisting such patients in developing positive dietary habits. Specific attention is given to the evaluation of personal impressions of the pediatric patient and potential negative influences produced by the family environment, possible dispositions towards genetic and environmental factors, specific guidelines for approaching a positive child-dietitian interaction; and practical weight loss methods through the improvement of patient self-esteem, nutrition education, and behavior modification to change negative attitudes and habits. Individualized meal plans, weight goals, and tutorial role models, also are discussed. (wz)

Q210

Weights of British and French children.

Stark, O.; Peckham, C.S.; Aches, A. Boston, Mass. Little, Brown and Company. The Lancet. Apr 12, 1986. v. 1 (8485) p. 862. Includes 3 references. (NAL Call No. DNAL 448.8 L22).

Abstract. A brief correspondence user a diagrammatic display to compare the body mass index (BMI) centiles at ages 7, 11, and 16 years for a cohort of British children born in 1958 with those of French children born in 1953. The results indicate that an 11 year-old British boy of average height whose BMI is on the 50th, 90th, or 97th centile would be about 0.7, 1.6, or 5.1 kg heavier, respectively, than his French counterpart. The possible health implications of this for British

children is questioned. (wz)

Q211

The you can do it! kids diet /Dee Matthews with Allan Zullo and Bruce Nash. --

Matthews, Dee; Zullo, Allan; Nash, Bruce. New York: Holt, Rinehart & Winston, c1985. Includes index. xiii + 256 p. ill., 24 cm. "Bibliography by Patty Campbell" p. 241-245. (NAL Call No. DNAL JRM222 2 M37 1985)

Abstract. Information and guidelines on dieting for teenagers are presented in this easy-to-read book. Twenty-four chapters detail the author's profile about being fat, the problems of being overweight, preparing for dieting, the diet guidelines, a sample 14-day menu, recipes, good eating habits, coping with temptation at home and away from home, maintaining weight loss, and a 7-day sample maintenance menu. Exercises, graphs, suggested reading list, and recipes are included. (mp)

Q212

You're someBODY how to be a slim kid /by Lynda Corby and Patti Clark

Corby, Lynda; Clark, Patti.; 1951- Saskatoon: Fifth House, 1985. 116 p. ill. (some col.); 28 cm. Includes bibliographical references. (NAL Call No. DNAL JRU399.C6C6)

Abstract. A children's activity book and companion's guide are the major components of this weight management program for children. The book is designed for a parent-child partnership but can also be used by a variety of professionals, including teachers, dietitians, nutritionists, home economists, and public health nurses. Self-discovery and the development of self-esteem are the two basic principles incorporated throughout the book. Self-discovery begins with a variety of record-keeping methods that children use 1) to identify eating and exercise behaviors that contribute to overweight, and 2) to develop a plan for change. Children's activities are self-explanatory for the most part, additional details and explanations are provided where necessary in the companion's guide. Appendices in the companion's guide include 1) children's food experience books, 2) nutrition and fitness resources for parents and children, 3) activities for children with handicapping conditions, and 4) addresses of federal and local resource groups.

-Nichols, J -Magarey, A 97
 Abbott, W G H 23
 Abernethy, E 151
 Abramson, Louis -Winter, Ruth, 201
 ADBEDS 32
 Adevanjo M 174 202
 Adler, V 137
 Agras, W S 61
 AJCNA 1 2, 24 25 33 89 90, 94
 100, 101, 104 121 135, 138, 155
 164 172, 173, 178 180, 197, 207
 AJDCA 31, 116, 156
 Andrews, J S -Hood, R -Williams,
 P J -Bagatell, C J 140
 Arbeit, M L 28
 Archibald, E H -Pencharz, P.E 164
 Annett, J -Rallo J S -Srikameswaran,
 S -Vulcano, B 80
 Artavia-Loria, E -Henry, S -Basdevant
 A -Castano L 115
 Avons, P 77
 AvRuskin, Theodore W 49
 Balentine M B 58
 Balnar, B C -Heyden S 126
 Bandini, L G 14
 Banilivy M M -Lifshitz F 95
 Barker, D C P 118
 Barr, Ronald G -Leduc Denis
 G -Borsjoly, Christiane-Movey, -White
 Lynne-Pless, J Barry 53
 Beaudette, T ec 124
 Becque, M D -Marks, C -Moorehead
 C -Rocchini, A 155
 Bell, Louise 170
 Bellisle, F 138, 148
 Berntson, G 33
 Blair, D J 4
 Blessing, P 38
 BMJDAE 54, 154
 Boakes, Robert A -Popp'ewe' David
 A -Burton, Michael J 68
 Bodurtha J N 109
 Bonds, D R 6
 Bouchard, C 110 180
 Bougneres, P F 115
 Boulton, T J C 97
 Braddon, F E M 154
 Bray, G A 4, 180
 Brooke, G.G. 151
 Brown, M.R 79
 Bull, N.L 199
 Burke, G.L -Harsha, D.W -Srinivasan,
 S.R.-Cresanta, J L -Webber,
 L.S -Berenson, G S 175
 Buskirk, E R. 104
 Carroll, A 70
 Chan, Linda-Pencharz, Paul B 170
 CHEJA, 107, 205
 CIRCAZ, 169
 Clark, Patti, 195, 212
 Cohen, Mindy, 201
 Collipp, Platon J... 37
 Corby, Lynda, 195, 212
 CPEDA, 122
 Craig-Schmidt, M.C.-Waete, J.D.-Clark,
 A.J. 90
 Crawford, M.A. 96
 Creswell, W.H. Jr. 174
 Creswell, W.H.-Stone, D.B.-Macrina, D.M
 202
 Crosby, L.O. 6
 Cross, Alan W. 105
 DACDA, 3
 David, P.-Vobecky, J 56
 Davis, S.M.-Ford, V.L.-Tso, H 34
 Dennison, B.A. 165
 Desmond, S M -Ruppert, E S -Stelzer,
 C.M. 183
 Desmono, S M -Steizer, C M 73
 Deurenberg, P 24
 DIAEA 23 85 115
 DICAD2 152
 Dietz Jr 60
 Dietz W H Jr 14 31 42, 113, 121
 139
 Dietz W H Jr -Sobol, A M -Wentler, C A
 116
 Downey, A M 112
 Doyle W -Drury, P J -Meadows, N 96
 Drake, M A 186
 Dugdale, A E 119
 Durnin, J V G A 30
 Dweck, H S -Holmes F -Kretchmer
 R -Mauer, A M -Reynolds, J W -Suskind,
 R M -Benson, J D -Miguel, S G -Purvis,
 G A 171
 ECFNB 75, 92 193
 Edelstein, P -Resnef A -Halfon,
 S T -Paiti, H 188
 Elliot, D.L. 132
 Epstein, L H 88, 157, 178
 Epstein, Leonard H 71, 196
 Fairburn C 66
 Falkner, F 100
 Falkner, Frank 101
 Fannier S C -Rogers C S 41
 Farris, Rosanne P 141
 FDACB 102
 Feldman, E.-Goodman, J T 48
 Feldman, W 48
 Finberg, L. 166, 171
 Flodmark, C E -Fex G -Henningsen, N C
 11
 Fodor, J G 126
 Forbes, G.B 79
 Forester, D. 65
 Fox, R.A -Matson, J -Menta, S -Baker,
 A -Lopuch, W R 29
 Francis, Dorothy E M 142
 Frank, G C.-Smoak, C G -Freedman,
 D.S -Berenson, G S 50
 Frank, G C -Webber L S -Harsha,
 C W -Virgilio, S J -Franklin,
 F.A -Berenson, G S 112
 Frank, Gail C.-Webber, Larry
 S.-Berenson, Gerald S 141
 Frankle, Reva T. 145
 Freedman, D.S 158, 172, 173, 175, 190
 Freedman, D.S -Burke, G.L -Harsha,
 D.W.-Berenson, G.S 26
 Freedman, D.S.-Burke, G.L -Harsha,
 D.W.-Webber, L.S -Berenson, G.S. 185
 Fripp, Raymond R 8
 Fujino, M.-Shindo, M.-Hiroki,
 T.-Arakawa, K. 69
 Ganse, R D. 198
 Garcia-Diz, L.-Cuesca, C.-Sanchez-Muniz,
 F.J.-Castro, A. 176
 Garcia, R.E. 181
 Garn, S.M. 92, 93
 Gillum, R.F. 47
 Glueck, C.J. 169, 182
 Godel, J C. 207
 Goldberg, L.-Kuehl, K.S -Hanna, C 132
 Goldbourt, U.-Cohen-Mandelzweig,
 L.-Katz, M.-Appel, S.-Harel,
 G.-Sperling, Z.-Lazarovici, M.-Hart,
 J.-Neufeld, H.N. 59
 Goni, I. 176
 Gortmaker, S.L. 116
 Goulart, Frances Sheridan, 20
 Graver, E.J.-Vargas, M.-Churella,
 H.R.-Paule, C.L. 103
 Greene, H.L. 159
 Griffin, G.-Murray, D.M 13
 Griffiths, M. 78, 150

Grinker, J. A. 200
 Gropman-Rubin, J. -Bose, K. 200
 Guinn, B. 9
 Guinn, Bobb, 74
 Guc, S. 89
 Hackett, A. F. 57
 Hann, P. 147
 Hale, E. 102
 Hamosn, M. 62
 Hansson, C. L. -Eck, L. H. -Duff, A. C. 1
 Harris, M. E. 34
 Harrison, G. G. 103
 Harsna, C. W. 204
 Hartung, R. 31
 Hauner, H. 2
 Havashi, T. 69
 Hediger, M. L. -Zemel, E. S. -Parks, J. S. 22
 Hevmsfiele, S. E. 140
 Hoogson, James L. -Kwiterovich, Peter
 C. -Werner, John C. -Schuler, H.
 Gregg-Wrightman, Victor 8
 Hofman, A. -Vaughan, R. D. -Wynder, E. L.
 133
 Ho Z. Monika-Brubacher, G. 121
 Horibe, H. -Sawada, S. -Uyama,
 M. -Matsuzawa, Y. -Yamamura,
 T. -Yokoyama, S. -Kasagi, F. -Kameda, K.
 189
 Hunt-Peow, J. 192
 Huttner, M. P. 160
 IJCBOP 17, 41, 69, 82, 83, 86, 110
 129, 160
 INDIO, 80, 206, 208
 Iton, R. 15
 Ivanovic, D. 144, 177, 191
 Ivanovic, R. -Buttton, C. 144
 IZVIK, 179
 JADAA, 5, 9, 10, 14, 28, 63, 64, 79, 87,
 88, 131, 145, 170, 186, 192
 JAMAA, 123, 175
 James, W. P. 177
 JFSTA, 93
 Johnson, C. C. -Arbeit, M. L. -Franklin,
 F. A. -Berenson, G. S. 63
 Johnston, E. M. 205
 JOPOA, 36, 39, 49, 53, 61, 71, 105, 132,
 165, 166
 JPGND6, 11, 21, 62, 103
 JSHEA, 74, 141
 Kahle, E. B. 85
 Kaplan, K. M. 39
 Katch, V. 17, 155
 Katch, V. -Schork, A. -Kelch, R. P. 120
 Katz, S. H. 22
 Key, J. -Bondie, D. -Chico, R. -Moorehead,
 C. -Katch, V. -Martin, M. 72
 Kies, C. 187
 Kikuchi, D. A. -Srinivasan, S. R. -Webber,
 L. S. -Berenson, G. S. 165
 Klaveren, H. van-Deurenberg, P. 32
 Klesges, R. C. 1
 Knip, M. -Paavilainen, T. 160
 Knwler, W. C. -Bennett, P. H. -Aleck,
 K. A. -Baird, H. R. 152
 Knulman, J. T. 81
 Koral, A. 106
 Korsch, B. 36
 Kowalski, Robert E. 43
 Kraemer, H. C. -Berkowitz, R. I. -Korner,
 A. F. -Hammer, L. D. 61
 Kramer, Michael S. 53
 Kretchmer, Norman, 99
 Kumanyika, Shiriki K. 131
 Labadarios, D. 163
 LaRosa, J. 166
 Larson, S. -Ginsberg-Fellner, F. 82
 Laskarzewski, P. M. -Suchindran,
 C. M. -Chambiess, L. E. -Barrett-Connor,
 E. -Stewart, P. -Heiss, G. -Tyroler, H. A.
 169
 Lauer, R. M. -Clarke, W. R. 128
 LeBow, M. D. 16
 LeBow, Michael D. 107
 Lee, J. 128
 Leichner, P. 80
 Lepage, G. -Roy, C. C. -Chessex, P. 197
 Levy, Robert I. 168
 Liebman, B. 125
 Littrell, J. M. 51
 Loader, Peter J. 40
 Lowenstein, M. K. 209
 Lundholm, J. K. 51
 Mackova, E. -Mackova, J. -Skopkova, M. 21
 Malina, R. M. 27, 83
 Mallick, M. J. 18
 Malvy, J. M. D. 179
 Marks, J. S. -de Romana, G. L. -Madrid,
 S. -Boutton, T. W. -Klein, P. D. 25
 Matthews, Dee. 211
 Mazeis, A. G. 82
 McCarron, D. A. 55
 Menta, S. K. 156
 Mellies, M. J. -Dine, M. -Perry,
 T. -Laskarzewski, P. 182
 Mellin, L. M. 5
 Meyerson, D. A. 19
 Mollaner, T. 94
 Moodie, D. S. 181
 Morley-Kotchen, J. -Wynder, E. 162
 Morris, C. 55
 Moses, N. 95
 Mourey, M. S. -Carlier, C. -Caces,
 P. -Dostalova, L. -Montagnon,
 B. -Amedee-Manesme, O. 179
 Mullins, J. B. 208
 Mullis, R. M. -Maile, M. C. 134
 Nader, P. R. 7
 Nader, P. R. -Sallis, J. F. -Patterson,
 T. L. -Rupp, J. W. 64
 NEJMAG, 6, 72, 76, 133
 Nicklas, T. A. 63, 135
 Nicklas, T. A. -Frank, G. C. -Webber,
 L. S. -Miner, M. H. -Berenson, G. S. 28
 NUHEB, 96, 98, 130
 NUREB, 56, 108, 126, 127
 NUREA, 45, 84, 114, 117, 149
 NURHA, 46
 NURIB, 15, 144, 176, 177, 187, 191
 O'Dorisio, T. M. -Walker, R. B. -Eisenman,
 P. A. -Reiser, S. -Cataland, S. -Zipf,
 W. B. 85
 Oka, J. -Echizen, H. -Murakami, K. -Aeba,
 H. 15
 Orlandi, M. A. -Vaccaro, D. -Wynder, E. 111
 Osmond, C. 118
 Palumbo, F. M. 42
 Parcel, G. S. 184
 Parizkova, J. 21
 Patterson, R. E. 87
 Peckham, C. S. -Ades, A. 210
 PEDIAU, 7, 8, 48, 50, 60, 95, 109, 128,
 159, 162, 171, 181, 182, 190, 204
 Perry, C. L. 13, 134
 Perusse, L. -Rivest, J. -Roy,
 R. -Morissette, J. -Allard,
 C. -Theriault, G. -Leblanc, C. -Tremblay,
 A. 110
 Pettitt, D. J. 152
 Peveler, R. 66
 Phillips, B. L. -Franck, L. -Fillmore,
 C. M. -Said, H. M. -Murrell, J. E. -Moore,
 M. E. C. -Briggs, R. 159
 PHRPA, 47
 Pillai, Shashikala-Juan,
 Christina-Kleinberg, David L. 49
 Poh Tan, S. 75

Poplawski C 206
 Potter C 127
 Prattala R 46 193
 Price, J H 73, 183
 Rankonen C -Rimpela M 46
 Rasanen L -Viikari, J -Akerblom,
 H K -Anjala M -Unari, M -Pasanen,
 M -Nikka, T 94
 Resnick K 111 162
 Riutalo U 30
 Rivers, J F W -Hornville E A 150
 Rivers J F W -Payne P R 78
 Roberts S E 76
 Rocchini A -Beccue, D -Marks
 C -Moorehead, K 17
 Rocchini, A P 72 120
 Rome, A F -Houtkooper, L 89
 Rodgers B -Wadsworth, M E J -Davies,
 J M C 154
 Rogers, C S 86
 Rolland-Cachera, M F 138
 Rolland-Cachera, M F -Deneeger,
 M -Guilloud-Bataillie, M 148
 Rosen, J C 206
 Rotatori A F 29
 Rugg-Gunn A J -Appleton, D R -Coombs
 A E
 Saha N 108
 Saitoz M T 191
 Sallis J F -Patterson, T -Kupp
 J -Atkins C -Nader, P R 12
 Santanello, N C 19
 Sarda, P 197
 Sasaki, J 129
 Savage J -Coward, W A -Chew B -Lucas
 A 76
 Schaefer, Lynda U 131
 Schieker, R -Segrest, J -Nance, W E 109
 Semper, T -Crofts, A 9
 Shear, C L 26, 185
 Shear, C L -Burke, G L -Srinivasan
 S R -Webber, L S -Harsha,
 D W -Berenson, G S 158
 Shephard G S -Hesseling
 F B -Hutchison, M E 163
 Srinoc, M -Tanaka, H -Ando, M -Arakawa
 K 129
 Simons-Morton, B G -O'Hara,
 N M -Baranowski, T -Kolbe, L J -Bee,
 D E 184
 Slinkard, L A -Irwin, C E Jr 5
 Smith, S M 203
 Smoak C G -Nicklas, T A -Webber,
 L S -Berenson, G S 204
 Sobnany, M S 86
 Squires, Sally 196
 Srinivasan, S.R. -Burke, G L -Shear,
 C L -Smoak, C.G -Harsha, D W -Webber,
 L S -Berenson, G.S. 172
 Srinivasan, S R. -Cresanta, J.L -Webber,
 L.S. -Berenson, G.S. 190
 Srinivasan, S.R. -Harsha, D.W. -Webber,
 L.S. -Berenson, G.S. 173
 Stallings, V.A. 164
 Stark, O. 210
 Stickler, G.B 122
 Stitt, K. -Bonner, J -Clark, L. 58
 Story, M. 10
 Strobl, W. 136
 Sullivan, T V -Hawthorne, V.M. 92, 93
 Sumner, S.K. 153
 Sveger, T. 11
 Taitz, L.S. 54, 98
 Tamin, D. 188
 Taras, H.L. 64
 Taras, H.L. -Sallis, J.F. -Patterson, T.L.
 7
 Thuillez, P -Howard, B.V. -Bennett,

F H -Salans, J.E -Cushman,
 S W -Reaver, G M -Foley, J E 23
 Tompkins, R A -Bass, M A -Wakefield
 L M 10
 Trowbridge, F L 25
 Typpo J T -Tvppc M H -Krause G F 87
 Valoski A -Koeske, R -Wing R R 88
 Valoski A -Wing R R -Perkins
 K A -Fernstrom, M -Marks, E -McCurley,
 J 157
 Van der Reek, M M 90
 Vega W A 12
 Vobecky, J S 56
 Vogl-Maier, B 208
 Wabitsch, M -Zwiazler, K -Widhalm,
 K -Pfeiffer, E F. 2
 Wadden, T A 39
 Walter, H U 133
 Watkins, J E 161
 Webber, L S 50
 Webber, L S -Thompson, B -Berenson, G S
 135
 Wells, J E -Beaver, D W -Hornblow, A.R
 75
 West, C E 81
 Weststrate, J.A 24, 32
 Whipple T W -Huerta E 18
 Widhalm K 127, 136
 Widhalm, Kurt -Naito, Herbert K 52
 William H -Gortmaker, Steven L 60
 Wing, R R -Cluss, P -Fernstrom,
 M H. -Penner, J -Perkins,
 K A -Nudelman, S -Marks, B -Valoski
 A 178
 Wing Rena R. -Penner, Barbara -Kress
 Mary Jeanne 71
 Wolbers, M 35
 Wolfe, R R 121
 Wolfie, J.A 41
 Wong, C T 108
 WRNDA, 199
 Wynn, A 130
 Wynn M 130
 Yamamoto, A 129
 Zahavi, I 59
 Zavaleta, A N -Little, B B 27, 83
 Zipf, W B 33
 Zullo, Allan -Nash, Bruce 211
 1923- 99
 1930- 201
 1932- 37
 1937- 168
 1938- 142
 1949- 195, 212
 1951- 195, 212