

ED 341 547

SE 052 352

AUTHOR Baker, Claude D., Comp.; And Others
 TITLE Experiential Training in Florida and the Florida Keys. A Pretrip Training Manual.
 PUB DATE May 91
 NOTE 82p.; For field trip guidelines, see ED 327 394.
 PUB TYPE Guides - Non-Classroom Use (055) -- Guides - Classroom Use - Teaching Guides (For Teacher) (052)

EDRS PRICE MF01/PC04 Plus Postage.
 DESCRIPTORS Animals; Classification; *Ecology; Environmental Education; Estuaries; *Field Trips; Higher Education; Ichthyology; *Marine Biology; Plant Identification; Plants (Botany); *Resource Materials; Science Activities; Science Education; Secondary Education
 IDENTIFIERS Coral Reefs; Dichotomous Keys; *Florida

ABSTRACT

This document is a pretrip instruction manual that can be used by secondary school and college teachers who are planning trips to visit the tropical habitats in South Florida. The material is divided into two parts: (1) several fact sheets on the various habitats in South Florida; and (2) a number of species lists for various areas. Factsheets on the classification of marine environments, the zones of the seashore, estuaries, mangroves, seagrass meadows, salt marshes, and coral reefs are included. The species lists included algae, higher plants, sponges, worms, mollusks, bryozoans, arthropods, echinoderms, vertebrates, insects, and other invertebrates. The scientific name, common name, and a brief description are supplied for all species. Activities on the behavior and social life of fish, a dichotomous key for seashells, and a section that lists useful references for Florida field trips are included. (KR)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *

A PRETRIP TRAINING MANUAL FOR:

**EXPERIENTIAL TRAINING IN
FLORIDA and the FLORIDA KEYS**

PART OF:

**AN INTERACTIVE FIELD BIOLOGY
PROGRAM CONSISTING OF PRETRIP
INSTRUCTION, SEARCH IMAGE
TRAINING, FIELD EXERCISES, AND
OBSERVATIONS OF TROPICAL HABITATS
AND CORAL REEFS**

COMPILED BY:

**CLAUDE D. BAKER, PROFESSOR OF BIOLOGY
BILL J. FORSYTH, PROFESSOR OF BIOLOGY
CARL CHRISTENSON, PROFESSOR OF BIOLOGY
DAVID E. STAPP, STUDENT ASSISTANT**

**DIVISION OF NATURAL SCIENCES
INDIANA UNIVERSITY SOUTHEAST
NEW ALBANY, INDIANA 47150**

MAY, 1991

**COMPLETED AS PART OF A SPRING 1991 SABBATICAL IN
FLORIDA BY CLAUDE D. BAKER**

"PERMISSION TO REPRODUCE THIS
MATERIAL HAS BEEN GRANTED BY

Claude D. Baker

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC)."

U. S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

This document has been reproduced as
received from the person or organization
originating it.
 Minor changes have been made to improve
reproduction quality.

• Points of view or opinions stated in this docu-
ment do not necessarily represent official
OEERI position or policy.

ED341547

SE052352



Notes to the user of this material.

The instructional materials developed for the field course are part of an award winning presentation for instructional development. The course guidelines were rearranged and submitted previously to the ERIC database. Copyrighted figures and certain other materials have been removed from this document..

The resulting pretrip instruction manual, however, can be used by secondary schools and college teachers who are planning trips to visit the tropical habitats in South Florida. The material is divided into two parts: (1) several fact sheets on the various habitats in south Florida and (2) a number of species lists for the various areas that we personally have visited.

In our own pretrip classes prior to departure, we use search image slide training to teach our students the major floral and faunal elements of south Florida . We also have video tapes of many of the habitats for students to peruse. While in Florida, we use the included checklists to summarize the elements that we see. These checklists are in a computer database; so we can update them every year. We also have a pretrip examination covering the material on the fact sheets.

When we travel to Florida for our field class, we normally take an extensive field library with us. These field guides are extremely helpful and many students delight in identifying additional organisms. We have included a suggested Florida-oriented library in the back of the instruction manual. Since many of these books are published by local Floridean companies, they may be difficult to locate. We suggest that individuals wanting a current list of what is available should contact: Mickler's Floridiana, Inc., P.O. Box 1450 Oviedo, Florida 32765. This is a company which specializes in Florida material. They offer a healthy discount for educational institutions.

Claude D. Baker, Ph.D.
Professor of Biology
Indiana University SE
4201 Grantline Road
New Albany, IN 47150
812-941-2251

TABLE OF CONTENTS

<u>Topic</u>	<u>Page</u>
Classification of Marine Environments	2
Supralittoral Zone	4
Littoral Zone	5
Sandy Shores	6
Muddy Shores	7
Estuaries	8
Sublittoral Zone	10
Mangrove Ecosystems	13
Seagrass Meadows	16
Salt Marshes	19
Coral Reefs	22
Fish Observed by 1991 Class	25
Required Fish List	26
1991 Species List	29
Species Descriptions (Figures Removed)	30
Vegetation Observed by 1991 Field Biology Class	40
Cultivated, Exotic and Otherwise Interesting	41
Plant List for Field Biology Fresh Water Swamps and Wetlands	42
Plant List for Field Biology Coastal Vegetation--Dunes	44
Field Biology Animal List	45
Field Biology Animal List	46
Common Birds of the Florida Keys	47
Field Biology Animal List--Birds	48
Plant List for Field Biology-- Scrub Forests	49

Plant List for Field Biology-- Pine Flatwoods	51
Plant List for Field Biology-- Hammocks and Tree Islands	52
Plant List for Field Biology-- Cultivated Plants	54
Plant List for Field Biology-- Disturbed Habitats and Ecotones	55
Study of Fishes on Coral Reefs	57
Part 1. Swimming	
Locomotion	58
Part 2. Eating	59
Part 3. Color	60
Protective Color and Shape	61
Bottom Dwellers	62
Part 4. Social Life	63
Behavioral Observation Exercise	64
Seashell Key	66
Florida Field Trip References	Appended at End

L303--FACT SHEET

CLASSIFICATION OF MARINE ENVIRONMENTS

ECOLOGY is a complex subject. A proliferation of terms prompted a committee to develop a standardized classification of marine environment.

The **OCEAN** is divided into **TWO** main divisions:

- 1. PELAGIC 2. BENTHIC**

Pelagic--open water or whole body of oceanic water.

Benthic--bottom region.

PELAGIC REGION is further divided into:

- 1. NERITIC PROVINCE**
- 2. OCEANIC PROVINCE**

Neritic--Water that overlies the continental shelf.

Oceanic--Beyond the continental shelf in the open ocean.

PELAGIC REGION is, because of varying conditions with depth, divided into several layers or zones.

- 1. EPIPELAGIC--0-200 METERS**
- 2. MESOPELAGIC--200-1000 METERS**
- 3. BATHYPELAGIC--1000-4000 METERS**
- 4. ABYSSOPELAGIC--4000-6000 METERS**
- 5. HADOPELAGIC-->6000 METERS**

BENTHIC DIVISION is divided into the following zones.

- 1. SUPRALITTORAL SPRAY ZONE**--Above the high tide mark where the ocean is affected by salt spray from the ocean. Sometimes called the spray zone.
- 2. LITTORAL OR INTERTIDAL**--The littoral lies roughly between the mean high tide and the mean low tide. Since the tide is out roughly half the time, the organisms living here are adapted to varying degrees of stress and exposure. This is the best studied marine zone.
- 3. SUBLITTORAL**--A familiar zone because it corresponds to the Continental Shelf; Not well studied close to shore. From mean low tide out to a depth of 200 meters.
- 4. BATHYAL**--The sides of the ocean basins down to a depth of 4000 meters. Environment is variable with very deep being

characterized by oozes formed from the skeletons of microscopic organisms. The upper section is very much like the littoral.

5. ABYSSAL--Essentially the bottom of the ocean. Virtually 1/2 of the earth's surface is covered by this zone from 4000 to 6000 meters.

6. HADAL--Located in relatively narrow slits called **trenches**. Often associated with the island arcs and volcanic activity. >6000 meters.

L303--FACT SHEET

SUPRALITTORAL ZONE

This is the spray zone above the high tide.

ANIMALS OF THE SUPRALITTORAL

The common snail found on mangrove branches and high on the rocks is the angulate or mangrove periwinkle. These snails can withstand long periods of desiccation. The other snail found on the fringe is the beaded periwinkle. Another group of animals common here have hard exoskeletons to help prevent drying. These crustaceans include arthropods like land crabs, amphipods, and isopods.

Obvious in Florida is the ghost crab-*Ocypode*. Burrows can be seen all along the sandy beaches.

Others--terrestrial hermit crabs like *Coenobita*, land crabs like *Carcinosoma*, the isopod--sea roach, *Ligia*, the beachhopper-*Orchestia*, an amphipod. Isopods have abbreviated or direct development. All are still tied to the ocean for breeding.

PLANTS OF THE SUPRALITTORAL

Many plants are able to survive in the harsh environment of the supralittoral.

Beach and dune grasses--the dominant temperate species is American beach grass, *Ammophila breviligulata*. This grass requires burial by sand for continued growth; can be covered a meter or so. Highest coastal dunes in the world along the Great Lakes are stabilized by American beach grass.

The Florida and tropical counterpart is sea oats, *Uniola*. This plant is not as effective a trap as beach grass, but they do protect against erosion. Therefore, they're protected now because in the past they were used by the little old ladies for dry plant arrangements. Other Florida forms in this area include: sea purslane, seagrapes, sea oxeye, beach morning-glory, saltwort, Spanish bayonet, railroad vine, prickly pear cactus and many others. The conspicuous exotic species on the dunes is *Casuarina*, the Australian pine. Currently, attempts are being made to remove this shallow rooted tree.

The dune is the major protection against storm surge. Rooted plants help keep the dune in place. *No major sand dunes are present in the Keys!*

L303--FACT SHEET **THE SEASHORE--LITTORAL ZONE**

Rachel Carlson, in her book the Edge of the Sea, described the low tide line as "the primeval meeting place of the elements of earth and water, a place of compromise and conflict and eternal change." Understanding comes when we can sense the long rhythms of the earth and sea that sculpted its land forms and produced the rock and sand of which it is composed.

1. **Rocky, sandy, muddy shore types.** These are all subjected to alternate submersion and exposure by the tides.
2. Organisms are adapted to withstand exposure to the air. High tide--a water world; low tide--a terrestrial with extremes in temperature and moisture.

ROCKY INTERTIDAL

The rocky intertidal is an "action zone" between the tides which may feature a low diversity of organisms with **high numbers of selected species**. Physical factors include twice daily changes in pH, salinity, temperature, light, etc.

The uppermost layers near the high tide mark are exposed to the widest ranges in exposure to air, temperature fluctuations, intense solar radiation and desiccation. The lowest fringes may only be exposed briefly. This range results in the often noted zonation of organisms--the most striking feature of the rocky shore.

Characteristic features--1. Zonation--Stephenson's classical zonation. Read section from *Life Between Tidemarks*. 2. Few species generally found in great abundance. 3. Difficult living conditions--exposed on the rocks is the worst place (pounding of the waves), but more niches here--in cracks, crevices, holes, under the rocks, boring into rocks, and tidal pools.

Factors affecting zonation--Physical and Biotic.

Physical include temperature, force on waves, and most importantly exposure.

Temperature change is severe. Ocean--66-88 F; on the rocks 45-100 F.

Biotic factors--The usual factors like competition, predation, and grazing, but also involved are biochemical pathways like the enzyme systems.

L303--FACT SHEET--SANDY SHORES

Many ocean shores have sandy beaches subject to the full force of the ocean. Actually, they are part of a continuum of particle sizes ending with muddy shores. On the open sandy beach, physical forces and migrating substrate produce an inhospitable environment, but some organisms have been remarkably successful in colonizing the sand.

FEW PLANTS LIVE HERE; ALMOST ALL ANIMALS ARE BURROWERS.

SANDY BEACHES are strongly heterotrophic, meaning the organisms get their energy or food from decomposing organic material--detritus. This may be transported in as beach wrack during high tide or water may drain off the beach or be transported in from rivers.

THE FLORIDA KEYS HAVE FEW SANDY BEACHES. LITTORAL DRIFT OF SAND ENDS AT KEY BISCAYNE AND VIRGINIA KEY NEAR MIAMI, FLORIDA.

A typical Floridian beach profile might include: ghost crabs along the upper edge (most common tropical supralittoral beach form). Ghost crabs are good scavengers--actually increasing in areas of human activity. The mole crab, *Emerita*, and the coquina, *Donax*, are the common forms in the swash zone. Whiting, pompano, and sand drum follow waves in to feed on the worms, crabs and snails are found in the surf zone. Adaptations include smooth shells and reduced spines. Worm rock is found on the sublittoral fringe along Florida's east coast.

Things drifting into the shore include boards, plastic bottles, Portuguese man-of-war jellyfish, by-the-wind-sailor jellyfish, sargassum or gulfweed and little old ladies picking up shells.

L303--FACT SHEET--MUDDY SHORES

MUDDY SHORES, located in secluded areas away from the pounding of the ocean in enclosed bays, lagoons, harbors, and estuaries, have little slope and are thus appropriately referred to as mud flats. Little or no wave action yields to deposition of smaller particles. Lack of water movement beneath the surface leads to decomposition and oxygen uptake leading to anaerobic conditions. The smell of life is H₂S and FeS.

The mudflat surface sometimes looks barren with millions of holes. Scientists have suggested that production is transported from elsewhere. The problem is that the highly nutritious algal biomass is rapidly converted into animal flesh. The invertebrates are then preyed upon by a variety of birds, crabs, and fishes.

ADAPTATIONS OF ORGANISMS IN THE MUDFLAT

Burrowers and organisms with permanent tubes. Crabs have light bodies and long legs. Clams, to fight acidic conditions, have a thick organic periostracum. Physiological adaptations for low oxygen tension.

FEEDING BIOLOGY

Deposit Feeders--*Arenicola*--lug worm, *Capitella*, a polychaete, and *Macoma*--a bivalve. The lug worm has a permanent u-shaped burrow where it pulls mud in one end and deposits feces on the other. The polychaete moves through the mud. The bivalve uses its siphons like a vacuum cleaner.

Suspension Feeders--*Chaetopterus*, the parchment worm lives in a parchment-like tube sticking about 1/2 inch above the surface. It sucks water in and traps the goodies in a mucous ball.

Predators--Fishes when the tide is in; birds when the tide is out. Crabs are scavengers. Although there are a lot of plants, herbivores are few.

In Florida, extensive portions of the shallow Gulf Coast, Florida Bay, and the Intracoastal are covered with only a few inches of water at low tide. Other organisms here include the razor clam, *Tagelus*, the quahog, *Mercenaria*, young stone crabs, *Menippe*, common mud crabs, *Panopeus*, common mud snail, *Nassarius*, hermit crabs, and a number of polychaetes.

L303-FACT SHEET--ESTUARIES

ESTUARIES are coastal areas where mixing with fresh water occurs. In Florida, estuaries comprise a complex system including wetlands, grasses, mangroves, oyster bars, sand, mudflats, and algae or seagrass meadows covering millions of acres. They are an immensely valuable resource (nursery and spawning ground, buffer against storms, recreational area, a detritus food source for offshore), but unfortunately they are a target for development and the release of effluents.

Tidal flux produces changes in salinity. Therefore organisms have a problem with osmoregulation. Abundance of true estuarine species is consequently restricted to euryhaline fishes like sardines, anchovies, seatrout, redfish, mullet, snook, tarpon, menhaden and several others. The importance of estuarine habitats is emphasized by the fact that 97+ percent of total Gulf of Mexico fisheries catch is comprised of species that are dependant on estuaries during some phase of their life cycle. Most are spawned offshore migrating into the estuaries where their juveniles use a variety of habitats. Commercial pink shrimp grow up here too.

In northern climates, Spartina or cord grass is a major detritus contributor. In Florida, manatee grass, turtle grass, shoal grass and eel grass, Zostera, are major contributors. Florida Bay is dominated by innumerable mangrove islands and mangrove forest fronting extensive marshes on the mainland. Red, black, and white mangroves inject significant energy into the system.

Florida contains over 1,300 miles of marine coastline. Adjoining the Gulf of Mexico are over 3,000,000 acres of estuarine water (11 major systems) which may vary in size from a few acres to a few hundred square miles. Many estuaries here are classical in the sense that they have pronounced salinity gradients. On the east coast, the open ocean is characterized by a high energy shoreline with some 18 river mouths and inlets into the back-barrier system. The St. John's River is the major estuarine river system on the east coast. The remainder of the system is characterized by estuarine lagoons with poor circulation. See the attached figure for a list of Florida's estuaries.

Shell mounds and oyster middens left behind by the Indians testify to the historical dependence upon the resources of the estuaries. Now this has grown to encompass commercial and recreational elements. Dominant estuarine-related species include: mullet,

spotted, sea trout, snook, pink shrimp, red drum, blue crabs, and oysters. The center of the oyster fishery is Apalachicola Bay. Crabs are harvested from the "Big Bend" area of the northeast Gulf of Mexico, from southwest Florida, and northeast Florida to the Georgia border. Smaller commercial pink shrimp taken from estuaries are often harvested and sold as bait shrimp.

L303--FACT SHEET--NERITIC PROVINCE, BENTHIC DIVISION SUBLITTORAL ZONE

The ocean is the reason why we visit Florida. Coupled with the warm tropical flow of the Florida Current, the ocean is responsible for the moderate climate. Florida has over a thousand miles of coastline, and **60,000 square miles of continental shelf**. These neritic waters may be divided into nearshore waters heavily influenced by freshwater inflow and those waters further offshore affected by inshore coastal water and offshore oceanic water. Oceanic water off Florida in the area of the Keys and the southern portion of the peninsula is influenced by the Loop Current and the Florida Current. After going through the Florida Straits, the Florida Current maintains a close proximity to land until it reaches the St. Lucie River area where the land mass angles westward.

Demersal or benthic habitats are those associated with the bottom. **A greater variety of benthic habitats occurs in the marine waters off Florida than anywhere in the United States.** These include: sediment areas, seagrass meadows, live bottom, *Oculina* coral reefs, worm rock reefs, vermetid worm snail reefs, tropical coral reefs, and numerous artificial structures like ship wrecks. These sites, of course, provide habitat for a diversity of fishes and invertebrates. In addition, they protect shorelines, stabilize sediments, and contribute to water clarity. The habitats are interrelated because organisms using one habitat might utilize another as a feeding or resting area.

The continental shelf off the east coast is narrow (40 km at Cape Canaveral, 1 at Palm Beach, and 13 at Key West). The Florida Keys archipelago begins in southern Biscayne Bay and arcs southward and westward to the Dry Tortugas. The sediment-covered shelf off the west coast is wide (up to 240 km of drowned karst topography). The bottom is irregular being marred by solution basins and sinkholes. An exception is the Florida Middle Ground, an area of great relief with pinnacles reaching up to 13 meters above the bottom.

Most of east Florida's shelf is covered with sand down to Biscayne Bay. In the Keys, the sediment is mostly carbonate sand from mollusks, coralline algae, etc. Off west Florida, the world famous "barking sands" of high quartz content were probably deposited during a period of lower sea level. Sand along the east coast is derived from land sources and carried southward by longshore currents. Florida Bay is dominated by calcareous mud and slits, much of it skeletal remains of bivalve mollusks and foraminiferans.

Significant food is available on the continental shelf. Worldwide, all big fisheries are found in this area. Species diversity is particularly high in Florida. Florida's lengthy shoreline and coastal habitats provide habitat for **over 1000 species of marine and estuarine fishes--one quarter of the entire number recorded for the northern portion of the Western Hemisphere!!!** This great diversity is partly due to Florida's location in a climatic transition zone. The waters contain both temperate fishes (and invertebrates) from the Carolinian Marine Province and tropical elements from the Caribbean West Indian Province.

The best Known and most profitable benthic or demersal fishery in Florida is the commercial pink shrimp industry. Other species taken include scallops, clams, conchs, stone crabs, flounders, croakers, tilefish, pompano, permit, bonefish, saltwater catfish, stingrays, and pinfish.

Live Bottom--This is a term used to describe bottoms where gorgonians, sponges, hydroids, anemones, corals and other organisms accumulate. This type of habitat provides excellent cover for fishes. Therefore, these are the major areas where we fish and SCUBA. Valuable food fishes found here include: sea basses, groupers, snappers, porgies, grunts, amberjacks, and triggerfish. Commercial sponges occur here also.

Oculina coral reefs--Unusual reefs or living deep water corals occur off the eastern coast at depths of 70-100 meters. These reefs consist entirely of bushy thickets of a single species--the ivory tree coral, *Oculina*. The coral supports a diverse and dense association of invertebrates which in turn support a dependant group of commercially valuable fishes.

Worm rock reefs--Sabellariid worms form large colonies cemented together with sand grains from the intertidal area to about 10 meters. A steady supply of sand is required for this activity; so you won't see these structures past Biscayne Bay. Spiny lobsters, snappers, groupers and a variety of fishes seek the protection of these interesting structures.

Vermetid worm snail reefs--Wormlike shell masses of the snail, *Petalocochus*. These reefs located on the lower west coast are now dead, but they were living in the 1890's. The reason for their death is unclear.

Mangrove rock reef--The only known example of fossilized mangroves is found near Bear Cut on Key Biscayne.

Tropical coral reefs--Communities found off southeast Florida are the only shallow water coral reefs found in North American waters. These are covered in detail in another section.

Artificial habitats--Bridges, piers, wrecks, pipelines, etc.

Fishery resources--Oceanic Pelagic. Important fishery species off Florida include the billfishes (sailfish, white and blue marlin, and swordfish), oceanic tunas (bluefin, yellowfin, bigeye, and skipjack tunas), dolphin fish, and sharks.

Fishery resources--Oceanic Neritic. Important species in the neritic environment include: the mackerels (king mackerel and Spanish mackerel), cobia, little tunny, and bluefish. Other important species include neritic clupeids such as Spanish sardines and thread herring, jacks, blackfin tuna, and bailyhoo. Mulletts and menhadens spawn in the neritic, but they spend much of their life in the estuaries.

L303--FACT SHEET--MANGROVE ECOSYSTEMS

The word-mangrove-has a complex etymology. It may have been derived from the Portuguese word for tree, mangue, and the English word for trees, grove. Other possible origins include the Malay--manggi or Taino--mangle. The word mangrove refers to a phylogenetically separate group of tropical salt tolerant trees. As a group, they are the **most numerous trees in the world**. In south Florida, mangroves serve two distinct roles: They serve as a habitat for juvenile fishes and they provide an enormous supply of leaf material (detritus) which is a source of energy for a detritus-based food web supporting numerous invertebrate species or small forage fish species that are food for other fish.

Four mangrove species are recognized in Florida: red mangrove (*Rhizophora mangle*), black mangrove (*Avicennia germinans*), white mangrove (*Laguncularia racemosa*), and buttonwood (*Conocarpus erecta*). Buttonwood is most often associated with the first three species at the transition zone to upland areas, and frequently contributes to the overall biomass but not in quantities comparable to other species.

Because they are phylogenetically unrelated, mangroves have solved the problem of salt tolerance in differing ways. They are subjected to a continual buffeting by the waves, and a tropical sun that beats down seemingly 100% of the time.

Red mangroves--These are generally the most seaward of the three species. They are easily recognized by their strong arching prop roots. Try to pull one of these suckers out of the mud, and you'll appreciate why they are a first line of defense against storms. The velocity of the waves really slows down as they go through the mangroves. As in the mudflats, the oxygen environment is low around the mangroves. Look for dots called lenticels on the roots which permit gaseous exchange, but which close when the water comes in. Some salt enters which is collected in leaves which are continually sacrificed to the environment. Look for orange leaves that are ready to be dropped. In emergency situations, the leaves can be used as a source of tobacco and/or tea. Normal leaves are bright green and waxy in appearance (to prevent desiccation). Reproduction of the red mangrove is in the Spring with the seed or propagule germinating directly on the trees (a characteristic of true mangroves). The pencil-shaped propagules reach a length of up to 18 inches, hanging conspicuously from the branches in late summer. The seeds float parallel out of the system and then float around

upright for a year. The seed is actually a germinated plant ready to take hold in the appropriate location.

Black mangroves--The black mangrove usually grows landward of the red. The blacks seem to be the hardiest of the species being somewhat resistant to cold and very resistant to salt. The black mangrove has a network of horizontal cable roots which give rise to characteristic finger-like pneumatophores arising from the soil around the plant. The pneumatophores allow for direct absorption of oxygen into the roots. The black mangrove apparently needs fresh water for photosynthesis; therefore, the salt is excreted onto the side of the leaf prior to the water entering the photosynthetic system. The leaves tend to be more pointed and gray. Rub the back of a leaf and taste the salt that's been excreted. Fruits of the black mangroves are lima bean-shaped--about an inch long. Bees seem to like the flowers; therefore, the species contributes to a great deal of honey production.

White mangroves--This plant has rounded, yellow green leaves with salt glands on the petiole. White mangrove seeds look something like dried raisins.

Buttonwood--These mangroves have pointed leaves with salt secreting cells all down the midrib. General characteristics are a rough bark and buttonlike seeds. The tree, in the past, was regarded as a source of charcoal. An ornamental variety--the silver buttonwood--is used as a hedge or yard plant in the Keys.

Estimates of total acreage occupied by mangrove communities in Florida vary widely between 430,000 to over 650,000 acres.

These systems provide a significant input of energy into the ecosystem. The leaves eventually are broken down releasing tannic acid around the roots which may protect the system from shipworms and gribbles. It is the detritus or fragments of leaves, stems, twigs, roots, fruits, and seeds and the associated bacteria and fungi that forms the basis of the energy flow in the mangrove food webs.

First, the detritus is fragmented, consumed, and excreted by a number of consumers beginning with small crustaceans such as amphipods, caridean shrimps, xanthid crabs leading through a relatively simple food web which includes small fish and then to secondary consumers like gray snapper, tarpon, snook, red drum and spotted sea trout. See the attached figure for a general breakdown.

Mangroves in Florida are extremely vulnerable to man-induced perturbations. They are a protected habitat requiring both state and federal permits for excavation or filling in the vicinity of the trees.

L303--FACT SHEET--SEAGRASS MEADOWS

Florida has an estimated 800,000 acres of seagrasses. The dominant species are turtle grass-*Thalassia testudinum*, manatee grass-*Syringodium filiforme*, and shoal grass-*Halodule wrightii*. The majority of these grasses are found in the Keys in Florida Bay down into the Dry Tortugas. The seagrasses are land type plants in that they have both flowers and seeds; however, they are pollinated underwater. Turtle grass can't grow on hard rock; so on the reef you'll find it in back areas or in potholes where sediment has accumulated.

Seagrasses are tremendous producers of carbon. The epiphytic algae attached to the seagrasses also contribute significantly to the energy flow within the system. In addition to providing a source of food, the seagrasses by a process of "winnowing up" stabilize the sediments in a particular area thereby providing a habitat for benthic fauna.

Seagrass meadows control or modify the ecosystem in the following ways. First, they provide food for a very limited number of organisms such as parrotfishes, surgeonfishes, queen conchs, sea urchins and some nudibranchs. Green sea turtles originally utilized the resource leading to the general use of the name. Second, the grasses serve as hosts for epiphytes which are used or skimmed off by mullet and snails. Next, large and significant quantities of detritus from dead grasses are the beginning of the food chain. The binding of the sediments listed above preserves the microbial flora and prevents erosion.

The limiting factor for seagrass growth is available sunlight. They don't grow well in areas where there is a lot of turbidity and siltation. Normally, you don't find these grasses below 30 feet. Also they are very susceptible to the action of storms. After a storm, you sometimes find 3 or 4 feet of beach wrack piled up on the shore. Herbivores also detach the blades. The turtle grass blades are straplike and the bite of a parrotfish or urchin will only remove a portion of the blade. However, a bite on the round manatee grass always will sever the blade, and the portion above the bite will float to the surface and drift away. This is why a major portion of the beach wrack is often manatee grass. Turtle grass doesn't float that well and sort of moves along the bottom. Turtle grass blades have been photographed in very deep water--3900 meters.

Major vertebrate consumers of the seagrasses include turtles and manatees. Neither is abundant these days, but occasionally one observes turtles mowing down large tracts of grass. The small grassbed parrotfish is *Spariosoma radians*. You might find all the types of sea urchins in the grassbeds. Sea urchins graze upon the outer region of the leaf which has ceased growing, but which is often heavily colonized by epiphytes of high food value.

A great many other animals utilize the area. Some 113 epiphytic forms counted included several mollusks. The mollusk counts have been suggested to approach 95 billion per square mile. The grasses also provide physical protection for small resident species like seahorses and transient juvenile species of sport fishes like snook and tarpon. Some have hypothesized that the abundance of the commercial pink shrimp is directly related to areal extent of the sea grasses. The food for these fishes and shrimp comes from a complex community of microalgae, amphipods, rotifers, copepods, and other small animals associated with decaying grasses and their decomposers.

Survival in the grass flat in the Florida Keys is especially difficult. In contrast to the coral reef, there is really no place to hide. A great deal of effort is put into not being observed initially. Most organisms are small, drab, or cryptically colored. The sea horse is a prime example of a grass flat organism. Ordinarily, they don't move very quickly. The eyes even move independently of one another. Other fishes are able to avoid detection by using the grass meadows only at night. In Florida, vast schools of snapper and grunts seen in the reefs during the day are apparently only resting. They migrate sometimes over long distances during the night into grass flats to feed. Migration routes to the beds may remain constant for several years.

Large, armor-protected queen conchs, *Strombus gigas*, eat the turtle grass. Because of overcollection of the shells and meat, they have a collection restriction until at least 1989. The horse conch, a large predatory tulip shell, and the Florida state shell, preys on the queen conch. The horse conch is the largest snail in the United States. The smaller lightning whelk, tulip shells, and apple murex are other examples of predatory gastropods in the grass beds. The lightning whelk is appropriately named because it can literally run down its prey. The apple murex anesthetizes its prey. Often when captured, they smell like rotten cabbage producing a yellow color which turns lavender when dried. I've been told that the Romans used the dye to color the royal robes.

Also, look for sea cucumbers in the grass flats. In Florida, it is estimated that there are 90 tons of sea cucumbers for every 2 square miles. This is an animal you shouldn't collect. The combination of toxins and evisceration will kill the remainder of the collection. *Carapus*--the pearl fish--lives in the anus of the five-toothed sea cucumber. The names come from the fact that the fish originally was found inside a pearl (and if you believe this, you'll believe anything).

L303--FACT SHEET--SALT MARSHES

Salt marshes are intertidal herbaceous plant communities that develop along margins of estuaries where wave energy is low. Essentially, they are a specialized type of wetland.

Florida has three physiographic types of marsh wetlands. 1. Carolinian Marine Province--(Atlantic Coast north of Cape Canaveral). This area is characterized by a temperate biota with some seasonal tropical elements. 2. West Indian--Cape Canaveral south through the Keys and north along the Gulf Coast to Cedar Key. This area has a tropical biota with a few temperate organisms. Coral reefs and mangrove are characteristic features of the Keys. 3. Louisiana-- Gulf Coast north of Cedar Key. Organisms are similar to the Carolinian.

Some 80 to 90% of the Atlantic and Gulf Coasts consists of estuaries and lagoons with shoreline marshes. South Atlantic and Gulf coast marshes represent the largest coastal area in the United States. In Florida, mangroves dominate the shoreline areas in the southern part of the state. Grasses, sedges, and rushes characteristic of marshes act as the pioneer community in the mangrove zone, eventually being shaded out by the mangroves. Along the southeast tip of the peninsula, however, large stands of black needle rush, Juncus roemerianus, are associated with the mangroves.

Vegetation in these marshes consists of grasses (Poaceae), sedges (Cyperaceae) and rushes (Juncaceae). The areas are typically intertidal with the same diurnal changes found on the rocky shoreline. However, with their high rates of primary production, marshes are among the most productive natural ecosystems in the world. As with mangrove systems, marsh vegetation fragments (detritus) are an important beginning of estuarine and marine food webs.

The marshes provide habitat and protection for juvenile fishes, birds, and invertebrates. They protect coastal areas from erosion. Briefly, the salt marsh functions in detrital export, nutrient export, sediment trapping, pollutant removal, and the critical habitat for juvenile fishes mentioned above.

Although it is unlikely that we will visit the Atlantic or Gulf Coast marshes, you should be aware of the following: Atlantic marshes are dominated by smooth cord grass, Spartina alterniflora. Gulf

Coast marshes are dominated by black needle rush, *Juncus roemerianus*. Only a small portion of the vegetative biomass is converted into herbivore biomass in either area. Aquatic marsh invertebrate detritivores are at the bottom of the dominant food chain.

Juvenile penaeid commercial shrimp are an important marsh component in late spring and summer months. Blue crabs spawn in marshes and the larvae return to the marshes for growth into juveniles. American oyster (the kind we eat) builds intertidal and subtidal reefs in marsh areas. Oysters require a mixture of salt water from the tides and freshwater from land drainage where they filter water taking in phytoplankton, detritus, bacteria, and other particulate matter. A positive relationship between marsh area and oyster production has been established. It is now clear that healthy oysters depend on a viable salt marsh environment.

A substantial portion of the salt marshes of Florida are now under governmental control and protection. But as the population of Florida continually expands, demands for residential and commercial developments near estuary and marsh habitat will be a certainty. It has been estimated that 8% or 60,000 acres of basic estuary habitat has been lost just to dredge and fill operations. Channelization and maintenance of navigation channels also represent a significant alteration in urbanized areas. Many other perturbations including diversion of water, construction of dams, ranchings and farming alter the coastal zone habitats. Hopefully, integrated water basin management plans will include salt marshes and estuaries.

Table 1. Comparison of South Atlantic and Gulf Coast Marshes.

<u>Item</u>	<u>South Atlantic</u>	<u>Gulf Coast</u>
Plant Angiosperms		
Dominant plant	<u>Spartina alterniflora</u>	<u>Juncus roemerianus</u>
High Marsh plant	<u>Salicornia-</u>	<u>S. patens-Distichlis-</u>
Inland Marsh plant	<u>Distichlis-Juncus</u> <u>S. cynosuroides-</u> <u>Typha</u>	<u>Scirpus association</u> <u>S. cynosuroides-</u> <u>P. communis-Cladium</u>
Fauna		
Dominant insect		<u>Orchelimum</u>
Benthic bivalves	<u>Orchelimum</u> <u>fidicinium</u> <u>Prokelisia marginata</u>	<u>Conocephalus spp.</u>
Snails	<u>Modiolus demissus</u>	<u>Polymesoda caroliniana</u> <u>Modiolus demissus</u> <u>Melampus bidentatus</u>
Crabs	<u>Littorina irrorata</u>	<u>Littorina irrorata</u>
Mammals	<u>Uca spp. and Sesarma</u>	<u>Uca spp. and Sesarma</u>
Salinity Range	<u>Procyon lotor</u>	<u>Ondatra zibethica</u> <u>Myocastor coypus</u>
Tide Cycle (Amplitude)	10-30‰	2-15‰ Diurnal (0.3 m)
Freshwater Input	Semi-diurnal (2.0 m)	
Major River System		Mississippi River
Annual Precipitation	Savannah River	40 cm
Primary Productivity	120 cm	1500-3000 g/m ² /yr
Secondary Productivity	1000-2000 g/m ² /yr	
Penaeid Shrimp yield	5,964 tonnes (13.2 kg/ha)	55,193 tonnes (33.9 kg/ha)
Total fish yield	144,245 tonnes (320.2 kg/ha)	790,625 tonnes (485.4 kg/ha)
Size of intertidal areas	450,500 ha	1,628,900 ha

L303--FACT SHEET--Coral Reefs

Coral reefs are shallow water, tropical marine ecosystems of biological origin characterized by a tremendous variety of plants and animals and high rates of production in nutrient-poor and plankton-impoverished tropical ocean water. **Tropical coral reef communities found off southeast Florida are the only shallow-water coral reefs found in North American waters.** They are similar to Caribbean coral reefs. Their existence this far north is a result of the moderating influence of the Florida Current and the availability of substrate for coral development.

The Keys and the remainder of the Atlantic were very much affected by the Pleistocene glaciation when the water cooled and the was lowered by maybe as much as 600 feet. These factors drastically reduced the number of coral species in the Atlantic. Currently, only 50 species of hermatypic (reef building) corals thrive in the Atlantic. The Pacific has some 80 genera and 700 species.

Corals are comprised of thousands of minute organisms called coral polyps (Cnidarian Class Anthozoa, Order Madreporaria or Scleractinia). In the hermatypic or hard corals, these tiny animals remove calcium carbonate from the water and form the reef. The other active components of the reef in the Keys are the fire corals (especially *Millepora complanata*) and some calcareous red algae of the coralline group.

Coral reefs are essentially a three-dimensional limestone framework which creates numerous niches for a diverse sedentary and mobile fauna. **Biological activity here has been described as being similar to a tropical rain forest with high species interactions like symbiosis, high gross primary productivity, and very rapid recycling of limiting factor nutrients like nitrogen and phosphorus. Most primary production is situated within the symbiotic algae with the coral tissue (zooxanthellae).**

In Florida, two major reef types are recognized, **inshore patch reefs and offshore bank reefs.** Patch reefs within sedimentary and seagrass communities are usually constructed by star and brain corals (*Monastraea annularis*, *Siderastrea siderea*, and *Diploria*). As noted below, the backbone of the Keys was an historic patch reef which over time built up to a thickness of 300 feet in some areas. Because this substrate is at present topographically low and porous, there is hardly any freshwater in the Keys. The

water is piped in from the north. The exception to this is Big Pine Key where a lens of Miami oolite overlies the Key Largo limestone. Patch reefs are normally irregularly shaped without a classic zonation pattern noted so often in the literature. Gorgonians like seawhips, plumes, feathers, and fans are often a conspicuous element in the patch reefs.

Bank reefs are normally 5-7 miles offshore and linearly parallel the Keys. Some like Looe Key possess well-defined biological zonation patterns controlled by depth, light, and wave force. Looe Key also is one of two reefs in the Keys with a spur and groove or finger and groove system with sand channels in between the coral formations. The shallowest part of the system is the reef flat which bears the brunt of the waves surging up through the finger and groove structures at Looe Key. Dead coral rubble, fire coral (do not touch), and some turtle grass may be found here. Near the top of the shallow portion of the spur and groove system, note the fire corals and mats of zoanthid or Cheerio coral. The spur and groove consists of a series of alternating ridges and channels. In this area, the ridges have a lot of elkhorn coral (*Acropora palmata*). Grooves are filled with sand and coral rubble. Look for mountainous star coral-*Montastraea annularis*-the major reef builder in the buttress zone in about 15 feet of water.

How can corals live in areas that are relatively low in nutrients? First, a relatively narrow set of environmental conditions must be met.

1. A hard surface or a previous coral reef for settlement and attachment of larvae to initiate development of the reef. Some corals in the Keys like rose coral and finger coral tolerate some turbidity and live in sandy areas, but they do not form reef structures.
2. Massive reef builders are only found in the tropics where the temperatures average 23-25° C. The reef distribution is shifted upward into Florida from the tropics because of the above noted ction of the Florida Current.
3. Full-strength seawater with a salinity of 32 to 42 parts per thousand. They have little or no tolerance to low salinities, nor do they like turbidity.
4. Major reef building occurs in shallow water from 5 to 150 feet. They require light for the development of the zooxanthellae.

Reefs are most often conspicuously arranged roughly parallel to the shore with the seaward edge in the direction of the prevailing winds demonstrating the most active growth. In Florida and particularly the Bahamas, the bank or platform type of fringing reef is often characterized by the spur and groove system mentioned above. The flow of water through these grooves is important in channeling water onto the reef and in the transport of nutrients and organic matter. In general, the Gulf of Mexico is a poor place for reef development. Little bedrock is available; most of the Gulf rocks off Florida's Gulf Coast is covered with a sediment veneer. Notable exceptions are the Middle Grounds mentioned earlier and the Flower Gardens way out in the Gulf.

5. Zooxanthellae (endozoic dinoflagellate algae symbionts) living in the coral tissue. The highly modified dinoflagellates are in some way associated with the rapid growth of corals. The corals may feed on the photosynthetic products of the algae; the algae may play a significant role in calcium precipitation.

Coral Biology Facts-(1) Growth rate-young colonies grow faster than older ones; branched faster than massive (1 foot every 28 years in mountainous star coral--a massive form; staghorn coral can grow as fast as 4 inches a year). (2) Corals reproduce sexually producing a free swimming planular and Edwardsia larvae and asexually by budding. In extratentacular budding, each polyp forms its own cup. Brain corals bud along the entire length of the polyp. (3) Predators include snails and nudibranchs, polychaete worms, a few barnacles, crabs and major coral crushers like the parrotfish. The major and best known coral predator lives in the Pacific--the crown-of-thorns starfish. (4) Feeding is at night. Polyps extend their tentacles at night securing zooplankton as food for themselves and their symbiotic zooxanthellae. During day, the algae photosynthesize thus becoming the major source of primary production on the reef. (5) Little phytoplankton is around, but you should look for several types of benthic algae like Halimeda and Penicillus which are also carbonate secreting. (6) Although nutrients in the water are consistently low, those in the reef system are tenaciously retained conservative, regenerative, and recycling mechanisms. (7) Secondary production comes from autochthonous production inside the system rather than materials transported in. Read the seagrass section for an exception to the statement. Grunts resting over the elkhorn corals positively stimulate the coral growth with their release of waste products. (8) Diversity is high because of the wide variety of habitats generated by the coral structures.

Fish Observed by the 1991 Field Biology Class

- | | | | |
|---|--|--|--|
| <input checked="" type="checkbox"/> Ocean surgeon | <input checked="" type="checkbox"/> Doctorfish | <input checked="" type="checkbox"/> Blue tang | <input type="checkbox"/> Barred cardinalfish |
| <input type="checkbox"/> Flamefish | <input type="checkbox"/> Belted cardinalfish | <input checked="" type="checkbox"/> Trumpetfish | <input checked="" type="checkbox"/> Orangespotted filefish |
| <input checked="" type="checkbox"/> Scrawled filefish | <input checked="" type="checkbox"/> Slender filefish | <input checked="" type="checkbox"/> Bar jack | <input checked="" type="checkbox"/> Queen angelfish |
| <input checked="" type="checkbox"/> Gray angelfish | <input checked="" type="checkbox"/> French angelfish | <input checked="" type="checkbox"/> Rock beauty | <input checked="" type="checkbox"/> Reef butterfly |
| <input checked="" type="checkbox"/> Four-eye butterfly | <input checked="" type="checkbox"/> Spotfin butterfly | <input checked="" type="checkbox"/> Saddled blenny | <input type="checkbox"/> Roughhead blenny |
| <input type="checkbox"/> Wrasse blenny | <input checked="" type="checkbox"/> Redlip blenny | <input checked="" type="checkbox"/> Neon goby | <input type="checkbox"/> Bridled goby |
| <input checked="" type="checkbox"/> Masked goby | <input type="checkbox"/> Goldspot goby | <input type="checkbox"/> Hovering goby | <input checked="" type="checkbox"/> Herrings |
| <input checked="" type="checkbox"/> Bermuda chub | <input checked="" type="checkbox"/> Spanish hogfish | <input checked="" type="checkbox"/> Slippery dick | <input checked="" type="checkbox"/> Yellowhead wrasse |
| <input checked="" type="checkbox"/> Creole wrasse | <input checked="" type="checkbox"/> Hogfish | <input type="checkbox"/> Clown wrasse | <input type="checkbox"/> Blackear wrasse |
| <input checked="" type="checkbox"/> Puddingwife | <input checked="" type="checkbox"/> Bluehead wrasse | <input checked="" type="checkbox"/> Schoolmaster snapper | <input checked="" type="checkbox"/> Mutton snapper |
| <input checked="" type="checkbox"/> Gray snapper | <input checked="" type="checkbox"/> Mahogany snapper | <input checked="" type="checkbox"/> Yellowtail snapper | <input checked="" type="checkbox"/> Yellow goatfish |
| <input checked="" type="checkbox"/> Spotted goatfish | <input checked="" type="checkbox"/> Glassy sweeper | <input checked="" type="checkbox"/> Sergeant major | <input checked="" type="checkbox"/> Blue chromis |
| <input checked="" type="checkbox"/> Brown chromis | <input checked="" type="checkbox"/> Yellowtail damsel | <input checked="" type="checkbox"/> Dusky damselfish | <input checked="" type="checkbox"/> Beaugregory |
| <input checked="" type="checkbox"/> Bicolor damselfish | <input checked="" type="checkbox"/> Threespot damsel | <input checked="" type="checkbox"/> Cocoa damselfish | <input checked="" type="checkbox"/> Black margate |
| <input checked="" type="checkbox"/> Porkfish | <input checked="" type="checkbox"/> Spanish grunt | <input checked="" type="checkbox"/> Tomtate | <input checked="" type="checkbox"/> Caesar grunt |
| <input checked="" type="checkbox"/> Smallmouth grunt | <input checked="" type="checkbox"/> Blue parrotfish | <input checked="" type="checkbox"/> White grunt | <input checked="" type="checkbox"/> Bluestriped grunt |
| <input checked="" type="checkbox"/> French grunt | <input checked="" type="checkbox"/> Queen parrotfish | <input type="checkbox"/> Bluelip parrotfish | <input checked="" type="checkbox"/> Midnight parrot |
| <input checked="" type="checkbox"/> Princess parrotfish | <input type="checkbox"/> Bucktooth parrot | <input type="checkbox"/> Striped parrotfish | <input checked="" type="checkbox"/> Redtail parrotfish |
| <input checked="" type="checkbox"/> Rainbow parrot | <input type="checkbox"/> Reef croaker | <input checked="" type="checkbox"/> Redband parrotfish | <input type="checkbox"/> Jackknife fish |
| <input type="checkbox"/> Redfin parrotfish | <input checked="" type="checkbox"/> Graysby | <input checked="" type="checkbox"/> Stoplight parrot | <input type="checkbox"/> Barred hamlet |
| <input checked="" type="checkbox"/> Highhat | <input checked="" type="checkbox"/> Harlequin bass | <input type="checkbox"/> Blue hamlet | <input type="checkbox"/> Sand tiger shark |
| <input checked="" type="checkbox"/> Butter hamlet | <input checked="" type="checkbox"/> Great barracuda | <input checked="" type="checkbox"/> Red grouper | <input checked="" type="checkbox"/> Nassau grouper |
| <input type="checkbox"/> Black grouper | <input type="checkbox"/> Bonnethead | <input type="checkbox"/> Lantern bass | <input type="checkbox"/> Tarpon |
| <input checked="" type="checkbox"/> Jolthead porgy | <input checked="" type="checkbox"/> Southern stingray | <input checked="" type="checkbox"/> Sharpnose puffer | <input checked="" type="checkbox"/> Saucereye porgy |
| <input checked="" type="checkbox"/> Nurse shark | <input checked="" type="checkbox"/> Spotted moray | <input checked="" type="checkbox"/> Sharksucker | <input type="checkbox"/> Sand diver |
| <input checked="" type="checkbox"/> Southern stingray | <input type="checkbox"/> Sand tilefish | <input checked="" type="checkbox"/> Yellow stingray | <input checked="" type="checkbox"/> Squirrelfish |
| <input checked="" type="checkbox"/> Green moray | <input type="checkbox"/> Reef squirrelfish | <input type="checkbox"/> Snook | <input type="checkbox"/> Blackbar soldierfish |
| <input checked="" type="checkbox"/> Trumpetfish | <input checked="" type="checkbox"/> Shy hamlet | <input checked="" type="checkbox"/> Houndfish | <input type="checkbox"/> Glasseye snapper |
| <input type="checkbox"/> Longspine Squirrelfish | <input type="checkbox"/> Tiger grouper | <input checked="" type="checkbox"/> Tobaccofish | <input type="checkbox"/> Red hind |
| <input checked="" type="checkbox"/> Fairy basslet | <input checked="" type="checkbox"/> Graysby | <input checked="" type="checkbox"/> Bigeye | <input checked="" type="checkbox"/> Yellowjack |
| <input type="checkbox"/> Marbled grouper | <input checked="" type="checkbox"/> Greater amberjack | <input type="checkbox"/> Yellowmouth grouper | <input checked="" type="checkbox"/> Permit/pompano |
| <input checked="" type="checkbox"/> Greater soapfish | <input checked="" type="checkbox"/> Mutton snapper | <input type="checkbox"/> Coney | <input type="checkbox"/> Dog snapper |
| <input checked="" type="checkbox"/> Bluerunner | <input checked="" type="checkbox"/> Cottonwick grunt | <input type="checkbox"/> African pompano | <input checked="" type="checkbox"/> Margate |
| <input checked="" type="checkbox"/> Lane snapper | <input checked="" type="checkbox"/> Spotted drum | <input checked="" type="checkbox"/> Vermillion snapper | <input checked="" type="checkbox"/> Redspotted hawkfish |
| <input checked="" type="checkbox"/> Sailor's choice grunt | <input checked="" type="checkbox"/> Spotted trunkfish | <input checked="" type="checkbox"/> Tripletail | <input checked="" type="checkbox"/> Porcupinefish |
| <input checked="" type="checkbox"/> Spotted goatfish | <input type="checkbox"/> Whitespotted filefish | <input checked="" type="checkbox"/> Atlantic spadefish | <input type="checkbox"/> Balloonfish |
| <input checked="" type="checkbox"/> Gray triggerfish | <input type="checkbox"/> Lesser electric ray | <input checked="" type="checkbox"/> Smooth trunkfish | <input checked="" type="checkbox"/> Needlefish |
| <input checked="" type="checkbox"/> Scrawled cowfish | <input type="checkbox"/> Pinfish | <input type="checkbox"/> Striped burrfish | <input checked="" type="checkbox"/> Mullet |
| <input checked="" type="checkbox"/> Flyingfish | <input type="checkbox"/> Scorpionfish | <input checked="" type="checkbox"/> Ballyhoo | <input type="checkbox"/> Lookdown |
| <input checked="" type="checkbox"/> Anchovies | <input checked="" type="checkbox"/> Blackfin tuna | <input checked="" type="checkbox"/> Lined seahorse | <input type="checkbox"/> King mackerel |
| <input checked="" type="checkbox"/> Mojarras | <input checked="" type="checkbox"/> Rainbow runner | <input type="checkbox"/> Sheepshead | <input checked="" type="checkbox"/> Blacktip shark |
| <input checked="" type="checkbox"/> Dolphinfin | <input checked="" type="checkbox"/> Dusky squirrelfish | <input type="checkbox"/> Little tunny | <input checked="" type="checkbox"/> Bigeye scad |
| <input type="checkbox"/> Sargassumfish | <input checked="" type="checkbox"/> Yellowfin mojarra | <input checked="" type="checkbox"/> Hammerhead shark | <input checked="" type="checkbox"/> False pilchard |
| <input type="checkbox"/> Green turtle | <input checked="" type="checkbox"/> Jewfish | <input checked="" type="checkbox"/> Almaco Jack | <input checked="" type="checkbox"/> Sharksucker |
| <input checked="" type="checkbox"/> Loggerhead turtle | <input checked="" type="checkbox"/> Night sergeant | <input checked="" type="checkbox"/> Cobia | <input checked="" type="checkbox"/> Pipefish |
| <input type="checkbox"/> Hawksbill turtle | <input checked="" type="checkbox"/> Sharksucker | | <input checked="" type="checkbox"/> Mosquitofish |

REQUIRED FISH LIST
FIELD BIOLOGY

LEARN TO RECOGNIZE THE FOLLOWING FISHES AND THEIR FAMILIES.
THEY COMPRISE OVER 80% OF THE FISH IN REEF AREAS.

ORECTOLOBIDAE--nurse sharks

Ginglymostoma cirratum--nurse shark

DASYATIDAE--stingrays

Dasyatis americana--southern stingray

Urolophus jamaicensis--yellow stingray

MURAENIDAE--morays

Gymnothorax funebris--green moray

SELONIDAE--needlefishes

Tylosaurus crocodilus--houndfish

EXOCOETIDAE--flyingfishes and halfbeaks

Hemiramphus brasiliensis--ballyhoo

FISTULARIIDAE--cornetfishes

Fistularia tabacaria--bluespotted cornetfish

AULOSTOMIDAE--trumpetfishes

Aulostomus maculatus--trumpetfish

HOLOCENTRIDAE--squirrelfishes and soldierfishes

Holocentrus ascensionis--squirrelfish

Holocentrus rufus--longspine squirrelfish

Holocentrus coruscus--reef squirrelfish

Myripristis jacobus--blackbar soldierfish

SERRANIDAE--seabasses

Epinephelus striatus--nassau grouper

GRAMMISTIDAE--soapfishes

Rypticus saponaceus--greater soapfish

LUTJANIDAE--snappers

Lutjanus apodus--schoolmaster

Lutjanus griseus--gray snapper

Lutjanus synagris--lane snapper

Ocyurus chrysurus--yellowtail snapper

PRIACANTHIDAE--bigeyes

Priacanthus cruentatus--glasseye snapper

APGONIDAE--cardinalfish

Apogon maculatus--flamefish

CARANGIDAE--jacks, scads, and pompanos

Caranx ruber--bar jack

Caranx bartholomaei--yellow jack

Caranx crysos--blue runner

GERRIDAE--mojarras

Eucinostomus argenteus--spotfin mojarra

POMADASYIDAE--grunts
Haemulon flavolineatum--french grunt
Anisotremus virginicus--porkfish
 SCIAENIDAE--drums
Equetus acuminatus--highhat drum
 MULLIDAE--goatfishes
Mulloidichthys martinicus--yellow goatfish
Pseudupeneus maculatus--spotted goatfish
 KYPHOSIDAE--sea chubs
Kyphosus sectatrix--bermuda chub
 CHAETODONTIDAE--butterfly fishes
Chaetodon capistratus--four eye butterflyfish
C. ocellatus--spotfin butterflyfish
C. sedentarius--reef butterflyfish
C. striatus--banded butterflyfish
 POMACANTHIDAE--angelfishes
Holocanthus bermudensis--blue angelfish
H. ciliaris--queen angelfish
H. tricolor--rock beauty
Pomacanthus paru--french angelfish
P. arcuatus--gray angelfish
 POMACENTRIDAE--damselfishes
Abudefduf saxatilis--sergeant major
Chromis cyanea--blue chromis
Pomacentrus fuscus--dusky damselfish
P. variabilis--cocoa damselfish
P. partitus--bicolor damselfish
P. planifrons--three spot damselfish
Microspathodon chrysurus--yellowtail damselfish
 LABRIDAE--wrasses
Bodianus rufus--spanish hogfish
Lachnolaimus maximus--hogfish
Halichoeres garnoti--yellowhead wrasse
H. bivittatus--slippery dick
Thalassoma bifasciatum--bluehead wrasse
 SCARIDAE--parrotfishes
Sparisoma viride--stoplight parrotfish
Scarus vetula--queen parrotfish
S. croicensis--striped parrotfish
 ACANTHURIDAE--surgeonfishes
Acanthurus chirurgus--doctorfish
A. bahianus--ocean surgeon
A. coeruleus--blue tang
 GOBIIDAE--gobies
Gobiosoma oceanops--neon goby

SCORPAENIDAE--scorpionfishes

Scorpaena plumieri--spotted scorpionfish

SPHYRAENIDAE--barracudas

Sphyraena barracuda--great barracuda

BALISTIDAE--triggerfishes

Balistes vetula--queen triggerfish

B. capriscus--gray triggerfish

MONACANTHIDAE--filefishes

Aluterus scriptus--scrawled filefish

A. schoepfi--orange filefish

OSTRACIIDAE--trunkfishes

Lactophrys quadricornis--scrawled cowfish

L. trigonus--truckfish

TETRADONTIDAE--puffers

Canthigaster rostrata--sharpnose puffer

DIODONTIDAE--porcupinefishes

Diodon histrix--porcupinefish

Chilomycterus schoepfi--striped burrfish

1991 Species

Green Algae

- _x_Disk Algae, *Halimeda*
- Caulerpa*
- _x_Sea grapes
- _x_Sea feathers
- _x_Mermaid's fan, *Udotea*
- ___Sea bottles, *Valonia*
- ___Green bubble algae, *Dictyosphaeria*
- _x_Neptune's shaving brush, *Penicillus*
- _x_Gigantic shaving brush, *P. dumentosus*
- _x_"Mossy algae", *Cladophoropsis*
- ___Dead man's fingers, *Codium*
- ___*Avrainvillea*
- _x_Mermaid's wine glass, *Acetabularia*
- Brown Algae**
- ___Strap browns, *Dictyota*
- _x_Brown whorls and fans, *Padina*
- _x_Gulfweed, *Sargassum*
- Red Algae**
- ___Goniolithon, coralline, crunch, crunch
- ___*Eucheuma*, cartilage
- ___Red coralline, *Jania*
- _x_Fragile coralline, *Amphiroa*
- _x_*Laurencia*
- Higher Plants**
- Sea grasses**
- _x_Turtle grass
- _x_Manatee grass
- _x_Shoot grass
- Mangroves**
- _x_Red mangrove
- _x_White mangrove
- _x_Black mangrove
- _x_Buttonwood
- Salt strand plants**
- _x_Sea purslane
- _x_Saltwort
- _x_Glasswort
- _x_Sea oxeye
- _x_Railroad vine
- _x_Sea grapes
- _x_Gumbo limbo
- _x_Spanish bayonet
- _x_Australian pine
- _x_Sea oats
- _x_Saw palmetto
- _x_Prickly pear cactus
- _x_Brazilian pepper
- _x_Cabbage palm
- _x_Spanish needles
- _x_Puncture vine
- _x_Strangler fig
- _x_*Melaleuca*, paper or punkwood tree
- ___Beach sunflower

Sponges

- _x_Loggerhead sponge
- _x_Vase sponge
- _x_Heavenly sponge
- ___Fire sponge
- ___Chicken liver sponge
- ___Stinker sponge
- _x_Green sponge
- ___Sheepswool sponge
- _x_Commercial sponge
- _x_Tube sponge
- ___Variable sponge
- Cnidarians**
- Hydrozoans**
- _x_Portuguese man-o-war
- ___By the wind sailor
- Hydrozoan "coral"**
- _x_Fire coral
- Scyphozoans**
- _x_Moon jellyfish
- _x_Upsidedown jellyfish
- Anthozoans**
- Gorgonians or soft corals**
- _x_Sea fans
- _x_Sea rods
- _x_Sea whips
- _x_Corky sea fingers
- Zoanthid mat anemones**
- _x_Mat anemone
- ___Cheerio coral
- Actinarians**
- _x_Pink tipped anemone
- _x_Ringed anemone
- Hard corals**
- ___Elkhorn coral
- ___Staghorn coral
- _x_Rose coral
- _x_Mountainous star
- _x_Brain
- _x_Starlet corals
- _x_Finger corals
- _x_Clubbed finger
- _x_Tube corals
- Comb jellies**
- _x_Lobate comb jelly
- Annelids**
- _x_Horned feather worm
- ___Green bristle worm
- Sipunculids**
- ___Peanut worm
- Mollusks**
- Chitons**
- _x_Fuzzy chiton
- ___Ischnochiton
- Sea hares**
- _x_Spotted sea hare
- Archeogastropod snails**
- _x_American star
- _x_Long-spined star
- _x_Nerites
- Mesogastropod snails**
- ___Black horn shells
- _x_Echinus littorinids
- ___Flamingo tongue
- _x_Stocky cerith
- _x_Queen conch
- ___Hawkwing conch
- _x_Beaded periwinkle
- ___Worm snail
- Neogastropods**
- ___Lightning whelks
- ___Horse conch
- _x_True tulip
- _x_Caribbean vase
- _x_Florida rock snail
- Pulmonate snail**
- ___Striped false limpet
- _x_Keyhole limpet
- Bivalves**
- ___Stiff pen shell
- _x_Turkey wing
- _x_Lister's tree oysters
- _x_Tulip mussel
- _x_Lucines
- _x_Cockles
- Cephalopods**
- ___Octopus
- _x_Reef squid
- Bryozoans**
- ___Whorled zoobotryon
- Arthropods**
- _x_Horseshoe crab
- Barnacles**
- _x_Ribbed
- ___Fragile or starred
- ___Ivory
- _x_Gooseneck
- Stomatopods**
- ___Mantis shrimp
- ___Swollen claw squilla
- Isopods**
- _x_Sea roach
- Amphipods**
- _x_Beach hoppers
- Penaeid shrimp**
- _x_Commercial pink
- Caridean shrimp**
- ___Grass shrimp
- ___Bumblebee shrimp
- _x_Snapping shrimp
- Lobsters**
- _x_Florida spiny
- _x_Spanish or shovel-nosed
- Hermit crabs**
- _x_Giant hermit
- ___Land hermit
- _x_Striped hermit
- Mole crab**
- ___Emerita
- Portunid crabs**
- _x_Blue crab
- Xanthid crabs**
- ___Common mud crab
- _x_Stone crab
- Grapsoid crabs**
- _x_Mottled shore crab
- ___Mangrove crab

Ghost and fiddler crabs

- _x_Ghost crab
- _x_Fiddler crab
- Spider crabs**
- _x_Arrow crab
- _x_Spider crab
- _x_Decorator crab
- ___Two-horned spider
- Echinoderms**
- Sea Stars**
- ___Cushion stars
- ___Thorny and spiny
- Sea urchins**
- ___Long spined
- ___Pencil
- ___Sea egg
- ___Variegated
- _x_Rock boring
- ___Sea bisquits
- Brittle stars**
- ___Slimy brittle star
- ___Basket starfish
- Sea cucumbers**
- ___Florida sea cucumber
- Sea squirts**
- _x_Black tunicate
- Vertebrates**
- Fish--Separate List**
- Birds--Separate List**
- Mammals**
- ___Whales
- _x_Dolphins
- _x_Key deer
- _x_Raccoons
- _x_Smallest gray squirrel
- _x_Rats
- Herps**
- _x_Brown anole
- _x_Green anole
- ___Bahamian curlytail
- _x_Ge kos
- ___Crocodile
- _x_Alligator
- ___Marine toad
- ___Other toads
- _x_Tree frogs
- Insects & other**
- Inverts**
- Butterflies**
- See separate list
- Dragonflies**
- ___Arrowhead damsel
- ___Forktail damsel
- ___Keys red
- ___Red-faced
- ___Antillean green-faced
- ___Marine
- ___Olive-faced
- ___Black-headed red
- ___White-faced
- ___Violet-faced
- ___Cosmopolitan
- ___Red and green-faced
- _x_Mosquitoes
- bigger than cars**
- _x_Palmetto bugs = roaches

KINGDOM PLANTAE, DIVISION CHLOROPHYTA, CLASS CHLOROPHYCEAE

GREEN ALGAE

1. THE HALIMEDAS

Halimeda opuntia=cactus Halimeda; often calcified with numerous branches running in different planes. 2-3 lobes. Look for branches running in different planes.

Halimeda incrassata. Thick, heavily calcified. 1 plane of growth. 3 lobes and ribbed. Look for 3 lobes and ribs.

Halimeda monile. Flat segments with a rounded upper joint. Look for flat segments with a rounded upper joint.

Halimeda tuna. One plane. Large flat, round to kidney-shaped joints. Look for large, flat kidney-shaped or round joints.

2. THE CAULERPAS

Caulerpa sertularioides. Open sandy areas. Beautiful with long feathery blades.

Caulerpa paspaloides. Four brushes on a stalk. Triangular in appearance.

Caulerpa racemosa. Found clinging to rock. Clusters of small green grapes.

3. PENICILLUS=SHAVING BRUSH ALGAE

Penicillus capitatus=Neptune's shaving brush. Distinctive.

Penicillus dumentosus=Large shaving brush. Distinctive.

Avrainvillea--Looks like shaving brush, but is large and fanlike. Dark brown to black in color.

4. UDOTEA=MERMAID'S FAN

Udotea flabellum=Mermaid's fan. Sandy areas and grass beds. Broadly fan-shaped on a short stalk.

5. ACETABULARIA=MERMAID'S WINEGLASS

Acetabularia crenulata=Mermaid's wineglass. Very distinctive.

6. CODIUM=DEAD MAN'S FINGERS

Codium taylori. Glistening green fingers or tubes.

7. VALONIA=SEA BOTTLES

Valonia ventricosa=Sea bottles. Elliptical, transparent sacs.

Dictyosphaeria cavernosa--Green bubble algae. Look for a large, green spherical mass growing in the intertidal and below

Cymopolia--Distinctive, unusual algae with tubular calcareous joints. Each branch ends in a tuft of green filaments

KINGDOM PLANTAE, DIVISION PHAEOPHYTA, CLASS PHAEOPHYCEAE

BROWN ALGAE

Dictyota dichotoma--Flat straplike branches always branching dichotomously.

Padina sanctaecrucis--Curled, fanlike branches from a single stalk.

DIVISION PHAEOPHYTA, CLASS PHAEOPHYCEAE

SARGASSUM OR GULFWEED

Sargassum filipendula--Gulfweed is familiar to everyone who has walked the beach or seen the floating masses at sea. This form, one of 15 or more species, is golden brown and found growing attached to the rocks in the Keys.

KINGDOM PLANTAE, DIVISION RHODOPHYTA, CLASS RHODOPHYCEAE

Laurencia papillosa--This red alga is often found growing along the edges of rocky platforms. It is densely clustered, olive-green to greenish purple. The branches have small tubercles.

KINGDOM PLANTAE, DIVISION RHODOPHYTA, CLASS RHODOPHYCEAE

CORALLINE RED ALGAE

The corallines are reds so heavily calcified that they have lost their resemblance to algae.

Goniolithon--Forms hard encrustations on coral and rock. We'll see a form at West Summerland Key that becomes masses of closely entangled thin branches. This form is mistaken for coral.

Amphiroa fragilissima--Slender coralline branchlets. As with the above, they go "crunch, crunch" under the foot.

Jania rubens--red coralline. Rose red, delicate, with a height of 1-2 inches.

Kingdom Plantae, Salt Strand Plants. Look for these plants living at the water's edge.

Batis maritima--Saltwort. Look for a pale green, succulent shrub. Crushed leaves give off a scent.

Sesuvium portulacastrum--Sea purslane. Succulent plants mostly lying on the ground.

Salicornia--Glasswort. Fleshy, shrubby. Crackle under the foot.

Kingdom Plantae, The Sea Grasses

Thalassia testudinum--Turtle grass. Look for larger flat blades.

Syringodium filiforme--Manatee grass. Look for round blades.

Halodule wrightii--Shoal grass. Look for smaller flat blades.

Kingdom Plantae, The Mangroves. An ecological rather than taxonomic designation. These are trees adapted for living in saltwater environments.

Conocarpus--Buttonwood. Shaggy bark often with a twisted trunk. Leaves a pale grayish green. My general impression is that they are narrower and more sharply pointed than the other mangroves.

Laguncularia--White mangrove. Rounded leaves with indented tips. A pair of swollen salt glands at the base of the leaf.

Rhizophora mangle--Red mangrove. Smooth leathery leaves. Prop roots. Distinctive seedling.

Avicennia germinans--Black mangrove. Lanceolate leaves. Characteristic aeration roots called pneumatophores.

Kingdom Animalia, Phylum Porifera, Class Demospongiae

Spherospongia vesparia--Loggerhead sponge

This is the largest sponge I have seen. Usually with a cake shape and a flat top and central depression. The dark holes at the top are called oscula. The species is the host of many marine animals--especially snapping shrimp.

Kingdom Animalia, Phylum Porifera, Class Demospongia

Ircinia campana--Vase sponge.

Phylum Cnidaria, Class Hydrozoa, Order Hydroida

CNIDARIA

This phylum includes jellyfish, sea anemones, hydrozoans, and corals. Although extremely variable in appearance, all members have a radially symmetrical body plan. The saclike body has a central stomach cavity with a single opening that is usually surrounded by food-capturing tentacles.

Classification of major reef benthic Cnidaria

PHYLUM CNIDARIA (COELENTERATA)

CLASS HYDROZOA

ORDER MILLEPORINA: fire corals

ORDER STYLASTERINA: hydrocorals

CLASS ANTHOZOA

SUBCLASS OCTOCORALLIA (ALCYONARIA) *sensu* Bayer 1961

ORDER STOLONIFERA

ORDER TELESTACEA

ORDER ALCYONACEA: fleshy soft corals

ORDER GORGONACEA: sea whips, sea feathers, sea fans, sea plumes, other gorgonian corals

ORDER PENNATULACEA: sea pens

SUBCLASS ZOANTHARIA (HEXACORALLIA)

ORDER ACTINIARIA: anemones

ORDER CORALLIMORPHARIA: false coral anemones

ORDER ZOANTHIDEA: carpet anemones

ORDER CERIANTHARIA: tube anemones, often parasitic in other organisms

ORDER SCLERACTINIA (MADREPORARIA): true stony corals

ORDER ANTIPATHARIA: black or thorny corals

Kingdom Animalia, Phylum Cnidaria, Class Scyphozoa

Cassiopeia xamachana--Upsidedown jellyfish. Distinctive.

Lies upside down to provide necessary sunlight to farm algae.

Aurelia aurita--Moon jellyfish. Commonly washed ashore.

Look for it in the water column around the reefs and grass flats.

Class Anthozoa, Order Zoanthidea

Zoanthus pulchellus--Mat anemone. Matlike pretty bluish to greenish colonies attached to rocks at low tide line.**Palythoa mammillosa**--Knobby zoanthidean or "cheerio coral".

Form on dead coral in the reef zone.

Kingdom Animalia, Phylum Cnidaria, Class Anthozoa, Subclass Alcyonaria

Pterogorgia anceps--Angular sea whip. Triangular with beautiful purple color. This is the common soft coral at West Summerland Key.

Kingdom Animalia, Class
Anthozoa, Subclass Alcyonaria
Gorgonia ventalina--Common sea
fan

Flat anastomosed colony; usually
purple in appearance. This one is
fairly common around the reefs
in the keys.

Kingdom Animalia, Phylum
Ctenophora

Mnemiopsis mccradyi--Lobate
comb jelly.

These transparent jellies are
often found by the thousands in
shallow water. Often they are
bioluminescent at night.

Phylum Cnidaria, Class Anthozoa, Subclass Zoantharia, Order
Madreporaria

Manicina areolata--Rose coral. The small coral is one that you will
find close to shore. Can tolerate some turbidity.

Siderastrea--Starlet coral. Look for small clumps in shallow areas.
The rounded masses may reach one to two feet in diameter.

Kingdom Animalia, Phylum Annelida.

Green bristle worms

Look for a squarish body,
greenish color, orange-red gills.
Handle this one with care--if at
all. The bristles easily

penetrate gloves to cause painful
stings.

Feather worms

Look for single opening of tube
which is glued together with
pieces of leaves, mangrove
detritus etc.

Kingdom Animalia, Phylum Mollusca, Class Polyplacophora

Ischnochiton papillosus--Slender, white chiton found at West
Summerland Key.

Acanthopleura granulata--Fuzzy chiton. Girdle with low, hairlike
spines.

Kingdom Animalia, Phylum Cnidaria, Class Anthozoa, Subclass
Zoantharia, Order Actinaria

Condylactis gigantea--Pink-tipped anemone. The tentacles are
always tipped in pink or blue.

Kingdom Animalia, Phylum Bryozoa

Zoobotryon verticillatum--Whorled Zoobotryon. These animals form
masses resembling algae on shoreline structures.

Kingdom Animalia, Phylum Mollusca, Class Gastropoda, Subclass Opisthobranchia.

Aplysia dactylomela--spotted sea hare. Distinctive. Punch it and watch it emit clouds of deep purple ink.

Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Amphipoda, Orchestia--beach hoppers

Kingdom Animalia, Phylum Chordata, Subphylum Tunicata, Class Ascidiacea Ascidia nigra--Black tunicate. Easily recognized by its blue to black test.

Kingdom Animalia, Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Isopoda.

Ligia exotica--Sea roach. Runs rapidly over rocks or sea walls.

Kingdom Animalia, Phylum Sipunculida

Phascolosoma antillarum--Peanut worm. Remarkable powers of extension and scrunching up. Bores into soft beach rock.

Phylum Arthropoda, Class Merostomata, Limulus polyphemus--horse-shoe crab. Familiar to all sandbar waders and beach walkers. Actually the horse-shoe crab is more related to spiders than to crabs.

Class Crustacea, Subclass Malacostraca, Order Stomatopoda

The mantis shrimp or squillas are often called "thumbsplitters" because they can use their raptorial claws to inflict a nasty wound. Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Decapoda, Suborder Natantia, Section Penaeidea

Penaeus duorarum--The commercial pink shrimp has characteristic grooves running along the carapace.

Caridean shrimp like the grass shrimp, Tozeuma, have a second or abdominal segment which overlaps both the preceding plate and the one behind.

Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Decapoda Suborder Natantia, Section Caridea

The Bumblebee shrimp lives on sea urchins in grassy areas. It has a cutoff look. The entire body is covered with deep chocolate bands separated by narrow yellow lines.

The common snapping shrimp is recognized by the red splotch at the end of the large claw. You can hear the snaps at night or around loggerhead sponges where these species often fill up the galleries and holes in the sponge.

Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Decapoda,

Suborder Reptantia, Division Palinura

Fanulirus argus--Florida spiny lobster, crawfish, "bug"

Scyllarides--Spanish or shovel-nosed lobster

Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Decapoda,

Suborder Reptantia, Division Anomura

Clibanaris vittatus--Striped hermit crab. Noticeable gray to white stripes on the legs.

Emerita talpoida--Mole crab. Catches the waves in the swash zone to grab detritus. It then quickly burrows back into the sand.

Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Decapoda, Suborder Reptantia, Division Anomura

Coenobita clypeatus--Land hermit crab. One of the few terrestrial hermit crabs. Purple left claw with orange dactyls.

Petrochirus diogenes--Giant hermit crab. Look for this monster in conch shells. The scaly chelipeds are diagnostic.

Phylum Arthropoda, Class Crustacea, Subclass Malacostraca, Order Decapoda

Suborder Reptantia, Division Brachyura--True Crabs

Crancoid crabs--Oval to circular in shape with spines on the anterior end.

Family Portunidae--Portunidae crabs are swimmers because the last appendage is modified into a paddle.

Grapsoid crabs are squarish with no or insignificant spines.

The ghost crab is the most common crab along the beaches--especially in the evening. The overall color is the color of a tan sand beach.

The mangrove or tree crab is small--about an inch. Look for a patch of hairs on the claws. Carapace turned inward posteriorly. Color on moveable claw extends back into palm. These are true crabs in the Family Xanthidae.

Panopeus is the common mud crab. Watch these. They have massive claws that can pinch the Hell out of you.

Menippe is the commercial crab you've heard so much about. The young are bluish purple with a white spot on the wrist. This is the common fiddler crab. Rough claw with no ridge. Also known as the mottled shore crab. The little crab has a converging posterior with tubercular lines on its carapace. The claws are smooth.

The Spider Crabs of the Superfamily Oxyrhyncha

The Arrow crab is a distinctive little crab which crawls slowly over the bottom.

Look for the big spider crabs in shallow water. The rostrum ends in two points. There are six spines on each side.

The grass crab or sponge crab is a type of "decorator crab". It attaches all sorts of objects to its rostrum and back. Sponges often grow on its back. Look for three spines at the rear.

The 2-horned spider has a spotted cheliped--even in alcohol..Spines at the angles are small.

Phylum Echinodermata, Class Asteroidea

Cushion star--Bright red starfish. Our best known starfish. Overcollected in many areas.

Phylum Echinodermata, Class Echinoidea, Subclass Regularia

Sea Urchins.

Both of the above are easily recognized. Look for long black spines and don't stick your hand or foot in the "Antillean crown of thorns".

Look for blunt spines in the slate-pencil urchin. You can pick this one up.

Phylum Echinodermata, Class Echinoidea, Subclass Regularia

Find an urchin with objects attached to its back, and you have found a variegated urchin. It is a master at camouflage.

The sea egg is distinctive with short white spines on a brown test.

Phylum Echinodermata, Class Echinoidea, Subclass Regularia

The Rock-boring urchin is elliptical in outline with strong, thick pointed spines. Look for in holes along the tide line.

Subclass Irregularia--Sea biscuits and sand dollars.

There are not many sand dollars in the Keys.

The sea biscuit is oval on top, but it is flattened underneath. This one is common in several habitats.

Basket starfish--Distinctive and easily recognized. Look for clinging to sea fans etc. By day, they may be curled into a tight knot.

Phylum Echinodermata, Class Ophiuroidea--Brittle Stars.

Vegetation Observed by the 1991 Field
Biology Class

Common Name	UK	LK	Vegetation continued		
___sweet acacia	—	P	___x_white mangrove	H	H,P
___pineland acacia	--	P	___wild sage	H	H,P
___torchwood		H	___x_wild tamarind	H	--
___pond apple	H	P	___wild dilly	H	H,P
___marlberry	H	P	___mastic	H	--
___crabwood	H	--	___gutta-percha mayten	H	--
___x_black mangrove	H	H	___x_polsonwood	H	H,P
___false willow	--	P	___twinberry stopper	H	--
___groundsel tree	—	H	___wax myrtle	—	H,P
___groundsel tree	H	H,P	___x_myrsine	H	H,P
___smooth strongbark	--	P	___lancewood	H	--
___strongbark	H	H	___red bay	H	P
___rough strongbark	H	--	___x_S. Fla, slash pine	—	H,P
___saffron plum	H	H,P	___x_Jamaica dogwood	H	H,P
___willow bustic	H	H,P	___pisonia	--	H,P
___x_gumbo limbo	H	H,P	___blackhead	—	P
___locust berry	H	H,P	___x_cat's claw	H	H
___beauty berry	H	--	___long-stalked stopper	—	H,P
___pale lidflower	H	--	___Bahama wild coffee	H	--
___myrtle-of-the-river	H	--	___x_live oak	H	--
___cinnamon-bark	H	--	___indigo berry	H	H,P
___Jamaica caper	H	--	___darling plum	H	H,P
___limber caper	H	H	___x_red mangrove	H	H,P
___seven-year apple	H	--	___sumac	—	P
___Bahama senna	—	H,P	___soapberry	H	--
___x_coco plum	H	P	___maiden bush	—	H
___satinleaf	H	H,P	___Florida boxwood	H	--
___fiddlewood	H	--	___gray twig	H	--
___pitch apple	—	H	___paradise tree H	H	
___pigeon plum	H	H,P	___Bahama nightshade	H	--
___x_sea grape	H	H,P	___potato-tree	H	H,P
___soldierwood	H	-	___necklace pod	—	P
___x_buttonwood	H	H,P	___x_mahogany	H	--
___x_Geiger tree	H	H	___West Indies trema	H	--
___rhacoma	H	H,P	___Florida trema	—	H,P
___pineland croton	—	P	___pearl berry	H	--
___cupania	—	H	___tallowwood	H	H
___varnish leaf	H	--	___wild lime	H	--
___milkbark	H	H,P	___yellow nicker	—	H
___Guiana plum	H	--	___x_gray nicker	H	--
___black torch	H	H,P	___snowberry	H	H
___x_white stopper	H	H,P	___fishpoison vine	H	--
___redberry stopper	H	--	___chew stick	H	--
___Spanish stopper	H	H,P	___doctor vine	H	--
___red stopper	H	--	___x_Virginia creeper	H	H
___princewood	H	H	___devil's claw	H	H
___inkwood	H	H	___greenbrier	H	H,P
___x_strangler fig	H	H,P	___soldier vine	H	H
___shortleaf fig	H	H,P	___x_poison ivy	H	H,P
___pineland olive	—	P	___muscadine	H	H
___lignum vitae	H	--	___x_silver palm	H	H,P
___blolly	H	H,P	___buccaneer palm	H	-
___velvetseed	H	H,P	___x_royal palm	—	—
___rough velvetseed	H	H,P	___x_cabbage palm	H	H,P
___false boxwood	—	H	___x_saw palmetto	H	H,P
___scarletbush	H	H	___x_Key thatch palm	H	H,P
___manchineel	—	H,P	___x_Florida thatch palm	H	H,P
___white ironwood	H	P			
___joewood	H	H,P			
___black ironwood	H				

Plant List for Field
Biology Fresh Water
Swamps and Wetlands

Common Name

Air plants or
Bromeliads

Alligator Flag or
Arrowroot
Arrowhead

Bald Cypress

Bladderwort
Bog Hemp or
Wild Hemp Vine

Butterfly Orchid

Cattail

Duckweed
Elderberry
Hat Pin or
Pipewort
Leather Fern

Milkweed Vine or
Climbing Milkweed
Milkweed

Mosquito Fern
Paper Tree
or Punkwood

Pennywort

Pickerel Weed

Pond Apple

Primrose Willow

Scientific Name

Tillandsia (11 spp.)

Thalia geniculata

Sagittaria (7 spp.)

Taxodium distichum

Utricularia (5 spp.)

Mikania (3 spp.)

Encyclia tampensis

Typha (3 spp.)

Lemna (2 spp.)

Sambucus simpsonii

Eriocaulon (3 spp.)

Acrostichum (2 spp.)

Cynanchum (4 spp.)

Asclepeas (12 spp.)

Azolla caroliniana

Melaleuca

quinquinervia

Hydrocotyle (3 spp.)

Pontedaria lanceolata

Annona glabra

Ludwigia peruviana

Family

Bromeliaceae

Pineapple Family

Marantaceae

Arrowroot Family

Alismataceae

Water plantain

Taxodiaceae

Bald Cypress Family

Asteraceae

(Compositae)

Aster Family

Orchidaceae

Orchid Family

Typhaceae

Cattail Family

Lemnaceae

Caprifoliaceae

Eriocaulaceae

Pipewort Family

Pteridaceae

Bracken Fern Family

Asclepeadaceae

Milkweed Family

Asclepiadaceae

Milkweed Family

Salviniaceae

Myrtaceae

(introduced

Myrtle Family from

Australia)

Apiaceae

Celery Family

Pontederiaceae

Pickerelweed Family

Annonaceae

Custard Apple Family

Onagraceae

Evening Primrose

Family

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Red Bay and Swamp Bay	<u><i>Persea</i> (2 spp.)</u>	Lauraceae Laurel Family
Red Maple	<u><i>Acer rubrum</i></u>	Aceraceae Maple Family
Royal Palm	<u><i>Roystonea elata</i></u>	Arecaceae (Palmae) Palm Family
Saw Grass	<u><i>Cladium jamaicensis</i></u>	Cyperaceae Sedge Family
Spanish Moss	<u><i>Tillandsia usneoides</i></u>	Bromeliaceae Pineapple Family
Spatterdock or Yellow Water Lilly	<u><i>Nuphar luteum</i></u>	Nymphaeaceae Water Lilly Family
Spider Lilly	<u><i>Hymenocallis</i> (2 spp.)</u>	Amaryllidaceae Amaryllis Family
Spike Rush	<u><i>Eleocharis</i> (7 spp.)</u>	Cyperaceae Sedge Family
Swamp Fern	<u><i>Blechnum serrulatum</i></u>	Blechnaceae Blechnum Family
Swamp Lilly	<u><i>Crinum americanum</i></u>	Amaryllidaceae Amaryllis Family
Water Arum	<u><i>Peltandra virginica</i></u>	Arum Family Araceae
Water Fern	<u><i>Salvinia rotundifolia</i></u>	Salviniaceae
Water Hyacinth	<u><i>Eichornia crassipes</i></u>	Pontederiaceae (Introduced from South America) Pickerelweed Family
Water Lettuce	<u><i>Pistia stratiotes</i></u>	Araceae Arum Family
Water Lilly	<u><i>Nymphaea</i> (3 spp.)</u>	Nymphaeaceae Water Lilly Family
Wax Myrtle or Bayberry	<u><i>Myrica</i> (2 spp.)</u>	Myricaceae Bayberry Family
Willow	<u><i>Salix caroliniana</i></u>	Salicaceae Willow Family

Plant List for Field Biology Coastal Vegetation-Dunes- Common Name	<u>Scientific Name</u>	<u>Family</u>
Australian Pine	<u>Casuarina (3 spp.)</u>	Casuarinaceae (Introduced from Australia) Beefwood Family
Beach Bean or Bay Bean	<u>Canavalia maritima</u>	Leguminosae (Fabaceae)
Beach Peanut	<u>Okenia hypogaea</u>	Bean & Pea Family Nyctaginaceae
Beach Sunflower	<u>Helianthus debilis</u>	Four O'clock Family Asteraeae
Century Plant	<u>Agave (3 spp.)</u>	(Agavaceae Agave Family)
Cocoplum	<u>Chrysobalanus icaco</u>	Chrysobalanaceae
Greenbrier	<u>Smilax (4 spp.)</u>	Smilacaceae Smilax Family
Ground Cherry	<u>Physalis (5 spp.)</u>	Solanaceae Nightshade Family
Heliotropes	<u>Heliotropium (3 spp.)</u>	Boraginiaceae
Inkberry	<u>Scaevola plumieri</u>	Goodeniaceae
Necklacepod	<u>Sophora tomentosa</u>	Fabaceae Pea & Bean Family
Poisonwood	<u>Metopium toxiciferum</u>	Anacardiaceae Cashew Family
Poor Mans Patches	<u>Mentzelia floridana</u>	Loasaceae
Prickly Pear Cactus	<u>Opuntia (6 spp.)</u>	Cactaceae Cactus Family
Puncture Weed	<u>Tribulus cistiodes</u>	Zygophyllaceae Caltrop Family
Railroad Vine or Beach Morning Glory	<u>Ipomoea pes-caprae</u>	Convolvulaceae Morning Glory Family
Saw Palmetto	<u>Serenoa repens</u>	Palmae (Arecaceae) Palm Family
Sea Grape	<u>Coccoloba uvifera</u>	Polygonaceae Buckwheat Family
Sea Oats	<u>Uniola paniculata</u>	Poaceae (Gramineae) Grass Family
Spanish Bayonet	<u>Yucca aloifolia</u>	Agavaceae
Spider Lilly	<u>Hymenocallis (2 spp.)</u>	Amaryllidaceae Amaryllis Family
Tread Softly	<u>Cnidocolus</u>	Euphorbiaceae

Field Biology Animal List

-Gastropods-

<u>Common Name</u>	<u>Scientific Name</u>	<u>Habitat</u>
Apple Snail	<u><i>Pomacea spp.</i></u>	
Tree Snails	<u><i>Liguus fasciatus</i></u>	

-Spiders-

<u>Common Name</u>	<u>Scientific Name</u>	<u>Habitat</u>
Crab Spider	<u><i>Gasteracantha cancriformis</i></u>	
Golden Orb Weaver	<u><i>Nephila clavipes</i></u>	
Orchard spiders	<u><i>Leucauge spp.</i></u>	

-Insects-

<u>Common Name</u>	<u>Scientific Name</u>	<u>Habitat</u>
Cabbage Butterflies	<u><i>Various genera & species</i></u>	
Honey Bee	<u><i>Apis mellifera</i></u>	
Monarch Butterfly	<u><i>Danaus plexippus</i></u>	
Paper Wasp	<u><i>Polistes sp.</i></u>	
Queen Butterfly	<u><i>Danaus gilippus</i></u>	
Swallowtails	<u><i>Papilio spp.</i></u>	
Sulfur Butterflies	<u><i>Var. genera & species</i></u>	
Viceroy Butterfly	<u><i>Basilarchia archippus</i></u>	
Zebra Butterfly	<u><i>Heliconius charionius</i></u>	

Field Biology Animal List

-Amphibians-

<u>Common Name</u>	<u>Scientific Name</u>	<u>Habitat</u>
Giant Toad	<u>Bufo marinus</u>	
Southern Toad	<u>B. terrestris</u>	
Tree Frogs	<u>Hyla (5 spp. inc. green & squirrel)</u>	
True Frogs	<u>Rana (2 spp. pig & leopard)</u>	

-Reptiles-

<u>Common Name</u>	<u>Scientific Name</u>	<u>Habitat</u>
Brown Anole	<u>Anolis sagrei</u>	
Green Anole	<u>A. carolinensis</u>	
Bahamian Curly Tail Lizard	<u>Leiocephalus carinatus</u>	
Southeastern Five-Lined Skink	<u>Eumeces inexpectatus</u>	
Alligator	<u>Alligator mississippiensis</u>	
Florida Softshell Turtle	<u>Trionyx ferox</u>	
Water Snakes	<u>Natrix (3 spp.)</u>	
Cotton Mouth or Water Moccasin	<u>Agkistrodon piscivorus</u>	
Coral Snake	<u>Micrurus fulvius</u>	
Pigmy Rattlesnake	<u>Sistrurus miliarius</u>	
Eastern Diamondback Rattlesnake	<u>Crotalus horridus</u>	

Field Biology Animal List

-Birds-

Common Name

Brown Pelican
 Cormorant
 Anhinga
 Turkey Vulture
 Black Vulture
 Hawks (7 spp.)
 Osprey (fish hawk)
 Black Skimmer
 Kingfisher
 Great Blue Herron/Great White
 Heron
 Snowy Egret
 Cattle Egret
 Louisiana Heron
 Little Blue Heron
 Green Heron
 Cardinal

Common Name

Black-Crowned Night Heron
 Yellow-Crowned Night Heron
 White Ibis
 Gallinules (2 spp. common and
 purple)
 American Coot
 Sea Gulls (Herring, ring-billed,
 laughing, and boneparte's)
 Blue Jay
 Terns (about 10 spp.)
 Common or American Egret
 Crows (fish and common)
 Red-winged Blackbird
 Grackles (common and boat-
 tailed)
 Sandpipers (about 20 spp.)

-Mammals-

<u>Common Names:</u>	White-tailed	O'possum	Armadillo
Raccoon	deer	Eastern Fox	
Marsh Rabbit	Key deer	Squirrel	
	Eastern Grey		
	Squirrel		

Plant List for Field Biology

Scrub Forests

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Lichens (Terrestrial)		
Lovevine	<u><i>Cassytha filiformis</i></u>	Lauraceae Laurel Family
Oak	<u><i>Quercus</i> sp</u>	Fagaceae Beech Family
Prickly Pear Cactus	<u><i>Opuntia</i> (6 spp.)</u>	Cactaceae Cactus Family
Rosemary	<u><i>Ceratiola ericoides</i></u>	Empetraceae Crowberry Family
Sand Pine	<u><i>Pinus clausa</i></u>	Pinaceae Pine Family
Spike "Moss"	<u><i>Selaginella</i> (4 spp.)</u>	Selaginillaceae Spike Moss Family

Coastal Vegetation

-Mangroves-

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Australian Pine	<u><i>Casuarina</i> (3 spp.)</u>	Casuarinaceae (Introduced from Australia) Beefwood Family
Black Mangrove	<u><i>Avicennia germinans</i></u>	Avicenniaceae
Red Mangrove	<u><i>Rhizophora mangle</i></u>	Rhizophoraceae Mangrove Family
Buttonwood	<u><i>Conocarpus erecta</i></u>	Combretaceae
White Mangrove	<u><i>Laguncularia</i></u>	Combretaceae
	<u><i>racemosa</i></u>	
Cordgrass	<u><i>Spartina</i> (4 spp.)</u>	Poaceae (Gramineae)

-Strand-

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Brazilian Pepper	<u><i>Schinus</i></u> <u><i>terebinthifolius</i></u>	Anacardiaceae (Introduced from S. America) Cashew Family
Coconut Palm	<u><i>Cocos nucifera</i></u>	Aracaceae (Palmae) Palm Family
Dalbergia	<u><i>Dalbergia</i> (3 spp.)</u>	Fabaceae (Leguminosae) Pea and Bean Family

-Strand-		
Glasswort	<u><i>Salicornia</i></u> (2 spp.)	Chenopodiaceae Goosefoot Family
Leather Fern	<u><i>Acrostichum</i></u> (2 spp.)	Pteridaceae Bracken Fern Family
Nickerbean	<u><i>Caesalpinia</i></u> (3 spp.)	Leguminosae (Fabaceae) Pea and Bean Family
Saltbush	<u><i>Baccharis</i></u> (4 spp.)	Compositae (Asteraceae) Aster Family
Saltwort	<u><i>Batis maritima</i></u>	Bataceae Saltwort Family
Sea Ox-eye or Sea Daisy	<u><i>Borrchia</i></u> (2 spp.)	Compositae (Asteraceae) Aster Family
Sea Purslane	<u><i>Sesuvium</i></u> (2 spp.)	Aizoaceae Carpetweed Family

Plant List for Field Biology

Pine Flatwoods

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Beauty Berry	<u><i>Callicarpa americana</i></u>	Verbenaceae
Blueberry	<u><i>Vaccinia myrsinites</i></u>	Ericaceae Heath Family
Bracken Fern	<u><i>Pteridium aquilinum</i></u>	Pteridaceae
Brazilian Pepper	<u><i>Schinus terebinthifolius</i></u>	Bracken Fern Family Anacardiaceae (Introduced from S. America) Cashew Family
Cabbage Palm	<u><i>Sabal palmetto</i></u>	Aracaceae (Palmae)
Cabbage Palm Fern or Golden Polypody	<u><i>Phlebodium aureum</i></u>	Palm Family Polypodiaceae Polypody Fern Family
Fetterbush	<u><i>Lyonia lucida</i></u>	Ericaceae Heath Family
Gallberry	<u><i>Ilex glabra</i></u>	Aquifoliaceae Holly Family
Greenbrier	<u><i>Smilax (4 spp.)</i></u>	Smilacaceae Smilax Family
Heliotropes	<u><i>Heliotropium (3 spp.)</i></u>	Boraginiaceae
Lovevine	<u><i>Cassytha filiformis</i></u>	Lauraceae Laurel Family
Marlberry	<u><i>Ardesia escalloniodes</i></u>	Myrsinaceae Myrsine Family
Partridge Pea	<u><i>Cassia fasciculata</i></u>	Leguminosae (Fabaceae)
Penny Royal	<u><i>Satureja rigida</i></u>	Pea & Bean Family Lamiaceae
Poison Ivy	<u><i>Toxicodendron radicans</i></u>	Mint Family Anacardiaceae Cashew Family
Saw Palmetto	<u><i>Serenoa repens</i></u>	Palmae (Arecaceae) Palm Family
Slash Pine	<u><i>Pinus elliottii</i></u>	Pinaceae Pine Family
Staggerbush	<u><i>Lyonia ferruginea</i></u>	Ericaceae Heath Family
Swamp Fern	<u><i>Blechnum serrulatum</i></u>	Blechnaceae

**Plant List for Field Biology
Hammocks and Tree Islands**

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Air Plants or Bromeliads	<u><i>Tillandsia</i> (11 spp.)</u>	Bromeliaceae Pineapple Family
Beauty Berry	<u><i>Callicarpa americana</i></u>	Verbenaceae
Boston Fern	<u><i>Nephrolepis</i> (3 spp.)</u>	Davalliaceae (Intro Pantropical)
Brazilian Pepper	<u><i>Schinus terebinthifolius</i></u>	Anacardiaceae (Intro S. America) Cashew Family
Butterfly Orchid	<u><i>Encyclia tampensis</i></u>	Orchidaceae Orchid Family
Cabbage Palm	<u><i>Sabal palmetto</i></u>	Aracaceae (Palmae) Palm Family
Cabbage Palm Fern or Golden Polypody	<u><i>Phlebodium aureum</i></u>	Polypodiaceae Polypody Fern Family
Citrus	<u><i>Citrus</i> (7 spp.)</u>	Rutaceae (Intro S. E. Asia) Rue Family
Cocoplum	<u><i>Chrysobalanus icaco</i></u>	Chrysobalanaceae
Coral Bean	<u><i>Erythrina herbaceae</i></u>	Leguminosae (Fabaceae)
Florida Holly or Dahoon Holly	<u><i>Ilex glabra</i></u>	Bean & Pea Family Aquifoliaceae Holly Family
Greenbrier	<u><i>Smilax</i> (4 spp.)</u>	Smilacaceae Smilax Family
Guava	<u><i>Psidium guajava</i></u>	Myrtaceae (Intro Tropical America) Myrtle Family
Gumbo Limbo	<u><i>Bursera simaruba</i></u>	Burseraceae Torchwood Family
Mahogany	<u><i>Swietenia mahogoni</i></u>	Meliaceae (West Indian)
Mastic	<u><i>Mastichodendron foetidissimum</i></u>	Sapotaceae Sapodilla Family
Marlberry	<u><i>Ardesia escalloniodes</i></u>	Myrsinaceae Myrsine Family
Moon Flower or	<u><i>Ipomoea</i> (2 spp.)</u>	Convolvulaceae
Morning Glory	<u><i>Ipomoea</i> (14 spp.)</u>	Morning Glory Family Convolvulaceae (includes moonflower and railroad vine)
Myrsine	<u><i>Myrsine guianensis</i></u>	Myrsinaceae

Papaya	<u><i>Carica papaya</i></u>	Caricaceae Papaya Family
Passion Flower	<u><i>Passiflora (5 spp.)</i></u>	Passifloraceae Passionflower Family
Pigeon Plum	<u><i>Coccoloba diversifolia</i></u>	Polygonaceae Buckwheat Family
Poison Ivy	<u><i>Toxicodendron radicans</i></u>	Anacardiaceae Cashew Family
Poisonwood	<u><i>Metopium toxiferum</i></u>	Anacardiaceae Cashew Family
Red Bay and Swamp Bay	<u><i>Persea (s spp.)</i></u>	Lauraceae Laurel Family
Snowberry	<u><i>Chiococca alba</i></u>	Rubiaceae Madder Family
Spanish Moss	<u><i>Tillandsia usneoides</i></u>	Bromeliaceae Pineapple Family
Stopper	<u><i>Eugenia (5 spp.)</i></u>	Myrtaceae Myrtle Family
Strangler Fig	<u><i>Ficus aurea</i></u>	Moraceae Mulberry Family
Swamp Fern	<u><i>Blechnum serrulatum</i></u>	Blechnaceae Blechnum Fern Family
Virginia Creeper	<u><i>Parthenocissus quinquefolia</i></u>	Vitaceae Grape Family
Wild Coffee	<u><i>Psychotria (4 spp.)</i></u>	Rubiaceae Madder Family
Wild Grape	<u><i>Vitis (4 spp.)</i></u>	Vitaceae Grape Family
Wild Lime	<u><i>Xanthoxylem fagara</i></u>	Rutaceae

Plant List for Field Biology

Cultivated Plants

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Australian Pine	<u><i>Casuarina</i> (3 spp.)</u>	Casuarinaceae (Intro Australia) Beefwood Family
Banana	<u><i>Musa sapientum</i></u>	Musaceae
Brazilian Pepper	<u><i>Schinus terebinthifolius</i></u>	Anacardiaceae (Intro S. America) Cashew Family
Citrus	<u><i>Citrus</i> (7 spp.)</u>	Rutaceae (Intro S. E. Asia) Rue Family
Coconut Palm	<u><i>Cocos nucifera</i></u>	Aracaceae (Palmae) Palm Family
Life Plant	<u><i>Bryophyllum pinnatum</i></u>	Crassulaceae (Intro Africa) Orpine Family
Mango	<u><i>Mangifera indica</i></u>	Anacardiaceae (Intro Tropical Asia) Cashew Family
Mother-in-laws-tongue or Snake Plant	<u><i>Sanseveria thyrsiflora</i></u>	Agavaceae Agave Family (Intro S. Africa)
Oleander	<u><i>Nerium oleander</i></u>	Apocynaceae Oleander Family (Intro Mediterranean)
Periwinkle	<u><i>Catharanthus roseus</i></u>	Apocynaceae (Intro Africa) Oleander Family
Royal Palm	<u><i>Roystonea elata</i></u>	Arecaceae (Palmae) Palm Family
Sea Grape	<u><i>Coccoloba uvifera</i></u>	Polygonaceae Buckwheat Family

**Plant List for Field Biology
Disturbed Habitats and Ecotones**

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Australian Pine (Australia)	<u>Casuarina</u> (3 spp.)	Casuarinaceae Beefwood Family
Balsam Pear or Apple	<u>Momordica charantia</u>	Curcubitaceae Cucumber Family
Beard Grass	<u>Andropogon</u> (10 spp.)	Poaceae (Gramineae) Grass Family
Beggar's Ticks	<u>Desmodium</u> (8 spp.)	Fabaceae (Leguminosae) Pea Family
Bracken Fern	<u>Pteridium aquilinum</u>	Pteridaceae Bracken Fern Family
Brazilian Pepper	<u>Schinus</u> <u>terebinthifolius</u>	Anacardiaceae (S. America) Cashew Family
Caesar Weed	<u>Urena lobata</u>	Malvaceae Mallow Family
Castor Bean	<u>Ricinis communis</u>	Euphorbiaceae (Old World Tropics) Spurge Family
Crowsfoot Grass	<u>Dactyloctenium</u> <u>aegyptium</u>	Poaceae (Gramineae) (Eurasia) Grass Family
Dog Fennel	<u>Eupatorium</u> (2 spp. of dog plus 12 add. spp.)	Compositae Asteraceae
Foxtail Grass	<u>Setaria</u> (6 spp.)	Poaceae (Gramineae) Grass Family
Kalanchoe	<u>Kalanchoe grandiflora</u>	Crassulaceae (Old World Tropics) Orpine Family
Lantana	<u>Lantana</u> (5 spp.)	<u>Verbenaceae</u> Crassulaceae
Life Plant	<u>Bryophyllum</u> <u>pinnatum</u>	(Africa) Orpine Family
Lovevine	<u>Cassytha filiformis</u>	Lauraceae Laurel Family
Mother-in-laws tongue (Snake Plant)	<u>Sanseveria</u> <u>thyrsiflora</u>	Agavaceae Agave Family (S. Africa)

Paper Tree or Punkwood	<u>Melaleuca</u> <u>quinquinervia</u>	Myrtaceae Myrtle Family (Australia)
Periwinkle	<u>Catharanthus roseus</u>	Apocynaceae (Africa) Oleander Family
Prickly Pear Cactus	<u>Opuntia</u> (6 spp.)	Cactaceae Cactus Family
Puncture Weed	<u>Tribulus cistoides</u>	Zygophyllaceae
Ragweed	<u>Ambrosia</u> (2 spp.)	Compositae (Asteraceae) Aster Family
Rattlebox	<u>Crotalaria</u> (8 spp.)	Leguminosae (Fabaceae) Pea and Bean Family
Red Top Grass	<u>Rhynchelytrum</u> <u>repens</u>	Poaceae (Gramineae) (Africa) Grass Family
Rosary Pea or Crabs Eye	<u>Abrus precatorius</u>	Fabaceae (Leguminosae) Pea and Bean Family
Saltbush	<u>Baccharis</u> (4 spp.)	Compositae (Asteraceae) Aster Family
Sandspur or Sandbur	<u>Cenchrus</u> (7 spp.)	Poaceae (Gramineae) Grass Family
Spanish Needles	<u>Bidens pilosa</u>	Compositae (Asteraceae) Aster Family
Spurge	<u>Chaemaesyce</u> (18 spp.)	Euphorbiaceae (Euphorbia) Spurge Family
Wedelia	<u>Wedelia trilobata</u>	Asteraceae (Compositae) Aster Family
Wild Poinsettia	<u>Poinsettia</u> <u>cyathophora</u>	Euphorbiaceae (Euphorbia) Spurge Family

STUDY OF FISHES ON CORAL REEFS

PART 1. SWIMMING

SHAPE OF BODY IS AN INDICATOR OF LIFE STYLE. LOOK FOR AND IDENTIFY FISHES WITH THE FOLLOWING SHAPES:

1. **FUSIFORM AND TORPEDO**=NORMALLY ASSOCIATED WITH PREDATORS AND PLANKTON FEEDERS (MANAGING STRONG CURRENTS).
2. **SLENDER AND COMPRESSED**=ROWING MOTION WITH PECTORAL FINS;HOVERING AND TURNING.
3. **WINGS**=RAYS WHICH "FLY" THROUGH THE WATER.
4. **SIDE TO SIDE SWEEP**=WATCH FOR JAWS.

TYPES OF SWIMMING STYLES

1. **EEL-FORM** (TECHNICALLY ANGUILLIFORM)=UNDULATIONS OF WHOLE LENGTH OF THE BODY
2. **JACK FORM**=NOSE OF AN AIRCRAFT WITH TAPERED POSTERIOR ENDING IN A FORKED TAIL. ADAPTED FOR ACCELERATION.
3. **TRUNKFISH STYLE**=SCULLING EFFECT AND MOVEMENT OF TAILS
4. **BALISTIFORM STYLE**=UNDULATING DORSALS AND ANAL FINS

LOCOMOTION

Aquatic animals have distinctive shapes and swimming appendages for efficient locomotion in their environments.

PELAGIC FISH

Streamlined "torpedo-shaped" bodies with wide spread "V" or sickle shaped tails, and narrow streamlined fins are adaptations of the fast swimming fish of the open oceans.

REEF FISH

Fish which must be able to maneuver between the coral heads, rocks and plants of their homes usually have bodies flattened from side to side (laterally compressed), with short fan-shaped tails and fins for abrupt, stop and start, darting movements.

BOTTOM DWELLERS

For swimming, undulating or crawling along the bottom, these aquatic animals usually have bodies flattened from top to bottom (dorsal-ventrally depressed). These flat shapes are particularly helpful to animals living in tide pools. Why?

Can you determine where the fish live by their shapes? How do reef fish use their fins to hover in one spot? Look at the flat fish. It started life as a compressed fish swimming upright, but now lives on the bottom. How is it adapted?

STUDY OF FISHES ON CORAL REEFS

PART 2. EATING

DIVERS RARELY SEE FISH EATING. PART OF THE REASON IS THAT FISH EATING FISH EAT IN THE LOW LIGHT HOURS OF DAWN AND DUSK (CREPUSCULAR). LOOK FOR THESE FEEDING PATTERNS.

1. **GRAZERS**--MOST FISH SEEN DURING THE DAY ARE GRAZERS. WATCH PARROTFISHES, ANGELFISHES, DAMSELFISHES, BUTTERFLYFISHES, SURGEONFISHES, TRUNKFISHES, AND TRIGGERFISHES. THEY CRUISE AROUND THE SURFACE OF THE CORAL EATING ALGAE, MOLLUSKS, WORMS, AND SMALL CRUSTACEANS.
2. PARROTFISH TEETH ARE FUSED TOGETHER IN A CHISEL-LIKE BEAK. THEY BREAK OFF CHUNKS OF CORAL TO EXTRACT ALGAE. THEY SLIDE THE MATERIAL DOWN THEIR THROATS INTO A PHARYNGEAL MILL. THE HARD CORAL IS EXCRETED AS SAND.
3. **HUNTERS**--JACKS, SNAPPERS, AND SEA BASS HAVE LEAN, MUSCULAR BODIES FOR HUNTING AND EATING PREY. DURING THE DAY THOUGH, THEY MAY JUST **PATROL** AMONG POTENTIAL PREY (PARTICULARLY JACKS). WATCH FOR THIS BEHAVIOR. BARRACUDAS WILL MAINTAIN A CERTAIN **DISTANCE** AWAY FROM DIVERS. WATCH FOR THIS.

TRUMPETFISH OFTEN USE OTHER FISH AS SHIELDS OR THEY FLOAT IN A VERTICAL POSITION ALONGSIDE A GORGONIAN. WATCH FOR THIS.

GOATFISHES HAVE BARBELS WHICH THEY GLIDE ALONG THE SURFACE OF THE SAND TO FIND PREY BENEATH THE SURFACE.

TURN OVER A ROCK AND WATCH WHICH FISH EAT THE BRITTLE STARS AND ASSORTED INVERTEBRATES UNDER THE ROCKS.

FEED A VIENNA SAUSAGE TO A MORAY EEL.

4. **PLANKTON OPPORTUNISTS**--HERRINGS, SILVERSIDES AND YOUNG HALFBEAKS SEEN NEAR REEFS FEED ON PLANKTON.

SMALL WRASSE, DAMSELFISH, GRUNTS, SQUIRRELFISH, CARDINALFISH, AND BASS EAT PLANKTON INSTEAD OF HUNTING OR GRAZING. LOOK FOR FORKED TAILS AND LARGE EYES. SOME OF THESE FISHES FEED NOCTURNALLY.

STUDY OF FISHES ON CORAL REEFS

PART 3. COLOR

THE NEW DIVER SEES COLOR FIRST AND REMEMBERS IT THE LONGEST. WHY ARE REEF FISHES SO BRILLIANTLY COLORED? IN GENERAL, THEY MAY SERVE AS SOCIAL SIGNALS FOR **INTRA- AND INTERSPECIFIC SPECIES IDENTIFICATION**. OTHER PATTERNS ARE FOR CAMOUFLAGE AND MIMICRY, COURTSHIP DISPLAYS, PROTECTION AGAINST A BACKGROUND, AND AGGRESSIVE ENCOUNTERS. LOOK FOR THE FOLLOWING COLOR PATTERNS.

1 IDENTIFY SEVERAL FISHES BY THEIR **MORPHOLOGICAL COLOR STAGE**. THIS IS THE NORMAL ADULT PATTERN.

2 IDENTIFY SEVERAL **JUVENILE COLOR PATTERNS**. CONSULT YOUR FISH GUIDE FOR SOME JUVENILE COLOR PHASES. JUVENILES MAY HAVE DIFFERENT COLOR PATTERNS TO MASK THEIR SPECIES MEMBERSHIP.

3. IDENTIFY THE **SUPERMALES AND FEMALES** OF A PARTICULAR SPECIES. WRASSES AND PARROTFISHES ARE HERMAPHRODITIC FISHES WITH 5 OR 6 COLOR PATTERNS OVER THEIR LIFE.

4 FIND EXAMPLES OF **MIMICRY AND CAMOUFLAGE**. MANY NOVICE DIVERS SEE A BROWN, GREEN, BLUE, OR YELLOW TRUMPETFISH--ALL THE SAME SPECIES.

5 FIND EXAMPLES OF FISHES WITH **VERTICAL BARS**. THESE FISHES BLEND INTO THE BACKGROUND OF THE QUILT-LIKE PATTERN OF CORALS, VEGETATION, AND GORGONIANS ON THE REEF.

6 FIND AN EXAMPLE OF A FISH WITH ANOTHER EYE (**OCELLUS**). THESE MAY PROTECT A FISH FROM PREDATION, OR THEY MAY AID IN SPECIES RECOGNITION.

7 FIND AN EXAMPLE OF A FISH WITH STRIPES. THESE CAN BE FISH THAT ARE FAST SWIMMERS. "NOW YOU SEE ME, NOW YOU DON'T".

8. FIND A FISH WITH **ORNAMENTED EYES**. A MAJORITY OF THE FISHES ON THE REEF HAVE EYE RINGS, EYEBROWS, CONTRASTING BACKGROUNDS ETC..

PROTECTIVE COLOR AND SHAPE

To avoid being seen is an important part of survival for many aquatic animals.

HOW NOT TO BE SEEN

COUNTERSHADING

Pelagic fish which live in the open ocean are darker on the top (making them difficult to see from above against the darker water), and are lighter on the bottom (making them difficult to see from below against the lighter surface).

CAMOUFLAGE

Many marine animals blend in with their surroundings, both in shape and color, and some can even change their color, like the octopus and the flat fish.

DISRUPTIVE COLORATION

Many fish that live in the changing lights and shadows of the shallow waters or reefs have lines or spots which disguise their body shapes and hide the distinctive round shape of their vulnerable eyes. Some even have false eyespots on the rear portions of their bodies to fool predators into thinking they are coming when they are really going. Look for false eye spots on fish. How many stone fish can you find? Some fish like the Garibaldi are brightly colored with no disruptive markings. They do not try to camouflage themselves. Can you find out why?

BOTTOM DWELLERS

The bodies of bottom dwelling animals are usually flattened from top to bottom, and their colors often blend with the bottom, making them very hard to see. Many of the slow moving animals that live on the bottom have hard shells, like the lobster, crab and starfish or have sharp, like the sea urchin, to protect them. Some even have poisonous spines, like the stone fish and the stingray.

Many of the animals you will see in the tide pools do not have eyes, but use their tube feet or tentacles to feel for food.

STUDY OF FISHES ON CORAL REEFS

PART 4. SOCIAL LIFE

FIND THE FOLLOWING SOCIAL INTERACTIONS.

1 **SCHOOLING**. THIS IS THE MOST STRIKING EXAMPLE OF SOCIAL LIVING AMONG FISHES. WATCH THE DISTANCE AMONG THE FISHES. ARE THEY A BODY LENGTH APART? MORE OR LESS?

2 **RESTING AGGREGATIONS**. WATCH FOR GROUPS OF FISHES RESTING IN A PROTECTED CORAL SETTING. USUALLY, THESE ARE FISHES THAT FEED AT NIGHT.

3 FIND A **FISH COLONY**. A GOOD EXAMPLE IS A COLONY OF SERGEANT MAJORS OF ALL SIZES LIVING IN A PARTICULAR LOCATION.

4 LOOK FOR A FISH **PAIR**. THE NOVICE DIVER ALWAYS FINDS BUTTERFLYFISHES AND ANGELFISHES IN PAIRS. MOST OF THESE ARE HAPPILY MARRIED COUPLES.

5 FIND AND STUDY A **DAMSELFISH TERRITORY**. THESE SMALL FISH ARE AMONG THE BRAVEST ON THE REEF.

6 FIND A DAMSELFISH **ALGAE FARM**. MANY DAMSELFISH CULTIVATE AND PATROL A SMALL ALGAE CROP.

7 FIND A FISH WITH A **SAFETY HOLE**. MANY FISH LIVE CLOSE TO A SAFE AREA THEY CAN DIVE INTO WHEN A PREDATOR COMES NEARBY.

8 FIND A NEON GOBY OR JUVENILE FRENCH ANGELFISH **CLEANING STATION**. WATCH THE KINDS OF FISHES WAITING TO BE CLEANED.

9. FIND **SPONGE DWELLING GOBIES**. THESE FISHES FIND PROTECTION WITHIN VASE AND BASKET SPONGES.

BEHAVIORAL OBSERVATION

Pick a minimum of one and a maximum of three fish . Allot 25 minutes for your total observation time. Below are some questions to help you in your observations.

1) IDENTIFY YOUR FISH:

2) QUICK SKETCH:

3) IS YOUR FISH

SCHOOLING -

SOLITARY -

4) IS YOUR FISH EATING? DESCRIBE FEEDING BEHAVIOR. If your specimen is not feeding look at mouth position and form and give possible prey items. Does your fish eat small/big prey; hard/soft prey; scavenger, predator, grazer?

5) Does your fish exhibit any defense displays? What possible predators might your fish have and how might it avoid being eaten?

6) Describe the fish's coloration. With this in mind describe the fish's habitat

7) Do you notice any special appendages/fin or other adaptations? If so, briefly describe.

8) As you observe the fish, also note any interactions with other animals (fish or invertebrates).

SEASHELL KEY

1. Shell had two halves and you have only half of it (bivalve)
GO TO NUMBER 2
1. Shell is a snail shell and does not have two halves (gastropod)
GO TO NUMBER 11
2. The shell's hinge has a straight row of many tiny, evenly spaced "teeth"
GO TO NUMBER 3
2. The shell's hinge does not have "teeth", but may have points or scars where the hinges join; or the hinge may not be visible at all
GO TO NUMBER 4
3. The hinge is straight (and the shell has brown zebra-like stripes)
TURKEY WING
3. The hinge is not completely straight (shell is white) ARK SHELL
4. Part of the hinge seems to be rolled outward (and the shell is white with many ribs) ANGEL WING
4. No part of the hinge rolls out
GO TO NUMBER 5
5. Shell is much longer than it is high (check diagram for difference between length and width); or may have a wedge shape
GO TO NUMBER 6
5. Shell is not much longer than it is high - is it more round or oval
GO TO NUMBER 8
6. Ends of the shell are blunt; shell very rectangular in shape
GO TO NUMBER 7
6. Ends of the shell are rounded; shell is wedge-shaped; shell is no longer than one inch
FLORIDA COQUINA

7. Shell is more than four times as long as it is high (and looks like an old-fashioned razor) JACKKNIFE or RAZOR CLAM

7. Shell is less than four times the height (and the beak is almost in the center) TAGELUS

8. Shell has rays GO TO NUMBER 9

8. Shell does not have rays GO TO NUMBER 10

9. Hinge line has ears SCALLOP

9. Hinge line does not have ears COCKLE

10. Shell is fragile, no hinge is visible, and you can almost see through it JINGLE SHELL

10. Shell is not as above, but has concentric growth rings DOSINIA

11. Shell is fairly flat and dish-like GO TO NUMBER 12

11. Shell is not as above (more round in cross section) GO TO NUMBER 14

12. Shell is coiled (but only at one end) GO TO NUMBER 13

12. Shell is not coiled (and is shaped like a small, low cone) LIMPET

13. Shell has a deck or shelf inside SLIPPER or BOATHELL

13. Shell not as above (but has a row of several open holes) ABALONE

14. Shell has visible spire at top (it may be tiny) GO TO NUMBER 15
14. Shell is smooth and has no spire at all GO TO NUMBER 21
15. Aperture is roundish or oval in outline (it is less than twice as long as it is wide) GO TO NUMBER 16
15. Aperture is not round, but is at least twice as long as it is wide GO TO NUMBER 18
16. Shell is compressed and has a deep umbilicus (a large, funnel-shaped hole underneath) SUNDIAL
16. Shell not as above GO TO NUMBER 17
17. Shell is globular (ball-shaped) and has smooth surface MOON SNAIL
17. Shell is top-shaped and dark in color PERIWINKLE
18. Shell has very flattened spire (and is quite thin, with a criss-crossed surface pattern) FIG SHELL
18. Shell not as above GO TO NUMBER 19
19. Shell has knobs on spire (or where spire begins) GO TO NUMBER 20
19. Shell has no knobs on spire OLIVE SHELL
20. Aperture is much wider at the top than the bottom KNOBBED WHELK
20. Aperture is wide all the way down and has a notch at the bottom CONCH

21. Teeth are present inside aperture (and outer lip of aperture curls inward) COWRIE

21. Teeth are not present inside aperture GO TO NUMBER 22

22. Shell has a ridge running at right angles to aperture (and is creme-colored with orange edges)FLAMINGO TONGUE

22. Shell does not have a ridge across its back; but the apex (where the spire should be) has a small hole in it BUBBLE SHELL

Florida Field Trip References

1. Florida Aquatic Habitat and Fishery Resources. Seaman, William, Jr., ed.: 543 pp. Florida Chapter of the American Fisheries Society; 1985.

Good reference written in a non-technical way. Much useful information.

2. Coastal Plants of Florida. A Key to Good Land Management. Bullard, Lacy F., ed.: 38 pp. Florida Department of Agriculture and Consumer Services; 1979.

Useful.

3. Abbott, R. Tucker. Collectible Shells of Southeastern U.S., Bahamas & Caribbean: 64 pp. American Malacologists, Inc.; 1984.

4. Abbott, R. Tucker. A Guide to Field Identification. Seashells of North America: 280 pp. Golden Press; 1968.

Excellent guide.

5. Amos, William H. and Stephen H. Amos. Atlantic and Gulf Coasts. The Audubon Society Nature Guides: 670 pp. Alfred A. Knopf; 1985.

Useful information

6. Anderson, Robert. Guide to Florida Wading Birds: 54 pp. Winner Enterprises; 1985.

7. Austin, Sandra K. Common Plants of Palm Beach and Broward Counties.: 73 pp. Florida Atlantic University Foundation, Inc.

8. Bell, C. Ritchie and Bryan J. Taylor. Florida's Wild Flowers and Roadside Plants: 308 pp. Laurel Hill Press; 1982.

One you shouldn't leave home without.

9. Bergeron, Eugene. How to clean seashells: 32 pp. Great Outdoors Publishing Co.; 1971.

Offers information on the cleaning of a variety of shells.

10. Brayfield, Lelia and William. A Guide for Identifying Fossil Shells and Corals. Brayfield Research Lab; 1986.

Good black and white photographs.

11. Carr, Archie. The Sea Turtle; So Excellent a Fische: 280 pp. University of Texas Press, Austin; 1984.

Classic reference on sea turtles

12. Chaplin, C. G. Fishwatchers Guide to West Atlantic Coral Reefs: 65 pp. Harrowood Books; 1972.

Waterproof Edition which we use quite a bit. New edition is out, but we haven't purchased it.

13. Craig, Robert M. Plants for Coastal Dunes of the Gulf and South Atlantic Coasts and Puerto Rico: 41 pp. United States Department of Agriculture. Bulletin 460.; 1984.

14. Dittmar, Lucy. Caribbean Tropical Nature Guide: 64 pp. Lucy Dittmar; 1978.

Elementary

15. Duncan, Wilbur H. and Marion B. Duncan. The Smithsonian Guide to Seaside Plants of the Gulf and Atlantic Coasts: 409 pp. Smithsonian Institution Press; 1987.

Does not cover the lower peninsula of Florida. We have problems with just using scientific names beneath the colored photographs.

16. Dunkle, Sidney W. Dragonflies of the Florida Peninsula, Bermuda and the Bahamas: 154 pp. Scientific Publishers; The Sandhill Crane Press, Inc.; 1989.

Great photographs.

17. Fleming, Glenn, Pierre Genelle and Robert W. Long. Wild Flowers of Florida: 96 pp. Banyan Books, Inc.; 1984.

18. Fotheringham, Nick. Beachcomber's Guide to Gulf Coast Marine Life: 124 pp. Lone Star Books; 1980.

A new edition recently came out.

19. Gerberg, Eugene J., and Ross H. Arnetts, Jr. Florida Butterflies: 90 pp. Natural Science Publications, Inc.; 1989.

I like this book, but we have had trouble using it because you have to catch and kill the butterflies to identify them.

20. Greenberg, Idaz and Jerry. Guide to Coral and Fishes of Florida, the Bahamas and the Caribbean: 65 pp. Seahawk Press; 1977.

We use this one all the time. Waterproof.

21. Greenberg, Idaz, Jerry and Simba. Gamefishes of the Tropical Atlantic: Card. Seahawk Press; 1989.

One of a series of plastic cards.

22. Hall, Francis Wyly. Birds of Florida: 34 pp. Great Outdoors Publishing Co.; 1979.

Outdated

23. Hall, Francis Wyly. Shells of the Florida Coasts: 29 pp. Great Outdoors Publishing Co.; 1981.

24. Hargreaves, Dorothy and Bob. Tropical Trees: 64 pp. Hargreaves Company, Inc.; 1965.

25. Hoese, H. Dickson and Richard H. Moore. Fishes of the Gulf of Mexico, Texas, Louisiana and Adjacent Waters: 327 pp. Texas A & M University Press; 1977.

This book can fill in where the others leave off

26. Hoffmeister, John E. Land from the Sea. The Geologic Story of South Florida: 143 pp. University of Miami Press; 1974.

27. Hotchkiss, Neil. Underwater and Floating-Leaved Plants of the United States and Canada: 124 pp. Bureau of Sport Fisheries and Wildlife; Resource Pub. 44; 1967.

An old standby

28. Humann, Paul. Reef Fish Identification. Florida, Caribbean, Bahamas: 257 pp. New World Publications Inc.; 1989.

One of the best new identification guides. Offers a complete identification system.

29. Idaz, Jerry and Michael Greenberg. Fishwatcher's Field Guide: Card. Seahawk Press; 1979.

Plastic underwater fish card

30. Kale, II, Herbert W. and David S. Maehe. Florida's Birds; A Handbook and Reference: 288 pp. Pineapple Press; 1990.

Don't leave home without this one. Florida is a great place for bird watching.

31. Lazell, Jr., James D. Wildlife of the Florida Keys: 253 pp. Island Press; 1989.

A helpful addition to the field guides we have. Lazell provides an interesting account of the major species in the Keys.

32. Littler, Diane Scullion, Mark M. Littler, K. E. Bucher, and J. N. Norris. Marine Plants of the Caribbean: 263 pp. Smithsonian Institution Press; 1989.

Excellent photographs of major algae

33. Lockey, M.D., Richard and Lewis Maxwell. Florida's Poisonous Plants, Snakes, Insects: 79 pp. Lewis S. Maxwell, Publisher; 1978.

34. McGeachy, Beth. Handbook of Florida Palms: 63 pp. Great Outdoors Publishing Co.; 1955.

We have used this occasionally.

35. Myers, Ronald L. and John J. Ewel. Ecosystems of Florida: 765 pp. University of Central Florida Press; 1990.

Good reference for all major ecosystems including coral reef systems.

36. Perrero, Laurie. The World of Tropical Flowers: 64 pp. Windward Publishing, Inc.; 1976.

Good photographs. We like this one.

37. Pope, Patricia E. Seashore and Wading Birds of Florida: 44 pp. Great Outdoors Publishing Co.; 1974.

New Florida bird books are more comprehensive

38. Proctor, Lucille. Handbook of Florida Flowers: 48 pp. Great Outdoors Publishing Co.; 1959.

We've only used this one a few times.

39. Robins, C. Richard and G. Carleton Ray. *A Field Guide to Atlantic Coast Fishes of North America.*: 354 pp. Houghton Mifflin Company; 1986.

If you can't find it in any of the other books, turn to this. Poorly illustrated. Many line drawings.

40. Romashko, Sandra. *The Shell Book*: 64 pp. Windward Publishing, Inc.; 1984.

Colorful plates.

41. Romashko, Sandra. *The Complete Collector's Guide to Shells and Shelling*: 112 pp. Windward Publishing, Inc.; 1984.

Covers Pacific Shells as well.

42. Romashko, Sandra. *The Coral Book. A guide to collecting and identifying the corals of the world*: 64 pp. Windward Publishing, Inc. ; 1975.

Has several habitat photographs which are of assistance.

43. Schuetz, Maxine F. *Flowering Trees for Central and South Florida Gardens*: 142 pp. Great Outdoors Publishing Co.; 1990.

44. Siekman, Lula. *Handbook of Shells*: 48 pp. Great Outdoors Publishing Company; 1981.

45. Stevenson, George B. *Trees of the Everglades National Park and the Florida Keys*: 32 pp. Banyan Books, Inc.; 1988.

Short and sweet. We use this one.

46. Stokes, F. Joseph. Divers and Snorkelers Guide to the Fishes and Sea Life of the Carribean, Florida, Bahamas and Bermuda: 160 pp. The Academy of Natural Sciences of Philadelphia; 1984.

An excellent, indispensable reference.

47. Stresau, Frederic B. Florida, My Eden: 299 pp. Florida Classics Library; 1986.

Native and Exotic Plants with Photographs

48. Tarver, David P., John A. Rodgers, M. J. Mahler and R. L. Lazor. Aquatic and Wetland Plants of Florida: 127 pp. Florida Department of Natural Resources; 1986.

Excellent photographs; closeups and at a distance. We like this book.

49. Van Meter, Victoria Brook. Florida's Wood Storks : 26 pp. Florida Power and Light Company; 1985.

Much additional information is available from the Florida Power and Light Company.

50. Voss, Gilbert L. Seashore Life of Florida and the Caribbean: 199 pp. Banyan Books, Miami; 1976.

Great reference for seashore life excluding seashells

51. Voss, Gilbert L. Coral Reefs of Florida: 80 pp. Pineapple Press; 1988.

Good nontechnical account.

52. Williams, Joy. The 1990-1991 Edition. Florida Keys from Key Largo to Key West: 226 pp. Random House; 1989.

Some of the historical information is informative. Watch the directions to Watson's Hammock. They'll lead you into a mosquito nest.

53. Williams, Winston. Florida's Fabulous Waterbirds. Their Stories.: 64 pp. World-Wide Publishing; 1983.

We really like these books. Part of a series: Flowers, waterbirds, reptiles and amphibians, seashells.

54. Williams, Winston. Florida's Fabulous Trees. Their Stories.: 64 pp. World Publications; 1986.

Beautiful photographs.

55. Wilson, Roberta and James Q. Wilson. Watching Fishes. Life and Behavior on Coral Reefs.: 275 pp. Harper and Row, Publishers.; 1985.

A fun to read account with oodles of interesting information and references.