



## Examination of the Correlation between Dynamic Balance and Leg Strength of 11 and 12-Year-Old Children Who Have Fencing Training

 Meryem Gülaç<sup>1</sup>

<sup>1</sup>Dulmuşınar University, School of Physical Education and Sports, Kutahya, Turkey

Email: [meryem.gulac@dpu.edu.tr](mailto:meryem.gulac@dpu.edu.tr)



### Abstract

In this study, it is aimed to examine the relationship between leg strength and dynamic balance of children who have fencing training. 59 epee athletes who are 11 and 12 years old (33 11-year-old, 26 12-year-old) participated in research groups. Star Excursion Balance Test was used to determine balance performances and Baseline brand leg-dynamometer used to determine the leg strength of athletes who had fencing training. First, the Shapiro-Wilk test was used to determine if the data obtained during the study had a normal distribution. Pearson Correlation test was performed for all data displaying a normal distribution. Significance levels were taken as  $P < 0,05$  According to the research findings it was determined that there is a meaningful relationship between leg strength and balance test of athletes who have fencing training. As a result, it was seen that balance test values increase as the leg strength increases.

**Keywords:** Leg strength, Dynamic balance, Star excursion balance test, Epee, Children, Sports

**Citation** | Meryem Gülaç (2019). Examination of the Correlation between Dynamic Balance and Leg Strength of 11 and 12-Year-Old Children Who Have Fencing Training. Asian Journal of Education and Training, 5(1): 39-43.

**History:**

Received: 27 September 2018

Revised: 24 October 2018

Accepted: 29 November 2018

Published: 21 December 2018

**Licensed:** This work is licensed under a Creative Commons

Attribution 3.0 License 

**Publisher:** Asian Online Journal Publishing Group

**Funding:** This study received no specific financial support.

**Competing Interests:** The author declares that there are no conflicts of interests regarding the publication of this paper.

**Transparency:** The author confirms that the manuscript is an honest, accurate, and transparent account of the study was reported; that no vital features of the study have been omitted; and that any discrepancies from the study as planned have been explained.

**Ethical:** This study follows all ethical practices during writing.

### Contents

1. Introduction .....	40
2. Method .....	40
3. Findings .....	41
4. Discussion .....	42
5. Limitations and Suggestions .....	42
References .....	42

## 1. Introduction

Balance is an important factor in preserving the body composition which is necessary for successful performance in sports. Therefore, it is the basis for dynamic sports which include sudden changes in the pattern of motion especially. All sports contain a certain level of balance (Eler, 1996). Electronic systems (force platforms, stabilimeters, etc.) via which the movements of the body center of gravity projection are examined (Tsigilis *et al.*, 2001; Geldhof *et al.*, 2006) and functional dynamic balance tests such as Y Balance Test (Plisky *et al.*, 2009) Star Excursion Balance Test (Filipa *et al.*, 2010) are used in balance assessment. Since the use of new system measuring equipment is limited in reflecting the balance performance encountered during sports movements, using the functional balance tests in the measurements is also worthwhile. Star excursion balance test is a test that is commonly used in the literature for dynamic balance and postural control and that has a high fundamental reliability (Hertel *et al.*, 2000; Gribble *et al.*, 2013).

Leg strength is important in fencing branch. For a successful score, factors such as a balanced en garde position, flexibility and leg length are effective rather than lengthening to the opponent with a weapon only (Lavoie *et al.*, 1985). A fencer who finishes his/her attack needs to reach maximum speed at the moment to make his/her move and needs to reach the target with his/her weapon as soon as possible. For this reason, having a good strength is also very important for fencers as well as other features (Gioux *et al.*, 1984). The fatigue that may occur late in the running battle during a fencing game may obstruct to maintain balance especially.

Leg strength is a subject that has been studied frequently in fencing sport (Ntai *et al.*, 2017). Within the scope of studies with leg strength, studies towards the effect of applied training programs, dominant and non-dominant leg strength comparison, differences between elite and non-elite fencers have been carried out (Tsolakis *et al.*, 2010; Tsolakis *et al.*, 2010; Tsolakis and Vagenas, 2010; Tsolakis *et al.*, 2011). Several studies have also been conducted to examine the relationship between lower extremity muscle strength and balance performance. These studies have been applied to different branches (Soyuer and Mirza, 2006; Mohammadi *et al.*, 2012; Celenk *et al.*, 2015; Siriphorn and Chamonchant, 2015; Aktuğ, 2016). Within the scope of studies related to fencing, anthropometric characteristics, variables such as age, sports age have been examined more. We carried out our study to investigate if there is a relationship between balance and leg strength, the two characteristics that are important for fencers.

## 2. Method

### 2.1. Participants

Measurements have been taken from 59 11 and 12-year-old epee athletes (33 11-year-old, 26 12-year-old) who have joined the Turkey Super Juniors and Juniors Championships held in the province of Eskisehir in 2017. The athletes, who have been receiving fencing training at least for 1 year and working out at least 3 days a week, have participated in this study in line with the information obtained from their coaches. After initially contacting the Federation, permissions have been taken from Eskişehir provincial representative, who made the organization, and coaches of the participants and then the athletes have joined the measurements voluntarily. During the measurement, the hand of a participant that he/she actively uses has been preferred.

### 2.2. Measurement Tool

*Dynamic balance measurement:* Star excursion balance test, which interests lower extremity and measures the dynamic balance of participants, has been used in this study. The test is carried out in eight directions. The test can be used to measure clockwise and anti-clockwise or both sides (Reiman and Manske, 2009). The outmost point that an athlete maintains his/her position for three seconds is recorded in centimeters (Lockie *et al.*, 2013).


*Measurement Of Leg Strength:* The measurement was made using Baseline brand leg-dynamometer. Following five minutes of warm-up, after placing their feet on the dynamometer table with their knees bent, the subjects have hove the dynamometer bar, which they have grabbed with their hands, vertically at a maximum using their legs while their arms were stiff, backs were straight and torsos were leaned forward slightly. This heaving has been repeated three times and the best value has been recorded for every subject.

### 2.3. Application

Athletes have been summoned one by one and taken to the private area prepared for measurement. First, their demographic information was recorded. Following that, star excursion balance test (with both feet) and strength measurement have been carried out respectively. For the star excursion balance test, the athletes have waited unshod in the starting position at the exact center of the star which had been previously fixed to the ground. They have reached out to the outmost point they could with the 45-degree angle for the 8 points which were pre-determined. The proper trial has been regarded as the participant touching the outmost point with a reach out move without unbalancing and coming back to the starting point without unbalancing again. The participant's getting support from the ground during touching or touching a different point with loss of his/her balance has not been considered as a proper trial and the trial has been repeated. For the assessment of one participant, a total of 16 scores, both right and left scores for 8 points, have been recorded considering the average of 3 tries for each direction.

### 2.4. Analysis

Statistical analysis was performed via SPSS 22 program. Shapiro-Wilk test has been used to determine if the data had a normal distribution. The relationship between leg strength and balance performance values has been analyzed using Pearson for the data that is deemed to have a normal distribution in the test result. Significance level has been taken as  $p < 0,05$



### 3. Findings

Table-1. Descriptive Statistics

		Descriptive Statistics		
			Mean	Std. Deviation
STAR EXCURSION BALANCE TEST	RIGHT	anterior	26,56	16,375
		anterolateral	52,41	20,383
		lateral	44,98	15,206
		posterolateral	51,93	16,211
		posterior	121,37	24,681
		posteromedial	65,81	11,751
		anteromedial	66,17	18,640
		medial	59,37	22,995
	LEFT	anterior	15,25	15,360
		anteromedial	67,66	13,811
		medial	68,05	20,077
		posteromedial	71,66	17,670
		posterior	123,58	24,146
		posterolateral	54,83	18,302
	lateral	50,83	12,248	
	anterolateral	46,14	15,964	
<b>THE LEG POWER</b>			52,43	18,793

Source: Author's field work

Table-2. The Relationship Between Star Excursion Balance Test and Leg Strength

Leg Strength		
Right Leg Star Excursion Balance	Pearson Correlation	Sig. (2-tailed)
Anterior	,290	,026
Anterolateral	,409	,001
Lateral	,369	,004
Posterolateral	,353	,006
Posterior	,428	,001
Posteromedial	,683	,000
Anteromedial	,484	,000
Medial	,518	,000
Left Leg Star Excursion Balance		
Anterior	-,023	,863
Anteromedial	,599	,000
Medial	,649	,000
Posteromedial	,626	,000
Posterior	,482	,000
Posterolateral	,454	,000
Lateral	,636	,000
Anterolateral	,423	,001

Source: Calculated from primary data \*Level of significance at p<0.05

According to the leg strength and star excursion balance test results, despite the non-existence of a significant relationship in the direction of Anterior Left Foot, it has been founded that there is a significant relationship between star excursion balance test and leg strength in all other parts.

## 4. Discussion

A sudden and severe forward move is the most characteristic leg move for a fencer. Because each attack develops on the basis of this movement. A forward move made fast and timely is regarded as the core principle of being a successful fencer and this sport (Weineck, 1998). Fencing is a sport which also improves balance by the nature of posture and footwork. Fencers move like they are walking on a thin line and they have to maintain their balances either at high speed or sudden deflections not to be eliminated from the game (Gökmen *et al.*, 1995). Within the scope of our study, it has been noted that there is a positive correlation between balance and leg strength characteristics and balance values increase as the leg strength increases. Studies in different branches seem to support our work. Soyuer and Mirza (2006) have found in their study that there is a relationship between balance and lower extremity muscle strength. Siriphorn and Chamonchant (2015) have stated in their study (2015) that, as a result of the Wii balance exercise that they had applied to the weight-lifters, the balance skill has been improved and the strength of lower extremity muscles has been increased. Mohammadi *et al.* (2012) have found in their study on young male athletes that the leg strength has increased and improvements on dynamic and static balance have occurred following a 6-week strength training toward leg muscles. Soyuer and Mirza (2006) have revealed in their study on subjects with Multiple Sclerosis that a relationship between balance and lower extremity muscle strength is existent. Bressel *et al.* have mentioned the existence of a relationship between dynamic balance and performance in their study, which was carried out in 2007 (Bressel *et al.*, 2007). Muscles transmit strength with the strength they produce by reaching out to adjacent body parts. These functions of muscles and joints are the basic mechanism that allows the emergence of the movement. A good posture and stability require the coordination of joint positions and movements. Different joints and different muscles, correspondingly, are used to maintain the balance and posture required for the work done (Kunduracioğlu, 1999). Celenk *et al.* (2015). They have stated that there is a positive correlation between Quadriceps muscle strength and balance performance. Mohammadi *et al.* (2012) have stated that a 6-week strength program has increased leg strength in young athletes, this situation has generated improvements on dynamic and static balance performance. Young *et al.* (2010) have stated that the relationship between lower extremity muscle strength and balance performance might be arising from the emergence of muscle coordination and the increase in motor unit contraction speed. Again, Liman (2008) has explained the positive correlation between strength and balance with the development of inner-muscle and intra-muscles coordination by the increase in muscle strength and this situation increasing synergist and antagonist working capacities of extensor and flexor muscles.

In some of the studies, a meaningful relationship has not been found. In Kurt's study (2015) that was performed with 30 male freestyle wrestlers, who are selected to the national team, a statistically significant difference between leg strengths of weights has been determined where a statistically significant difference between balance results has not been determined in the scope of Freestyle Wrestling. Kapşigay *et al.* (2013) couldn't have detected a statistically significant difference between dominant and non-dominant lower extremity body balance in their study that they have performed on footballers.

As a result, in our study, which was conducted with fencers, it was found that there is a positive correlation between leg strength and balance. It has been revealed that the balance characteristic also improves as the leg strength increases. Looking at the results of studies, that have been conducted on different disciplines, it has been seen that there is a necessity of strength exercises to be included in training programs (Manolopoulos *et al.*, 2004; Asadi *et al.*, 2016). Under the circumstances, the inclusion of specific strength exercises in training programs along with technical tactical practices is recommended to be a good fencer.

## 5. Limitations and Suggestions

It is thought that there are not enough studies on fencing branch and that the studies related to this field should be increased. The number of samples can be increased.

## References

- Aktuğ, Z.B., 2016. Investigating the relationship between isokinetic leg strength and balance performance LEG volume and leg mass in professional football players. Erciyes University, Health Sciences Institute, Department of Physical Education and Sports, PhD Thesis, Kayseri.
- Asadi, A., H. Arazi, W.B. Young and E.S. De Villarreal, 2016. The effects of plyometric training on change-of-direction ability: A meta-analysis. *International Journal of Sports Physiology and Performance*, 11(5): 563-573. Available at: <https://doi.org/10.1123/ijsp.2015-0694>.
- Bressel, E., J. Yonker, J. Kras and E. Heath, 2007. Comparison of static and dynamic balance in female collegiate soccer, basketball, and gymnastics athletes. *Journal of Athletic Training*, 42(1): 42-46.
- Celenk, Ç., İ. Marangoz, Z.B. Aktuğ, E. Top and M. Akil, 2015. The effect of quadriceps femoris and hamstring muscular force on static and dynamic balance performance. *International Journal of Physical Education Sports and Health*, 2(2): 323-325.
- Eler, S., 1996. The examination of some motoric and physiological parameters of elite class handball players throughout one season training period. Gazi University, Health Sciences Institute, Department of Physical Education and Sports, Master Thesis, Ankara.
- Filipa, A., R. Byrnes, M.V. Paterno, G.D. Myer and T.E. Hewett, 2010. Neuromuscular training improves performance on the star excursion balance test in young female athletes. *Journal of Orthopaedic & Sports Physical Therapy*, 40(9): 551-558. Available at: <https://doi.org/10.2519/jospt.2010.3325>.
- Geldhof, E., G. Cardon, I. De Bourdeaudhuij, L. Danneels, P. Coorevits, G. Vanderstraeten and D. De Clercq, 2006. Static and dynamic standing balance: Test-retest reliability and reference values in 9 to 10 year old children. *European Journal of Pediatrics*, 165(11): 779-786. Available at: <https://doi.org/10.1007/s00431-006-0173-5>.
- Gioux, M., P. Ame and M.V.A. Dogui, 1984. Biomechanical and electrogoniographical characteristics of the human quadriceps in relation with sport performance. *Proceedings of the World Congress of Sports Medicine, Urban and Schwarzenberg*, pp: 699-705.
- Gökmen, H., T. Karagül and F.H. Aşçı, 1995. Psychomotor development. Ankara: General Directorate of Youth and Sports Publications, 14.
- Gribble, P.A., S.E. Kelly, K.M. Refshauge and C.E. Hiller, 2013. Interrater reliability of the star excursion balance test. *Journal of Athletic Training*, 48(5): 621-626. Available at: <https://doi.org/10.4085/1062-6050-48.3.03>.
- Hertel, J., S.J. Miller and C.R. Denegar, 2000. Intratester and intertester reliability during the star excursion balance tests. *Journal of Sport Rehabilitation*, 9(2): 104-116. Available at: <https://doi.org/10.1123/jsr.9.2.104>.
- Kapşigay, B., B. Özgül, Z. Sari and M.G. Polat, 2013. Investigation of the effect of the dominant and non-dominant lower limb on body balance in football players. 7th Congress of national sports Physiotherapists, 7-9 November, Ankara.

- Kunduracioğlu, B., 1999. Stabilometric evaluation of lower extremity proprioception before and after cycling and running exercises. Ankara University Faculty of Medicine, Department of Sports Medicine, Master Thesis, Ankara.
- Lavoie, J.M., L. Leger, R. Pitre and J.F. Etmarin, 1985. Fencing competition Epee. Analysis of travel times and distances. *Sports Medicine*, 59(5): 279-283.
- Liman, Ö.N., 2008. The effects of aerobic-step and pilates exercise on strength, flexibility, anaerobic power, balance and body composition. Gazi University Health Sciences Institute, Department of Physical Education and Sports, Master Thesis, Ankara.
- Lockie, R.G., A.B. Schultz, T.M. Luczo, S.J. Callaghan and M.D. Jeffriess, 2013. Effect of unilateral dynamic stability on lateral jump performance in team sport athletes. *Serbian Journal of Sports Sciences*, 7(4): 159-166.
- Manolopoulos, E., C. Papadopoulos, K. Salonikidis, E. Katartzi and S. Poluha, 2004. Strength training effects on physical conditioning and instep kick kinematics in young amateur soccer players during preseason. *Percept Mot Skills*, 99(2): 701-710. Available at: <https://doi.org/10.2466/pms.99.2.701-710>.
- Mohammadi, V., M. Alizadeh and A. Gaieni, 2012. The effects of six weeks strength exercises on static and dynamic balance of young male athletes. *Procedia-Social and Behavioral Sciences*, 31: 247-250. Available at: <https://doi.org/10.1016/j.sbspro.2011.12.050>.
- Ntai, A., F. Zahou, G. Paradisis, A. Smirniotou and C. Tsolakis, 2017. Anthropometric parameters and leg power performance in fencing. Age, sex and discipline related differences. *Science & Sports*, 32(3): 135-143. Available at: <https://doi.org/10.1016/j.scispo.2016.06.011>.
- Plisky, P., P. Gorman, R. Butler, K. Kiesel, F. Underwood and B. Elkins, 2009. The reliability of an instrumented device for measuring components of the star excursion balance test. *North American Journal of Sports Physical Therapy*, 4(2): 92-99.
- Reiman, M.P. and R.C. Manske, 2009. *Functional testing in human performance: 139 tests for sports, fitness, and occupational settings*. Champaign, Illinois: Human Kinetics Books.
- Siriphorn, A. and D. Chamonchant, 2015. Wii balance board exercise improves balance and lower limb muscle strength of overweight young adults. *Journal of Physical Therapy Science*, 27(1): 41-46. Available at: <https://doi.org/10.1589/jpts.27.41>.
- Soyuer, F. and M. Mirza, 2006. Relationship between lower extremity muscle strength and balance in multiple sclerosis. *Journal of Neurological Sciences*, 23(4): 257-263.
- Tsigilis, N., E. Zachopoulou and T. Mavridis, 2001. Evaluation of the specificity of selected dynamic balance tests. *Perceptual and Motor Skills*, 92(3): 827-833. Available at: <https://doi.org/10.2466/pms.2001.92.3.827>.
- Tsolakis, C., G.C. Bogdanis, A. Nikolaou and E. Zacharogiannis, 2011. Influence of type of muscle contraction and gender on postactivation potentiation of upper and lower limb explosive performance in elite fencers. *Journal of Sports Science and Medicine*, 10(3): 577-583.
- Tsolakis, C., A. Douvis, G. Tsigganos, E. Zacharogiannis and A. Smirniotou, 2010. Acute effects of stretching on flexibility, power and sport specific performance in fencers. *Journal of Human Kinetics*, 26: 105-114. Available at: <https://doi.org/10.2478/v10078-010-0054-x>.
- Tsolakis, C., E. Kostaki and G. Vagenas, 2010. Anthropometric, flexibility, strength-power, and sport-specific correlates in elite fencing. *Perceptual and Motor Skills*, 110(3C): 1015-1028. Available at: <https://doi.org/10.2466/pms.110.3c.1015-1028>.
- Tsolakis, C. and G. Vagenas, 2010. Anthropometric, physiological and performance characteristics of elite and sub-elite fencers. *Journal of Human Kinetics*, 23(1): 89-95. Available at: <https://doi.org/10.2478/v10078-010-0011-8>.
- Weineck, J., 1998. *Sports anatomy*. Ankara: Bağrgan Publishing House. pp: 225- 227.
- Young, M.D., D. Jordan and M.A.Y. Metz, 2010. Strength training for the young athletes. *Medial Pediatric Annals*, 39(5): 293-299. Available at: [10.3928/00904481-20100422-10](https://doi.org/10.3928/00904481-20100422-10).