

## DOCUMENT RESUME

ED 313 596

CG 022 114

AUTHOR Starzyk, Patricia M.  
 TITLE Alternative Maternity Services in Washington State.  
 PUB DATE 17 Nov 88  
 NOTE 12p.; Paper presented at the Annual Meeting of the American Public Health Association (116th, Boston, MA, November 13-17, 1988).  
 PUB TYPE Reports - Research/Technical (143) -- Speeches/Conference Papers (150)  
 EDRS PRICE MF01/PC01 Plus Postage.  
 DESCRIPTORS \*Birth; \*Health Services; \*Obstetrics; \*Pregnancy; \*Social Services  
 IDENTIFIERS \*Washington

## ABSTRACT

The nature of maternity services has changed in the past 20 years, with a movement away from traditional (physician delivery in a hospital) towards other alternative services. This study examined alternative maternity services in Washington State, which ranks eighth in the country in the use of such services. Data were collected from birth and infant and fetal death certificates. A delivery was considered alternative if it occurred outside of a hospital and/or was delivered by a non-physician attendant, typically a midwife. The findings revealed that alternative maternity services in Washington were used by a select group of low-risk women with generally favorable pregnancy outcomes. Trend data showed that more low-risk women were choosing these services every year. This increase, coupled with the fact that 45% of Washington women having babies are in this low-risk category and could thus choose alternative services, suggests that any consideration of new delivery technologies must take into account the desire for "low-tech" alternatives. (NB)

\*\*\*\*\*  
 \* Reproductions supplied by EDRS are the best that can be made \*  
 \* from the original document. \*  
 \*\*\*\*\*

ED313596

To be presented: 9:35 AM Thursday November 17, 1988  
Session title: Trends in Traditional and Alternative Approaches  
to Maternity Services and Care  
Primary sponsor: Maternal and Child Health  
Presented at: American Public Health Association Annual Meeting,  
Boston, Mass.

ALTERNATIVE MATERNITY SERVICES IN WASHINGTON STATE

(Patricia M. Starzyk, PhD, Washington State Department of Social  
and Health Services, Health Data Unit, Olympia, Washington)

INTRODUCTION

The nature of maternity services has changed in the past two  
decades, with a movement away from the traditional (physician  
delivery in a hospital) towards other, alternative, services (1).  
As a result, there is increased interest in these services, both  
by groups wanting to make them more widely available and by  
groups wanting tighter regulation of obstetric practice. This  
increased interest has created a need to put alternative  
maternity services into perspective and define what their role  
should be, as a guide to public health policy decisions.  
Accordingly, this paper presents data on alternative maternity  
services in Washington State, which ranks eighth in the country  
in the use of these services (1). Three areas are covered:  
trends in service use, characteristics of service users, and  
pregnancy outcomes for service users.

CG 022114

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

- This document has been reproduced as  
received from the person or organization  
originating it.
- Minor changes have been made to improve  
reproduction quality.

• Points of view or opinions stated in this docu-  
ment do not necessarily represent official  
• OERI position or policy.

"PERMISSION TO REPRODUCE THIS  
MATERIAL HAS BEEN GRANTED BY

Patricia M.  
Starzyk

TO THE EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC)."



The results presented here are based on data reported on birth and infant and fetal death certificates. A delivery was considered alternative if it occurred outside of a hospital and/or was delivered by a non-physician attendant, typically a midwife. As used in this paper, 'midwife' refers to both certified nurse midwives and licensed midwives. Certified nurse midwives account for about 80 percent of the midwife deliveries. They practice mainly in hospitals, whereas licensed midwife deliveries occur primarily at home. Lay midwives, who deliver about 100 babies annually, are not included in this midwife classification.

## RESULTS

1. Trends in service use: Between 1975 and 1985, the use of traditional maternity services declined from 99 percent of births to 95 percent, representing a decrease of about 2,700 deliveries per year. Two separate changes have been responsible for this decline (Figure 1). At first, the decline was due to an increase in out-of-hospital deliveries (home, freestanding birthing center). However, out-of-hospital delivery reached its peak in 1981 and has since declined in popularity, largely due to a decrease in deliveries in freestanding birthing centers. Thus, women are returning to hospital delivery, but not to traditional services. All of the recent increase in hospital delivery is due to hospital deliveries by midwives, which had a twelve-fold increase between 1979 and 1985 (Figure 2).

How do these trends fit together? Initially, women seeking 'different' maternity services found them outside of the hospital, in birthing centers or their own homes. More recently, changes in hospital-based obstetrics (setting up birthing rooms, allowing more midwifery practice) have allowed these women to return to the hospitals, deserting birthing centers, but not home births, which remain relatively constant at 1.5 percent of births. However, even the nature of home delivery has changed with the increase in midwifery practice. Decreasing physician home deliveries left a gap which at first was filled by unlicensed attendants (fathers, lay midwives). In fact, unlicensed attendants were the most common home delivery attendant for the years 1977-82 (Figure 3). However, recent increases in midwife home deliveries have changed this pattern so that midwife deliveries now outnumber unlicensed attendant deliveries by 2:1.

2. Characteristics of service users: Trend data have shown that midwives are the key to many of the recent changes in alternative service use. What type of woman uses these midwife services? Birth certificate data show that Washington State women who have midwife deliveries tend to be older (non-teen), white, married women (Table 1). Compared to women using traditional services, they have timely prenatal care more often and are less likely to smoke during pregnancy, and they are also less likely to have had a previous delivery with adverse outcome (late fetal loss, premature infant, or previous live birth now dead).

The salient point emerging from this analysis is that these midwife users are women who could have chosen a traditional delivery if they had wished. Thus, they probably used alternative services not because other services were not available, but because a personal choice. This distinction has consequences in terms of pregnancy outcomes to be expected.

3. Pregnancy outcomes: The most important consideration in any analysis is still the outcome - how did the baby do? Since women having midwife deliveries are in general the low-risk mothers, their outcomes should be better than average. In this case, the issue becomes: how do these women compare to similarly low-risk women who use traditional services? Even among these low-risk mothers, infants delivered by midwives have low birth weight considerably less often (Table 2). Perinatal mortality rates are also significantly lower [ $p(X^2) < .001$ ] for babies delivered by midwives (even adjusting for birth weight differences). In contrast, postneonatal mortality rates, which depend more on environmental factors than on health care, have little variation by type of attendant.

## DISCUSSION

These results indicate that, at least in Washington State, alternative maternity services are used by a select group of low-risk women with generally favorable pregnancy outcomes. Furthermore, trend data show that more low risk women are choosing these services every year. This increase, coupled with the fact that 45 percent of the Washington State women having babies are in this low-risk category and could thus choose alternative services, suggests that any consideration of new delivery technologies must take into account the desire for 'low-tech' alternatives.

These changes also have consequences for traditional maternity services. As more low-risk women choose alternative services, physicians are left to deal with relatively more high-risk deliveries. In this respect, Washington State is moving closer to the European experience, where trained professional midwives deliver the majority of babies and physicians primarily handle problem deliveries, in keeping with their training and talents (2). Unfortunately, these changes come at a time when more obstetricians are curtailing delivery services because of malpractice insurance increases (3). Whereas low-risk pregnancies can be accommodated by alternative services, creative solutions will still need to be found for dealing with these high-risk pregnancies.

## REFERENCES

1. Vital Statistics of the United States, Volume I-Natality.  
US DHHS, Public Health Service, Hyattsville, Md, published  
annually.
2. Midwifery Outside the Nursing Profession: The Current Debate  
in Washington. Health Policy Analysis Program, University of  
Washington School of Public Health and Community Medicine,  
Seattle, Wa, October, 1980.
3. Reported in The Nation's Health. American Public Health  
Association, Washington, DC, April, 1988.

FIGURE 1.

### TRENDS IN OUT-OF-HOSPITAL DELIVERY WASHINGTON STATE RESIDENTS, 1975-86






FIGURE 2.

### TRENDS IN HOSPITAL DELIVERY BY MIDWIVES WASHINGTON STATE RESIDENTS, 1975-86




FIGURE 3.

### TRENDS IN HOME DELIVERY ATTENDANT WASHINGTON STATE RESIDENTS, 1975-86





TABLE 1.

**PROFILES OF WOMEN WITH HIGH AND LOW MIDWIFE USE  
WASHINGTON STATE RESIDENTS, 1985**



•AGE-ADJUSTED PERCENTS

TABLE 2.

**BIRTH OUTCOMES FOR LOW-RISK WOMEN  
WASHINGTON STATE RESIDENTS, 1984-85**

|  | <b>MD</b>  | <b>MIDWIFE</b> |
|--|------------|----------------|
| <b>PERCENT LOW BIRTH WEIGHT</b>              | <b>3.2</b> | <b>0.8</b>     |
| <b>DEATH RATES (per 1,000 live births)**</b> |            |                |
| <b>PERINATAL</b>                             | <b>9.2</b> | <b>3.5</b>     |
| <b>POSTNEONATAL</b>                          | <b>3.4</b> | <b>3.8</b>     |

**•LOW-RISK - WHITE SINGLETON INFANT BORN TO MARRIED MOTHER  
AGED 20-39 WHO BEGAN PRENATAL CARE IN FIRST TRIMESTER**

**\*\*ADJUSTED FOR BIRTH WEIGHT DIFFERENCES**