

DOCUMENT RESUME

ED 409 347

TM 026 830

AUTHOR Triska, Olive H.; And Others
TITLE Clinicians' Explanations of Students' Reasoning while Solving Multiple-Choice Items.
PUB DATE 26 Mar 97
NOTE 22p.; Paper presented at the Annual Meeting of the American Educational Research Association (Chicago, IL, March 24-28, 1997).
PUB TYPE Reports - Research (143) -- Speeches/Meeting Papers (150)
EDRS PRICE MF01/PC01 Plus Postage.
DESCRIPTORS *Distractors (Tests); Higher Education; *Medical Education; Multiple Choice Tests; *Physicians; *Problem Solving; Protocol Analysis; *Student Attitudes; Test Construction; Thinking Skills

ABSTRACT

A study was conducted to determine whether competently reasoning clinicians (clinical instructors in medical instruction) could identify reasons competently reasoning students would eliminate distractors, and explain how students would reason to select the keyed response when solving multiple-choice items. The think-aloud protocols of clinicians (the number varying from 5 to 11) for 4 multiple-choice items were analyzed by categorizing and coding the clinicians' explanations and then comparing them to the justifications of students (varying from 18 to 32). Results show that expert clinicians were more successful in identifying the students' justifications for eliminating distractors than were novice clinicians, and novice clinicians were more successful than expert clinicians in explaining how students reasoned to select the keyed response. These results suggest that the expert and novice clinicians' decisions complemented each other in identifying the distractor elimination and perceptions of how students solved items. (Contains eight tables and eight references.) (Author/SLD)

* Reproductions supplied by EDRS are the best that can be made *
* from the original document. *

**CLINICIANS' EXPLANATIONS OF
STUDENTS' REASONING WHILE SOLVING
MULTIPLE-CHOICE ITEMS**

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

This document has been reproduced as
received from the person or organization
originating it.

Minor changes have been made to
improve reproduction quality.

• Points of view or opinions stated in this
document do not necessarily represent
official OERI position or policy.

PERMISSION TO REPRODUCE AND
DISSEMINATE THIS MATERIAL
HAS BEEN GRANTED BY

Olive H. Triska

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC)

Olive H. Triska*,
Faculty of Education & Faculty of Medicine, University of Alberta, Edmonton Alberta, Canada

Thomas O. Maguire,
Faculty of Education & Faculty of Medicine, University of Alberta, Edmonton Alberta, Canada

Charles C. Harley
Faculty of Medicine, University of Alberta, Edmonton Alberta, Canada

**Presented at the Annual Meeting of the
American Educational Researchers Association**

**Chicago, Illinois
Wednesday, March 26, 1997.**

*

This research was supported in part by the National Health Research and Development Program, Canada, through a National Health Fellowship to Olive H. Triska.

TM026830

Clinicians' Explanations of Student Reasoning While Solving Multiple-choice Items

O.H. Triska¹, T.O. Maguire^{1,2}, C.C. Harley².

Faculty of Education¹, and Faculty of Medicine², University of Alberta, Edmonton Alberta, CA

Abstract

The purpose of this study was to determine if competently reasoning clinicians could: (1) identify reasons competently reasoning students used to eliminate distractors; and, (2) explain how students would reason to select the keyed response when solving multiple-choice items. Clinicians (n=5 to 11) think aloud protocols for four multiple-choice items were analyzed by categorizing and coding the clinicians' explanations. and compared to students' (n=18 to 32) justifications.

The results showed that: (1) expert clinicians were more successful in identifying the students' justifications for eliminating distractors than novice clinicians; and, (2) novice clinicians were more successful than expert clinicians in explaining how students reasoned to select the keyed response. These results suggest that the expert and novice clinicians' decisions complimented each other in identifying the distractor elimination and perceptions of how students solved items.

CLINICIANS' EXPLANATIONS OF STUDENT REASONING WHILE SOLVING MULTIPLE-CHOICE ITEMS

O.H. Triska¹, T.O. Maguire^{1,2}, C.C. Harley².

Faculty of Education¹, and Faculty of Medicine², University of Alberta, Edmonton Alberta, CA

Introduction

To a large extent the quality of multiple-choice (MC) items depends upon the ability of clinical instructors who create the items to place themselves in the role of students, and create distractors that represent the kinds of misconceptions that students are likely to have. Since clinicians are further along the continuum of professional knowledge than students, it may be difficult for them to move back along the continuum to reason at a student level. The purpose of this study was to determine if competently reasoning clinicians could: (1) identify reasons competently reasoning students used to eliminate distractors in MC items, and, (2) explain how students would reason to select the keyed response when solving MC items. For this study, competence was defined as those individuals who chose the keyed response.

The theoretical foundation underlying this study was an information processing theory proposed by John Anderson (1983; 1993), the ACT* theory. According to Anderson (1993), cognitive processes used by problem solvers can be viewed as a sequence of internal states of knowledge successively being transformed from declarative knowledge processing and deeper problem representation through to practice and expert problem solving skills. Declarative knowledge becomes very large and well organized while the procedural knowledge becomes more

specialized and efficient.

Anderson (1983, 1993, 1994) explained that declarative knowledge and procedural knowledge are distinct. The initial stages of cognitive skill development are termed the interpretive stages. Declarative knowledge is conceptual and factual and is the result of acquiring information about something. Novices use declarative knowledge with general procedural knowledge to solve domain-related problems. Eventually, the declarative knowledge becomes elaborate and well organized. However, as the complexity of problems increases, general knowledge is insufficient to reach a solution. When individuals reach a state where there are no adequate solutions, they will search for an example of a similar problem-solving situation and try to solve the problem by analogy to that example. Specific scenarios are recalled and interpreted. Knowledge is retrieved from declarative memory. Individuals rehearse pivotal issues from the situation and substantial verbalization can occur. As the verbalization lessens, the cognitive skills are encoded procedurally. This transition is termed knowledge compilation and is the associative link between the interpretive stage and autonomous stage of procedural cognitive skill acquisition.

The cognitive skill assessment model proposed by Glaser, Lesgold, and Lajoie (1985) functioned in concert with the ACT* theory (Anderson, 1983; 1993) to provide a six dimension continuum for determining levels of competent reasoning in medical examinations. As noted earlier, assessment of competency requires clinician instructors to project how far along the continuum of medical education the student should be. This task requires clinician instructors to assess the level of clinical reasoning skills of the medical students about to enter residency. These skills tend to be sequential in a domain-specific area of knowledge and span several dimensions, from knowledge differences of novices to experts, with intermediate stages, and transitions from level to level. First, beginners' knowledge is fragmented. It consists of isolated

facts, definitions, and a superficial understanding of the domain vocabulary. At the first level of assessment, knowledge organization and structure, can be examined to determine the elements and components in a domain of knowledge. The objective is to determine the interconnections between the chunks of information and fragments of knowledge.

Second, one can assess the depth of problem representation an individual possesses. Novices typically recognize the surface features of a problem or task. More proficient individuals identify inferences, principles, and concepts imbedded below the surface structure. Individuals who immediately recognize underlying principles solve problems rapidly by spending little time on details. They arrive at the correct answer quickly.

Third, the quality of mental model can be investigated. Individuals develop mental models of scenarios and situations consistent with their domains of knowledge. The nature of these representations is determined by the tasks necessary to execute the performance of skills at a specific level. As tasks become more complex, the model is amended to incorporate these new skills. The mental model not only indicates the level of task complexity but also the level of cognitive ability needed to solve the problem.

Fourth, the efficiency of carrying out procedures can be looked at. As individuals' mental models become more refined, tasks are performed more effectively. Well-practiced procedures are important for understanding and comprehension of a scenario. At this level, assessment should focus of the relationship between understanding the requirements on a task and performing the task.

Fifth, automaticity should be present. At higher levels of cognitive development, previously learned basic skills are practiced enough so that they become automatized and are performed with little conscious attention. When performing complex tasks, individuals' attentional demands are taxed simultaneously. As this occurs, the efficiency of the overall task is compromised. The

criterion for assessment is whether the automaticity of the basic process has progressed to a point where the subtasks have minimal interference on the total performance.

Sixth, metacognitive activities should be present. At this level, individuals monitor their performance and identify strategies for solving problems using procedural skills. Individuals reflect on and control their performance. They know what they know or don't know. They can predict the outcome of their performance. They can plan their activities in advance and efficiently apportion their time. They can check and monitor the outcomes of their problem solutions and attempts to learn. These regulatory skills develop with experience and practice. Evaluating metacognitive skills could be an important facet of predicting problem-solving abilities that result in learning.

Glaser et al., (1985), explained that these skills tend to be indexed in a domain of knowledge. Because these skills are not discreet but on a continuum, identifying assessment tools appropriate for each level is difficult, particularly at the transitions points and at higher levels of cognitive development.

Differentiating among the higher levels of knowledge is demanding because experts' cognitive structures are well developed. The speed of processing knowledge is so fast that the borders among levels are indistinct, particularly between efficiency of procedures and automaticity. Metacognitive activity can be difficult to identify when individuals change their train of thought, for example when they generate a hypothesis, then consider peripheral issues and offer another solution to a problem. In responding to a MC item, experts might read an item stem and generate a hypothesis. If after thinking further about the scenario, they change the hypothesis? Did the experts just change their minds, or did they reconsider the information presented in the stem? After rethinking the scenario, perhaps they did incorrectly view the situation. Are they monitoring their behaviour based on experience, or did they consider

additional information retrieved from their elaborated schemata and just change their opinion? This situation is difficult to classify in the assessment of skills hierarchy.

Royer, Cicero, and Carlos (1993) offered a strategy to assess cognitive development using the theory that Glaser et al., (1985) proposed and linked it to Anderson's (1983) ACT* model for acquisition of knowledge. These researchers suggest that knowledge acquisition runs parallel to the assessment of clinical reasoning skills along a continuum. The authors of the present study propose that knowledge acquisition, assessment of these skills, and medical education could be overlaid on the same continuum.

When clinician instructors design items they are attempting to discriminate between competent reasoning and incompetent reasoning. They must create alternatives that competently reasoning students would know to be incorrect, but that other students might be attracted to. Similarly, when they choose from a bank, they should be able to identify which options competently reasoning students should eliminate as obviously incorrect. They should know which options are moderately wrong, an indication of partial knowledge, and which option is the correct response, the keyed option. For clinicians to identify options within these categories, they must be further along the knowledge acquisition continuum than students (Anderson, 1983), but if they are further along the continuum they may not appreciate the complexity and possible ambiguity that would be apparent to competently reasoning students. When clinicians examine distractors, can they provide a rationale for eliminating a distractor as incorrect that coincided with the students' reasons? Did the clinicians' explanations of student reasoning parallel the students' rationales for choosing the keyed alternative? If clinicians successfully performed these two tasks, it would verify their ability to move down the continuum to think from a student's perspective.

Method

Selection of the Subjects

Each clinician was a physician: (1) licensed to practice medicine, (2) with more than five years of clinical practice experience, (3) with a minimum of five years of current experience in medical education, (4) who had contact with medical students in his/her daily practice, and (5) who had instructional experience. Of the 12 physicians in this study, there were four from family practice, two each from general internal medicine, infectious diseases and pulmonary medicine, and one each from haematology and gastroenterology. Forty medical students who had just completed a medicine clinical rotation in their third year of medical school were asked by Skakun (1994) to think aloud as they responded to 30 multiple-choice items selected from an item bank in the Department of Internal Medicine. Results from the four items are reported in Table 1.

Data Collection

All of the clinicians who consented to be in the study were sent preliminary materials to review. The information package included an overview of the interview, a consent form, and an example of an item being solved using a think aloud strategy. Clinicians were first asked to think aloud as they solved the item themselves, and then asked to go back and offer reasons students would use to eliminate distractors. Each clinician instructor was interviewed independently without interruption and completed in one session. Because the present study was part of a larger study on standard setting, the total time required to gather data from each clinician instructor was between two and three and one-half hours. Competent reasoning was operationally defined as choosing the keyed response for this study.

All interviews were recorded on audiotapes and field notes were taken. Transcriptions were done immediately after the interviews. After each interview was transcribed, it was verified by

listening to the audiotape while reading the protocol for accuracy (Triska, 1996). Students transcripts (i.e., protocols), obtained from Skakun (1994) were used to compare clinician reasoning and student reasoning for the same MC items.

Analysis

Once competence was established, identification was removed from all of the competently reasoning clinician and student protocols. The clinician and student protocols were randomly interspersed for each item and the items were sorted by the researcher into two categories, novice or expert, based on the reasoning displayed in the protocol (Royer, et. al., 1993). The purpose of the dichotomous sort was to determine if clinicians using expert reasoning could be separated from student novice reasoning (Table 2). In the discussion that follows, the terms expert and novice clinicians and students refer to students and clinicians who demonstrated expert and novice reasoning on a particular item.

Since the focus of this inquiry was on the behaviour of clinicians, the analysis centred on whether the reasons clinicians gave agreed with the reasons students gave for eliminating distractors. To determine if the clinicians' insights into the students' reasoning processes were accurate or inaccurate, a content analysis of each protocol was done.

For the distractor analysis, the clinicians' reasons for eliminating each distractor were categorized and coded. Students' protocols were examined to see if their reasons paralleled the clinicians' reasons. The results of the categorization and coding of the clinicians' reasons for elimination and students' protocols were verified by a clinician. The interrater agreement, using proportions, for the distractor analysis ranged from 75% to 92%.

Due to the qualitative nature of the analysis of the clinicians' and students' protocols, and due to the spontaneous nature of the clinicians' explanations, the criterion used to determine if

clinicians could reason like students was set at a fairly low level. When clinicians did not identify more than half of the reasons students used to eliminate each distractor as incorrect the data were taken as not supporting the theory. The proportion of agreement between the two groups was calculated by using the formula:

$$\text{Proportion of Agreement} = \frac{\text{Number of Reasons Listed by Clinicians and Used by Students}}{\text{Number of Reasons the Students Listed by Clinicians} + \text{Other Reasons Listed by Students}}$$

Skakun (1994) recorded only the final justification for eliminating a distractor. In the few cases where they gave more than one reason, the final one was the dominant factor in their decision.

For the keyed response analysis, clinician protocols were examined to determine how a clinician thought a student would choose the keyed alternative. The clinicians' reasons were categorized into four broad areas. An atomistic approach involved solving the item by examining each piece of information and using facts, descriptions, and recalling definitions to arrive at the correct response. In a holistic approach, the individual uses concepts and principles to solve the item. Other clinicians stated what they expected students to know to solve the item. The final group of clinicians gave no explanation of how a student solved the item.

Students' item solving strategies were compared to the clinicians' explanations. The result of the content analysis was verified by another rater (Tables 3, 4, 5, 6). The interrater agreement for the clinicians' perceptions of how students reasoned ranged from 54% to 80% (mean=67%). The interrater agreement for how students reasoned ranged from 60% to 77% (mean=69%).

Results

Clinicians were never specifically asked to disclose why a student would reject an alternative rather, they were asked to indicate which alternatives students should know to be incorrect. The results of the distractor analysis showed that clinicians do not usually offer a reason to eliminate an alternative as incorrect spontaneously. This clinician behaviour could be attributed to the clinicians' position on the knowledge continuum, that is, the individual's automaticity and efficiency of processing information. It appeared that the expert clinicians' knowledge structures were so well formed that the declarative knowledge was not proximal and they were using procedural knowledge and composed knowledge only. Where justifications were presented, clinicians did so without prompting.

A summary of the results of the proportion of agreement of reasons for eliminating alternatives between the clinicians and students showed that as a group, clinicians did not identify reasons students used to eliminate distractors for 12 of the 16 distractors (75%) for the four items (Table 7). Contrary to expectations, the expertly reasoning clinicians however were more successful in identifying reasons students used to eliminate distractors as compared the novice clinicians. However, in a general sense, with both the expert and novice clinicians, neither group effectively viewed the item from a student's levels of knowledge according to the ACT* theory (Anderson, 1983).

Clinicians captured all of the reasons students stated for only three distractors (19%). Clinicians did not identify any of the students' reasons for eliminating one distractor. This pattern of judgements suggests that the clinicians' decisions on which alternatives are implausible or plausible may depend on the item's content, the clinicians' domain knowledge, and the clinicians' ability to view the item from a student's perspective. It may be that clinicians are unsure of justifications students use to eliminate alternatives, although this was not specifically

probed in the study. Also, it may be the case that if clinician instructors cannot identify students' probable misconceptions, the standard of performance demanded by the item might be at a level inconsistent with the students' knowledge levels at this point in their medical training.

The results of the clinicians' perceptions of how students reasoned to choose the keyed alternative for the four items varied (Table 8). The novice reasoning clinicians thought students would use a variety of strategies to reason to the keyed response. Novice clinicians gave more explanations that were parallel to students' solutions whereas expert clinicians described students' solutions in more general terms (i.e., expectations). This finding suggests that novice clinicians were more likely to move down the knowledge continuum to the students position on the ACT* continuum. This group of clinicians' judgements, the competent novice reasoning clinicians, would be in concert with the students cognitive development.

None of the expertly reasoning clinicians said that students would solve any of the items using an atomistic approach. These clinicians thought that students would use a higher level of knowledge to reason to the keyed response, that is, students would use a conceptual, principled approach to solving the item. The expert reasoning clinicians tended to state their expectation of the students' knowledge levels, and frequently did not explain how students would reason to select the keyed response. This group of competently reasoning clinicians did not move to the students positions on the ACT* continuum. Clinicians in this category might select items for an examination beyond the capabilities of the students at a given level.

The findings of the clinicians' perceptions of students' reasoning for solving items revealed two prominent issues. First, expertly reasoning clinicians were more successful in identifying the students' justifications for eliminating distractors than novice reasoning clinicians. There were fewer novice reasoning clinicians in this study, therefore a note of caution in interpreting the results is needed. Second, novice reasoning clinicians were more successful than expert reasoning

clinicians in explaining how students reasoned to select the keyed response. These results suggest that the competently reasoning expert and novice clinicians' decisions complimented each other in identifying the distractor elimination and perceptions of how students solved items.

In view of the results of this study, further research should be conducted in linking cognitive psychology with: (1) measurement, and (2) standard setting. In measurement, the effect of cognitive psychology training and item writing for clinicians on item analysis and distractor analysis should be investigated. Distractor analyses should be done to compare the effect of clinician responses before and after prompting to explain misconceptions in distractors, then compared with reasons students gave to eliminate distractors. Item statistics should be compared before and after clinicians were given instruction on the students' position on the knowledge continuum.

In setting standards of performance, clinicians require an understanding of certain aspects of cognition, particularly knowledge acquisition and the position of clinicians and students on the knowledge continuum. Clinicians must be made aware of their level of automaticity, efficiency of processing information, and highly proceduralized and composed knowledge structures, and how they differ from the students. Without this awareness, clinicians may set unrealistic standards for students at a particular level of training.

Standard setting committee members should be aware of the role of the experts' and novices' knowledge structures and decision-making characteristics to use both effectively. This study showed that experts' judgements had higher average consistency than the novices' judgements, however the novices were sometimes more successful in viewing an item from a student's perspective. Having both experts and novices reach consensus supports the reproducibility and validity of the standard setting judgements.

The time has come to integrate cognitive psychology with psychometrics in order to produce valid MC items and reproducible standard setting judgements of performance.

References

- Anderson, J. R. (1983). *The architecture of cognition*. Cambridge, MA: Harvard University Press.
- Anderson, J. R. (1993). Problem solving and learning. *American Psychologist*, 48(1), 35-44.
- Anderson, J. R. (1995). *Cognitive psychology and its implications* (4th ed.) New York: W. H. Freeman.
- Glaser, R., Lesgold, A., & Lajoie, S. (1985). Toward a cognitive theory for the measurement of achievement. In R. R. Ronning, J. Glover, J. C. Conoley, & J. C. Witt (Eds.) *The influence of psychology on testing and measurement* (pp. 41-85). Hilldale, NJ: Lawrence Erlbaum Associates Publishers.
- Nedelsky, L. (1954). Absolute grading standards for objective tests. *Educational & Psychological Measurement*, 14, 3-19.
- Royer, JM, Cicero, CA, & Carlo, MS. (1993). Techniques and procedures for assessing cognitive skills. *Review of Educational Research*, 63, 201-243.
- Skakun, E. N. (1994). *Strategy choices in responding to multiple items*. Unpublished doctoral dissertation. University of Alberta, Edmonton, Alberta, Canada.
- Triska, O. H. (1996). *The validity of clinicians' judgements in standard setting*. Unpublished doctoral dissertation. University of Alberta, Edmonton Alberta, Canada.

Table 1

Multiple Choice Item Used in the Study

ITEM 1

A previously healthy 27 year-old female is suddenly seized with pleuritic pain in the left chest and shortness of breath. The most likely cause is:

1. mycoplasma pneumonia.
- * 2. spontaneous pneumothorax.
3. pulmonary embolism.
4. acute pericarditis.
5. pleurodynia.

ITEM 2

A 28 year-old environmental activist has a history of having had pneumonia four times in the past twenty years. She has had a productive cough "all her life" which is worse in the winter. Physical examination reveals dullness, diminished breath sounds and numerous crepitations below T3 bilaterally. Her fingers are clubbed. The most likely diagnosis is:

1. hypogammaglobulinemia.
2. congenital heart disease.
3. bronchiolitis obliterans.
- * 4. bronchiectasis.
5. cystic fibrosis.

ITEM 3

A 56 year-old man presents with a month history of intermittent right facial pain. On examination he is found to have a diminished right corneal reflex and a slight hearing defect on the same side. The diagnosis is:

1. right cerebral tumour.
2. trigeminal neuralgia.
3. otitis media.
- * 4. acoustic neuroma.
5. multiple sclerosis.

ITEM 4

A 24 year-old airline flight attendant complains of feeling tired and losing weight in spite of a good appetite. For the past year she has noticed voluminous, pale, foul-smelling stools. She recalls being told of having bowel difficulty in early childhood and of being fed a diet largely consisting of bananas. Radiological examination discloses an abnormal small bowel follow through. Biochemical analysis of the stool shows an increased amount of fat. The blood picture shows anemia. Which of the following diets would you select for this patient?

- * 1. Gluten free
2. Lactose free
3. Low fat
4. Low residue
5. High residue

Table 2

Item 1: Clinicians' and Students' Problem Solving Strategies

Group	Expert Reasoners	Novice Reasoners	Total
Clinicians	8	3	11 of 12
Students	16	5	21 of 40

Item 2: Clinicians' and Students' Problem Solving Strategies

Group	Expert Reasoners	Novice Reasoners	Total
Clinicians	3	2	5 of 12
Students	14	8	14 of 40

Item 3: Clinicians' and Students' Problem Solving Strategies

Group	Expert Reasoners	Novice Reasoners	Total
Clinicians	3	5	8 of 12
Students	9	9	18 of 40

Item 4: Clinicians' and Students' Problem Solving Strategies

Group	Expert Reasoners	Novice Reasoners	Total
Clinicians	9	2	11 of 12
Students	19	13	19 of 40

Table 3

Item 1: Summary of Reasons for Eliminating Distractors

Stem: A previously healthy 27 year-old female is suddenly seized with pleuritic pain in the left chest and shortness of breath. The most likely cause is:

Keyed Option: 2. Spontaneous pneumothorax.

Alternative 1. Mycoplasma pneumonia.

Reasons for elimination.	Eliminated by		Students
	Clinicians E*	N**	
1. Mycoplasma pneumonia is not characterized as a catastrophic event.	X***	O****	X
2. Additional symptoms are present with mycoplasma pneumonia.	X	X	X
3. The number of pneumonias is unusual.	X	O	X
4. The presentation does not fit with mycoplasma pneumonia.	O	O	X

Alternative 3. Pulmonary embolism.

1. No risk factors are stated.	X	O	X
2. The history does not indicate a pulmonary embolism.	O	O	X
3. The age is inconsistent with pulmonary embolism.	O	O	X

Alternative 4. Acute pericarditis.

1. Suddenly seized is inconsistent with acute pericarditis.	X	O	X
2. Acute pericarditis is quite unlikely due to the absence of symptoms if a viral infection or bacterial infection.	O	X	O
3. Given the framework of this question, this is incorrect.	O	O	X
4. The age is inconsistent with acute pericarditis.	O	O	X

Alternative 5. Pleurodynia.

1. Can be associated with a prodrome or premonitory symptoms, such as a viral infection or fever.	X	O	O
2. Pleurodynia is not associated with a sudden onset of pain.	X	O	X
3. Pleurodynia is not associated with shortness of breath.	X	O	O
4. Pleurodynia can present like this.	X	O	O
5. Do not know what pleurodynia is.	O	O	X

Note: E* = Expert reasoning clinicians; N** = Novice reasoning clinicians.
X*** = Reason given; O**** = No reason given

Table 4

Item 2: Summary of Reasons for Eliminating Distractors

Stem: A 28 year-old environmental activist has a history of having had pneumonia four times in the past twenty years. She has had a productive cough "all her life" which is worse in the winter. Physical examination reveals dullness, diminished breath sounds and numerous crepitations below T3 bilaterally. Her fingers are clubbed. The most likely diagnosis is:

Keyed Option. 4. Bronchiectasis.

Alternative 1. Hypogammaglobulinemia.

Reasons for elimination.	Eliminated by		
	Eliminated by Clinicians (E)	Students (N)	
1. Skin lesion infections are often associated with hypogammaglob.	X	O	O
2. Does not present with these physical findings.	X	O	X
3. Recurrent infections are present with hypogammaglobulinemia.	O	O	X

Alternative 2. Congenital heart disease.

1. Additional findings would be present.	X	O	X
2. Congenital heart disease presents with shortness of breath, cyanosis, and clubbing.	X	O	X
3. History indicative of pulmonary problems.	X	O	X

Alternative 3. Bronchiolitis obliterans.

1. A disease not known by the students.	X	O	X
2. Students could be confused and think that this on one of the manifestations of bronchiolitis obliterans.	O	O	O
3. This is the wrong presentation for bronchiolitis obliterans.	O	O	X
4. This disease is more of an autoimmune problem.	O	O	X
5. Not enough information to diagnose bronchiolitis obliterans.	O	O	X
6. Bronchiolitis obliterans progresses faster.	O	O	X

Alternative 5. Cystic fibrosis.

1. In comparing bronchiectasis with cystic fibrosis, bronchiectasis would be much more likely.	X	O	O
2. Age is inconsistent with cystic fibrosis.	O	O	X
3. Cystic fibrosis affect the upper respiratory tract.	O	O	X
4. More problems are associated cystic fibrosis.	O	O	X
5. Not information to diagnose cystic fibrosis.	O	O	X

Note: E* = Expert reasoning clinicians;
X*** = Reason given;

N** = Novice reasoning clinicians.
O**** = No reason given.

Table 5

Item 3: Summary of Reasons for Eliminating Distractors

Stem: A 56 year-old man presents with a month history of intermittent right facial pain. On examination he is found to have a diminished right corneal reflex and a slight hearing defect on the same side. The diagnosis is:

Keyed Option. 4. acoustic neuroma.

Alternative 1. Right cerebral tumour.

Reasons for elimination.	Eliminated by		
	Clinician E*	N**	Student
1. Symptoms should be on the opposite side.	X***	X	O****
2. Doesn't explain the pain and diminished corneal reflex	X	O	X
3. Doesn't cause right facial pain.	O	X	O
4. Symptoms suggest cranial nerve involvement, not cerebellum.	O	O	X
5. Symptoms suggest that the problem is not in the rt. cerebellar area.	O	O	X
6. The symptoms would be more localized.	O	O	X
7. The symptoms would be more generalized.	O	O	X
7. Do not know how this relates to the scenario.	O	O	X

Alternative 2. Trigeminal neuralgia.

1. Physical findings are not totally explained by trigeminal neuralgia.	X	X	X
2. Does not have a diminished corneal reflex.	X	O	O

Alternative 3. Otitis media.

1. Doesn't explain the corneal reflex involvement.	O	X	X
2. Not a probable presentation of otitis media.	X	O	X
3. No prior history of chronic ear inflammation.	X	O	X

Alternative 5. Multiple sclerosis.

1. Multiple sclerosis is more central.	X	O	X
2. Multiple sclerosis does not produce pain.	X	O	O
3. Multiple sclerosis is more specific.	X	O	X
4. Not a typical presentation of multiple sclerosis.	O	O	X
5. Optic nerve involvement is not present.	O	O	X

Note: E* = Expert reasoning clinicians; N** = Novice reasoning clinicians.
X*** = Reason given; O**** = No reason given.

Table 6

Item 4: Summary of Reasons for Eliminating Distractors

Stem: A 24 year-old airline flight attendant complains of feeling tired and losing weight in spite of a good appetite. For the past year she has noticed voluminous, pale, foul-smelling stools. She recalls being told of having bowel difficulty in early childhood and of being fed a diet consisting largely of bananas. Radiological examination discloses an abnormal small bowel follow through. Biochemical analysis of the stool shows an increased amount of fat. The blood picture shows anemia. Which of the following diets would you select for this patient?

Keyed Option 1. Gluten free

Alternative 2. Lactose Free.

Reasons for elimination.	Eliminated by	
	Clinicians	Students
	E*	N**
1. Lactose intolerance does not result in an abnormal small bowel follow though.	X*** X	O****
2. Excess fat in the stool not associated with lactose intolerance.	X O	O
3. Weight loss not associated with lactose intolerance.	X O	O
4. Fatigue not associated with lactose intolerance.	X O	O
5. Anemia not associated with lactose intolerance.	X O	O
6. This scenario is not a presentation of lactose intolerance.	O O	X
7. Diarrhea is not present.	O O	X

Alternative 3. Low Fat.

1. Not going to help.	X O	X
2. This is not a malabsorptive diagnosis.	X O	X
3. No pancreatic or gall bladder disease is present.	O O	X
4. The mucosa is not damaged.	O O	X
5. This diet will not address the underlying pathophysiology.	O O	X
6. Fat is needed in the diet.	O O	X
7. Do not know what this will do for the problem.	O O	X

Alternative 4. Low Residue.

1. Not going to help.	X O	X
2. Not a large bowel problem.	O O	X
3. Do not know what this will do for the problem.	O O	X

Alternative 5. High residue.

1. Not going to help.	X O	X
2. Will make their voluminous stools more voluminous.	X O	O
3. Not a large bowel problem.	O O	X
4. No diarrhea is present	O O	X
5. Do not know what this will do for the problem.	O O	X

Note: E* = Expert reasoning clinicians;
X*** = Reason given;

N** = Novice reasoning clinicians.
O**** = No reason given

Table 7

Summary of the Agreement of Distractors Elimination Between the Clinicians and Students

Item No.	Distractor			
	1st	2nd	3rd	4th
1	0.75	0.33*	0.50*	0.50*
2	0.50*	1.00	0.20*	0.00*
3	0.17*	1.00	1.00	0.50*
4	0.29*	0.28*	0.33*	0.25*

Note: *Clinicians did not successfully identify >0.50 of the reasons students gave for eliminating a distractor.

Table 8

Summary of the Clinicians' Perceptions of the Students' Reasoning to Solve Items

Item	Novice Clinicians				Expert Clinicians			
	A	H	E	NE	A	H	E	NE
1	O*	X**	O	O	O	X	X	X
2	X	X	O	O	O	O	X	O
3	X	X	X	O	O	X	X	X
4	O	O	X	X	O	X	X	X

Note: O* = No clinicians; X** = Clinicians in this category.

FM 26830



U.S. Department of Education
Office of Educational Research and Improvement (OERI)
Educational Resources Information Center (ERIC)



REPRODUCTION RELEASE

(Specific Document)

I. DOCUMENT IDENTIFICATION:

Title: <i>CLINICIAN'S EXPLANATIONS OF STUDENTS' REASONING WHILE SOLVING MULTIPLE-CHOICE ITEMS</i>	
Author(s): <i>D.H. TRISKA, T.O. MAGUIRE, L.R. HARLEY</i>	
Corporate Source: <i>UNIV. OF ALBERTA, EDMONTON, AB CANADA, T6G 2G5</i>	Publication Date: <i>MARCH 26/97</i>

II. REPRODUCTION RELEASE:

In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, *Resources in Education* (RIE), are usually made available to users in microfiche, reproduced paper copy, and electronic/optical media, and sold through the ERIC Document Reproduction Service (EDRS) or other ERIC vendors. Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to the document.

If permission is granted to reproduce and disseminate the identified document, please CHECK ONE of the following two options and sign at the bottom of the page.

The sample sticker shown below will be affixed to all Level 1 documents

The sample sticker shown below will be affixed to all Level 2 documents



Check here
For Level 1 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical) and paper copy.

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY

Sample

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 1



Check here
For Level 2 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical), but not in paper copy.

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN OTHER THAN PAPER COPY HAS BEEN GRANTED BY

Sample

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 2

Documents will be processed as indicated provided reproduction quality permits. If permission to reproduce is granted, but neither box is checked, documents will be processed at Level 1.

"I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate this document as indicated above. Reproduction from the ERIC microfiche or electronic/optical media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries."

Sign here → please

Signature: <i>Dr. Olive H. Triska</i>	Printed Name/Position/Title: <i>DR. OLIVE H TRISKA PSYCHOMETRICIAN</i>	
Organization/Address: <i>C.R.A.M.E., FACULTY OF EDUCATION 3-104 EDUCATION CENTRE NORTH UNIV. OF ALBERTA EDMONTON, AB CANADA T6G 2G5</i>	Telephone: <i>(403) 482-3762</i>	FAX: <i>(403) 492-0061</i>
	E-Mail Address:	Date: <i>March 25/97</i>

III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of the document from another source, please provide the following information regarding the availability of the document. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:
Address:
Price:

IV. REFERRAL OF ERIC TO COPYRIGHT/REPRODUCTION RIGHTS HOLDER:

If the right to grant reproduction release is held by someone other than the addressee, please provide the appropriate name and address:

Name:
Address:

V. WHERE TO SEND THIS FORM:

Send this form to the following ERIC Clearinghouse:

ERIC Clearinghouse on Assessment and Evaluation
210 O'Boyle Hall
The Catholic University of America
Washington, DC 20064

However, if solicited by the ERIC Facility, or if making an unsolicited contribution to ERIC, return this form (and the document being contributed) to:

ERIC Processing and Reference Facility
1100 West Street, 2d Floor
Laurel, Maryland 20707-3598

Telephone: 301-497-4080
Toll Free: 800-799-3742
FAX: 301-953-0263
e-mail: ericfac@inet.ed.gov
WWW: <http://ericfac.piccard.csc.com>