

DOCUMENT RESUME

ED 410 335

UD 031 825

AUTHOR Singleton, Edward G.; Dale, Grady A., Jr.
 TITLE Lack of Co-occurring Interpersonal Violence
 Emotionally-Related Difficulties, or Alcohol and Other-Drug
 Problems among African American Youth with Conduct Disorder.
 PUB DATE [97]
 NOTE 15p.
 PUB TYPE Reports - Evaluative (142)
 EDRS PRICE MF01/PC01 Plus Postage.
 DESCRIPTORS Alcoholic Beverages; *Behavior Disorders; Behavior Problems;
 *Black Youth; Case Studies; *Drinking; *Drug Abuse;
 Etiology; Males; Research Methodology; *Violence
 IDENTIFIERS African Americans

ABSTRACT

Case studies of African American youth with conduct disorder were examined in the context of a descriptive evaluation of co-occurring substance-related problems and mental disorders. The purpose of this study was to extend the findings of the Epidemiologic Catchment Area (ECA) study of the National Institute of Mental Health into the area of comorbid assaultive violence and to evaluate recommendations made by J. D. Yoder and A. S. Kahn (1993) for noncomparative and descriptive research designs. Case records of 257 male youth offenders, aged 12-17 years, were reviewed. Variables were documented alcohol- or drug-related problems, referrals or counseling for emotional problems, and incidents of interpersonal violence. Half of these participants had at least one prior offense, and 14% were adjudicated for drug-related offenses not related to alcohol. Thirty-three percent had a history of violent crime. Subjects with an alcohol problem were 18 times more likely to have a co-occurring drug problem, 10 times more likely to have a co-occurring drug-related or mental health problem, and 5 times more likely to have an emotional problem than subjects without a history of alcohol-related difficulties. However, there was no indication of comorbidity of any of these disorders with assaultive violence. Substance-related and mental disorders may share some common pathways, but the mechanisms underlying interpersonal violence appear to be different. (Contains 32 references.) (SLD)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *

ED 410 335

Lack of Co-occurring Interpersonal Violence Emotionally-Related Difficulties, or Alcohol and Other-Drug Problems among African American Youth with Conduct Disorder

Edward G. Singleton

Morgan State University

Grady A. Dale, Jr.

Atlantic Coast Behavioral Health Services

and

University of Maryland at Baltimore

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

This document has been reproduced as received from the person or organization originating it.

Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE AND
DISSEMINATE THIS MATERIAL
HAS BEEN GRANTED BY

Edward G. Singleton

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC)

BEST COPY AVAILABLE

ED 410 335
ERIC
Full Text Provided by ERIC

This investigation reviewed the archives of more than 400 adjudicated juvenile cases. Case records of all African-American youth in the pool (i.e., 317 males and 72 females) were selected from this pool on the basis of diagnosed conduct disorder following court ordered referrals for psychological evaluation. Variables included in the study were documented alcohol-or-other-drug-use-related problems or referrals, referrals or counseling for emotional problems or mental disorders, and incidents of interpersonal violence. Incidence rates per 100 and risk ratios (unadjusted odds ratios) were calculated and compared separately for youth with and without each classification and within each class of comorbid disorders. Odd ratios were adjusted to reflect disparities in the gender and age distribution. Increased rates of incidence and risk of co-occurring substance abuse and emotional problems were noted; however, no significant relationship was found between comorbidity of the same disorders with interpersonal violence. Results were also consistent with the findings of previous investigations of psychiatric comorbidity such as the Epidemiologic Catchment Area (ECA) study of the National Institute of Mental Health which reported a wide range of co-occurring substance abuse and mental disorders, and they parallel conclusions of studies that cite frequent arrests but few violent crimes among population groups that experience emotional difficulties. With regards to study youth, therapeutic optimism not pessimism is warranted.

Considerable attention has been paid to the concepts of risk and vulnerability in discussions of societal problems, and both are essential to our understanding of the etiology of mental disorders. For example, childhood disadvantages such as adverse familial and parental circumstances serve as markers for development of emotional problems, and factors like poverty or low socioeconomic status are typically mentioned as risk factors for drug and alcohol use, while conduct disorder and

early childhood violence are often cited as precursors of antisocial personality disorder (American Psychiatric Association, 1994; Arkin & Funkhouser, 1990; Brunswick, Messeri, & Titus., 1992; Clayton, 1992; Hammond & Yung, 1993; Hawkins, Lishner, Jenson, & Catalano, 1987; Newcomb, 1992; Rodney, Mupier, & Crafter, 1996; Romano, Bloom, & Syme, 1991).

Epidemiology studies are the hallmark of risk and vulnerability research and are particularly useful in identifying groups with special intervention needs. One such investigation, the National Institute of Mental Health Epidemiological Catchment Area (NIMH ECA) Study, noted substantial comorbidity among substance-related and mental disorders, and reported that rates were noticeably higher among groups like incarcerated populations (Regier, et al., 1990). Another major thrust provided National Center for Health Statistics data which indicated that deaths caused by interpersonal violence among African American youth were seven times higher than those found among their White counterparts (Fingerhut & Makuc, 1992). Epidemiologic findings were also used to generate Healthy People 2000 (U.S. Department of Health and Human Services, 1990). Racial differences in negative health outcomes were established and emphasis was placed on the needs of special populations like African Americans. The assumption was that because of their disadvantaged status blacks experienced higher levels of risk making them more vulnerable to health problems and mental disorders.

Unfortunately, the focus on risk has also helped perpetuate a dim view of some American families and the behaviors of their children (Collier, 1992; Harper, 1991; Rhodes & Singleton & Miller, 1995; Singleton, 1989). The American public relies heavily on the judgments of science; the public, however, lacks the sophistication to truly understand the negative implications of scientific

findings on general perceptions. A particular case concerns comments made by Frederick Goodwin (the Director of NIMH at the time and one of the co-authors of the ECA study) which suggested that inner-city ghetto males were like jungle monkeys with regard to violent and sexual behaviors (see Jackson & George, 1996 for details). Clearly, research has value implications and the "...perceptual, comprehensive and systematic process of destruction can be seen in what is being done to our children and youth who are much more vulnerable to research abuse than most people in our communities realize" (Jackson & George, 1996, p. 12).

Group comparisons often lead to exaggerated distinctions and an incorrect conclusion of deficiencies (Yoder & Kahn, 1993) because racial differences may be confounded with unknown and unmeasurable combinations of nationality, ethnicity, gender, age, socioeconomic status, culture, biology, and heredity (Betancourt & Lopez, 1993; Jones, LaVeist, Lillie-Blanton, 1991; Zuckerman, 1990). For this reason, some researchers (Yoder & Kahn, 1993) have recommended that research needs to be conducted by using noncomparative and descriptive designs to help balance the literature and counteract negativism. Studies of this type should focus on members of various social categories alone. This would foster insight into how high-risk populations behave within their own group, not as they exist relative to other races who may or may not operate in a different personal, family, community, or social contexts (Jones, Laveist, Lillie-Blanton, 1991). Thus, this study examined the case records of African American male youth with conduct disorder within the context of a descriptive evaluation of co-occurring substance-related problems and mental disorders to extend the findings of ECA into the area of comorbid assaultive violence and to provide appropriate evaluation of the merit of the recommendations of Yoder and Kahn (1993).

Method

Participants

We examined records of youth offenders in the custody of the Department of Juvenile Services in the inner-city of a large Northeastern city. Records contained results of court-ordered psychiatric evaluations and profiled offenses that came before the juvenile courts between 1987-89. Only cases that met the Diagnostic and Statistical Manual for Mental Disorders criteria (see DSM-III-R, American Psychiatric Association, 1987) for the diagnosis of conduct disorder were included in the study. Consistent with the methods proposed by Yoder and Kahn (1993), records of females and racial groups other than black were excluded from the analyses. The final sample contained data on 257 male, African American adolescents who were 12-17 years of age (Mean = 14.5, S.D. = 1.6).

Procedures

Variables included in the study were documented alcohol or other-drug-use-related problems, treatment or referrals; notes, referrals, or counseling for emotional problems or mental disorders, and past or current adjudication for violent crime. According to procedures utilized in the ECA study (Regier, et al., 1990), comorbidity of alcohol, alcohol- or other-drug-use-related problems or referrals, referrals or counseling for emotional problems or mental disorders, and incidents of interpersonal violence were assessed by calculating the rates per 100 participants of each disorder (such as alcohol-related problems) with all possible combinations of the other disorders (e.g., alcohol with any drug problem; alcohol with any drug or mental disorder; alcohol with any drug or mental problem; alcohol with any drug or mental disorder or violent incident). The assessments were made separately for subjects who had and did not have a documented history of the problem.

Then, relative risk was determined by computing odds ratios (rate per 100 with the disorder divided by the rate per 100 without the disorder) for all comorbid combinations (here for example, odds ratio = 71% with alcohol-related problem or any drug-related problem divided by 4% without alcohol-related problem but who had a documented drug-related problem = 17.85). Loether and McTavish (1976) indicate that 95% confidence intervals were used to define a critical region bounding each odds ratio (for the previous example, 15.4 - 20.1). A statistically significant association among two or more disorders existed for odds ratios with critical regions that did not include 1.0, a value indicating that the risks were identical (Morris, 1970). Psychological prototypes of subjects with and without each disorder were compared and contrasted using graphical profiles of T-scores from clinical scales on the adolescent normed personality inventory. Twenty-one demographic, educational/vocational, behavioral, home environment and discipline, peer group, and community variables were entered into a forward stepwise multiple regression to identify smaller subgroups of risk factors specific to each disorder under investigation. Only significant findings will be reported to conserve space.

Results

Demographics

Half of the participants had at least one prior offense. Fourteen percent were adjudicated for drug-related offenses not related to alcohol. Thirty-three percent had a past or current history of committing a violent crime according to public health criteria where force or other means were used by the youth with intent of causing harm, physical injury, or death to another person. Most (87%) were attending school and many (66%) were identified as problem students. Eighty-three percent

had a mother or other adult female present in the home and more than one-fifth (21%) had a sibling with a prior arrest history living with the family. Average household income was \$473 per month.

Alcohol

Subjects with an alcohol problem (8.6%) were 18 times more likely to have a co-occurring drug problem, 10 times more likely to have a co-occurring drug-related or mental disorder, and five times more likely to have had an emotional problem than subjects without a history of alcohol-related difficulties. The most significant risk factor of alcohol disorder was a comorbid drug-related problem (Alcohol model, $R^2 = .47$, $p = .001$). Psychological profiles of subjects with alcohol-related problems suggested they were more assertive and less irritable, and they exhibited less severe emotional symptoms than those without a documented history of referrals or intervention for difficulties with alcohol use.

Other Drugs

Among subjects with an other-drug-related problem (7%), the risk of having a co-occurring alcohol-related and mental disorder was 10 times greater than those with no drug-related difficulty. The most prominent risk of having this disorder was a documented history of an alcohol-related problem (Drug model, $R^2 = .66$, $p = .001$). This group showed higher levels of social sophistication, and less anger and resentment than subjects without alcohol-related disorder.

Mental Disorders

Personality problems placed this group at higher risk for mental disorder (Mental disorder model, $R^2 = .42$, $p = .001$). Subjects who exhibited signs of a mental disorder (4.5%) were more than four times as likely to have had a co-existing alcohol-related problem.

Violence

Profiles of adolescents with a history of interpersonal violence suggested lack of insight, social introversion, and isolation in contrast to non-violent offenders. The greatest risk factor for assaultive violence (Violence model, $R^2 = .47$, $p = .001$) was the presence of an older sibling in the home. More importantly, this group had less or similar overall risk of having any comorbid disorder when compared to subjects who had no history of violent crime.

Discussion

This investigation examined the case histories of youth with conduct disorder within the context of a descriptive evaluation of documented co-occurring substance-related problems and mental disorders to extend the findings of ECA into the area of comorbid assaultive violence, and to assess the utility of noncomparative and descriptive designs (Yoder & Kahn, 1993). Here, reliance on documented incidences may limit the findings to the extent that the rates of any disorder may have been underreported and biased with regard to the criteria of this specific adjudication process. Other research (Betancourt & Lopez, 1993) has demonstrated that symptoms rather than diagnoses may be better indicators in studies of this kind because they may be more sensitive to sociocultural influences.

Nevertheless, results for adolescents with alcohol-related, other-drug-related, and mental disorders were generally consistent with the findings of ECA and other comorbidity studies that have reported a wide range of increased risk in co-occurring substance abuse and mental disorders (Young & Werch, 1990), especially among incarcerated populations (Regier et al., 1990). Significant multiple correlations from the multiple regression models for these disorders supported

the conclusions regarding increased risk, accounting for 18% to 44% of the explained variance. The findings strengthen the argument for the inclusion of drug and alcohol interventions within mental health programs and vice versa.

However, there was no indication of comorbidity of these disorders with assaultive violence. Substance-related and mental disorders may share some common pathways, but the mechanisms underlying interpersonal violence appear to be different. This supports the current trend in treating violence as a separate public health problem (Nagayama, Hall & Barongan, 1997) and bolsters the contention that more prevention programs be targeted at violence among African-American youth (Rodney et al., 1996). Findings also augment the general psychiatric literature about the low rates of violence in mentally disordered groups in general (Kaplan & Sadock, 1991).

On the average, this group of African-American adolescent males exhibited psychological profiles that were only slightly elevated or well within ascribed normal limits. This is not unusual because conduct disorder is symptomatic of behavioral problems and not gross psychopathology. In fact, the psychological profiles of youth with alcohol-related other-drug-related, and mental disorders were less suggestive of pathology than their non-problem counterparts. Thus, there is reason for therapeutic optimism not negativism with regards to intervention outcome. Overall, findings fail to support the notion that violence, substance abuse, and defective personalities are co-occurring (Ito, Miller, & Pollack, 1996) in the exclusive domain of African-American male youth.

The question of why the most prominent risk factor for assaultive violence was the presence of an older sibling in the home is a matter of speculation at this time. Notably, only 12% of the variance was attributed to variables in the violence risk model. This result suggests that the model

may have been underspecified. At present, the literature on violence remains underdeveloped (Hammond, 1993) and we would do well to look for sources of information other than juvenile justice records in determining the true nature of interpersonal violence (Singleton & Douglas, 1995). Many researchers (Brown & Rhodes, 1991; Clayton, 1992; Cowen, 1991; Hawkins, et al, 1987; Garmezy & Neuchterlein, 1972) have stressed taking a look at a broad range of protective factors as one alternative to the traditional risk paradigm. A protective factors model may well provide a better framework for understanding how high-risk youth avoid rather than engage in violent behavior, and yield information critical to the development of more effective intervention programs.

References

American Psychiatric Association (1987). Diagnostic and statistical manual of mental disorders iv. Washington, DC: American Psychiatric Press.

American Psychiatric Association (1994). Diagnostic and statistical manual of mental disorders iv. Washington, DC: American Psychiatric Press.

Arkin, E. B., & Funkhouser, J. E. (Eds.). (1990). Communicating about alcohol and other drugs. Strategies for reaching populations at risk. (DHHS Publication No. ADM 90-1665). Office for Substance Abuse Prevention. Washington, DC: U.S. Government Printing Office.

Betancourt, H., & Lopez, S. R. (1993). The study of culture, ethnicity, and race in American psychology. American Psychologist, 48(6), 629-637.

Brown, W. K., & Rhodes, W. A. (1991). Factors that promote invulnerability and resiliency in at-risk children. In W. A. Rhodes & W. K. Brown (Eds.). Why Some Children Succeed Despite the Odds. New York: Praeger.

Brunswick A. F., Messeri, P. A., & Titus, S. P. (1992). Predictive factors in adult substance abuse: A prospective study of African American adolescents. In M. Glantz & R. Pickens (Eds.). Vulnerability to Drug Abuse. Washington, DC: American Psychological Association.

Clayton, R. R. (1992). Transitions in drug use: Risk and protective factors. In M. Glantz & R. Pickens (Eds.), Vulnerability to Drug Abuse. Washington, DC: American Psychological Association.

- Collier, M. A. Z. K. (1992). What we must do to save our youth. Visions, 2, 6-10.
- Cowen, E. L. (1991). In pursuit of wellness. American Psychologist, 46, 404-408.
- Fingerhut, L. A., & Makuc, D. M. (1992). New from NCHS: Chartbook on minority health. American Journal of Public Health, 82, 1 167-1 170.
- Garmezly, N., & Neuchterlein, K. (1972). Invulnerable children: The fact and fiction of competence and disadvantage. American Journal of Orthopsychiatry, 42, 328-329.
- Hammond, W. R., & Yung, B. (1993). Psychology's role in the public health response to assaultive violence among young African-American men. American Psychologist, 48, 142-154.
- Harper, F. D. (1991). Substance Abuse and the Black American family. Urban Research Review, 13, 1-5.
- Hawkins, J. D., Lishner, D. M., Jenson, J. M., & Catalano, R. F. (1987). Delinquents and drugs: What the evidence suggests about prevention and treatment programming. In B. S. Brown & A. R. Mills (Eds.), Youth at High Risk for Substance Abuse (DHHS Publication No. ADM 87- 1537). National Institute on Drug Abuse. Washington, DC: U.S. Government Printing Office.
- Ito, T. A., Miller, N., & Pollock, V. E. (1996). Alcohol and aggression: A meta-analysis on the moderating effects of inhibitory cues, triggering events, and self-focused attention. Psychological Bulletin, 120, 60-82.
- Kaplan, H. I., & Sadock, B. I. (1991). Synopsis of psychiatry: Behavioral sciences, clinical psychiatry. Baltimore, MD: Williams & Wilkins.
- Jackson, J. F., & George, P. Y. (1996). Unethical research: A weapon against African

American children and youth. Psyc Discourse, 27, 10-16.

Johnston, L. (1992). How epidemiology helps us to grasp the phenomena of drug use. In Dolentium Hominum: Church and Health in the World, 19, 67-71.

Jones, C. P., Laveist, T. A., Lillie-Blanton, M. (1991). "Race" in the epidemiologic literature: An examination of the American Journal of Epidemiology, 192 1-1990. American Journal of Epidemiology, 134, 1079-1084.

Loether, H. J., & McTavish, D. G. (1976). Descriptive and inferential statistics: An introduction. Boston: Allyn and Bacon, Inc.

Nagayama Hall, G. C., & Barongan, C. (1997). Prevention of sexual aggression: Sociocultural risk and protective factors. American Psychologist, 52, 5-14.

Newcomb, M. D. (1992). Understanding the multidimensional nature of drug use and abuse: The role of consumption, risk factors, and protective factors. In M. Glantz & R. Pickens (Eds.). Vulnerability to Drug Abuse. Washington, DC: American Psychological Association.

Regier, D. A., Farmer, M. E., Rae, D. S., Locke, B. Z., Keith, S. K., Judd, L. L., & Goodwin, F. K. (1990). Comorbidity of mental disorders with alcohol and other drug abuse. JAMA, 264, 2511-2520.

Rhodes, W. A., Singleton, E. G., & Miller, F. (1995). Substance use by students at a Historically Black College. Journal of Research on Minority Affairs, 5, 65-78.

Rodney, H. E., Mupier, R., & Crafter, B. (1996). A preliminary study of the relationship between alcohol and conduct disorder. Challenge, 7, 77-86.

Romano, P. S. Bloom, J., & Syme, S. L. (1991). Smoking, social support, and hassles in

an urban African-American community. American Journal of Public Health, 81, 1415-1422.

Singleton, E. G. (1989). Substance use and Black youth: Implications of cultural and ethnic differences in adolescent alcohol, cigarette, and illicit drug use. In R. L. Jones (Ed.). Black Adolescents. Hampton, Virginia: Cobb & Henry.

Singleton, E. G., & Douglas, T. E. (1995). Identifying factors that prevent and reduce drug use to counteract ethnic bias and racial stereotyping. Proceedings of the Annual Conference, Optimal Family Mental Health: Building on the Strengths of African American Families. Baltimore: Black Mental Health Alliance.

US Department of Health and Human Services (1990). Healthy People 2000: National health promotion and disease prevention objectives. Washington, DC: (author).

Yoder, J. D., & Kahn, A. S. (1993). Working toward an inclusive psychology of women. American Psychologist, 48, 846-850.

Young M., & Werch, C. E. (1990). Relationship between self-esteem and substance use among students in fourth through twelfth grades. Wellness Perspectives: Research, Theory and Practice, 1, 31-44.

Zuckerman, M. (1990). Some dubious premises in research and theory on racial differences. American Psychologist, 45, 1297-1303.



U.S. Department of Education
Office of Educational Research and Improvement (OERI)
Educational Resources Information Center (ERIC)



REPRODUCTION RELEASE

(Specific Document)

UD031825

I. DOCUMENT IDENTIFICATION:

Title: LACK OF CO-OCCURRING INTERPERSONAL VIOLENCE WITH EMOTIONALLY-RELATED DIFFICULTIES, OR ALCOHOL- AND OTHER-DRUG-USE-RELATED PROBLEMS AMONG AFRICAN AMERICAN YOUTH WITH CONDUCT DISORDER	
Author(s): EDWARD G. SINGLETON AND GADY A. DALE, JR.	
Corporate Source:	Publication Date:

II. REPRODUCTION RELEASE:

In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, Resources in Education (RIE), are usually made available in microfiche, reproduced paper copy, and microfiche/teletype media, and sold through the ERIC Document Reproduction Service (EDRS) or other ERIC vendors. Credit is given in the source of each document, and, if reproduction release is granted, one of the following notices is affixed to the document.

If permission is granted to reproduce and disseminate the identified document, please CHECK ONE of the following two options and sign at the bottom of the page.



Check here

For Level 1 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical) and paper copy.

The example notice shown below will be affixed to all Level 1 documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY

Sample

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 1

The example notice shown below will be affixed to all Level 2 documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN OTHER THAN PAPER COPY HAS BEEN GRANTED BY

Sample

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 2



Check here

For Level 2 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical), but not in paper copy.

Documents will be processed as indicated provided reproduction quality permits. If permission to reproduce is granted, but neither box is checked, documents will be processed at Level 1.

I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate this document as indicated above. Reproduction from the ERIC microfiche or electronic media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries.

Sign here please	Signature: <i>Edward G. Singleton</i>	Printed Name/Address: EDWARD G. SINGLETON CONSULTING PSYCHOLOGIST	
	Organization/Address: E.G. SINGLETON, PH.D., PSYCHOLOGIST 5 ELDERBERRY COURT BALTIMORE, MD 21228-1008	Telephone: 410-744-0963	FAX: 410-744-4604
		E-mail Address: EDGS@msn.com	Date: 6/4/97

(11/95)