

DOCUMENT RESUME

ED 422 554

CG 028 661

AUTHOR Soley, Georgia; Marshall, Renee; Chambliss, Catherine
TITLE Training Therapists about Client Expectations of
Psychotherapy.
PUB DATE 1998-00-00
NOTE 37p.
PUB TYPE Reports - Research (143) -- Tests/Questionnaires (160)
EDRS PRICE MF01/PC02 Plus Postage.
DESCRIPTORS Counselor Client Relationship; Counselor Training;
*Expectation; Outcomes of Treatment; *Parent Attitudes;
*Psychotherapy; Termination of Treatment; *Therapists

ABSTRACT

Research has indicated that premature termination of therapy is sometimes due to a conflict in goal and outcome expectations between therapists and family members of clients. The present study requested both therapists and parents of child clients to complete questionnaires to determine if there is congruence between therapist and parental expectations. The questionnaire was designed to measure the similarity in expectations for both parties or if lack of congruence is present that may possibly lead to premature termination of therapy for the child client. Also included in the study were responses from adult clients and therapists of adult clients. Within subject t-tests of a sample of 63 parents, 18 therapists, and 43 adult clients produced significant findings. Therapists and parents of child clients attached greater significance to the behavior change sub-scale, while adults rated the five sub-scales as equally important. Appendix A contains the Parent Attitude Questionnaire with a discussion of "Parents' Motivations for Seeking Treatment for Their Children." Appendix B provides the Client Attitude Questionnaire and a discussion of "Clients' Motivations for Seeking Treatment." More than 130 references are listed. (Author/EMK)

* Reproductions supplied by EDRS are the best that can be made *
* from the original document. *

Training Therapists about Client Expectations of Psychotherapy

Georgia Soley

Renee Marshall

Catherine Chambliss, Ph.D.

Ursinus College

1998

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

"PERMISSION TO REPRODUCE THIS
MATERIAL HAS BEEN GRANTED BY

C. CHAMBLISS

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC)."

Abstract

Research has indicated that premature termination of therapy is sometimes due to a conflict in goal and outcome expectations between therapists and family members of clients. The present study requested both therapists and parents of child clients to complete questionnaires to determine if there is congruence between therapist and parental expectations. The questionnaire was designed to measure the similarity in expectations for both parties or if lack of congruence is present that may possibly lead to premature termination of therapy for the child client. Also included in the study were responses from adult clients and therapists of adult clients. Within subject t-tests of a sample of 63 parents, 18 therapists, and 43 adult clients produced significant findings. Therapists and parents of child clients attached greater significance to the behavior change sub-scale, while adults rated the five sub-scales as equally important.

Why do parents seek psychotherapy for their children? Although this knowledge would seem fundamental to the development of successful helping strategies, relatively little empirical research has addressed this issue. The previous research data has contributed to our current knowledge of consumer satisfaction with child psychotherapy; however, data has rarely been collected to analyze parental expectations for their child's treatment.

Critics of various schools of therapy have long argued that treatments that fail to address specific concerns of parents of child clients may be coherent conceptually, but of limited practical value. Premature termination and attrition from therapy outcome studies have often been attributed to treatments value to conform to parents' expectations of child psychotherapy, although parents who terminate their child's treatment prematurely have rarely been interviewed to ascertain their exact reasons for departing.

As consumers, we want to be satisfied with the quality of the products or services we are spending money to obtain. In order to ensure our satisfaction, consumerism has provided an abundance of product alternatives from which to choose. When the thought of product choice comes to mind, tangibility usually accompanies that thought; but, as consumers we have an array of intangible services that hopefully enhance lifestyle. This intangibility may affect the ability to accurately measure its quality. One of these consumer goods is child psychotherapy. What is the best way to measure the success of therapy given to children? Some of our measurements would look at therapist flexibility, pathology change, behavior improvement, and parent perception of child improvement. The production of true results can be quite a challenge as demonstrated by some studies

designed to measure the satisfaction of consumers of mental health services. Depending on the definition used for consumer satisfaction which can be construed in a very broad sense, it would be a safe assumption to conclude that parents of antisocial children would stop pursuing treatment if they are not finding a substantial behavioral change in their child. It is possible to equate the parental drop-out rate with dissatisfaction of the services provided while at the same time stating that this is a very simplistic view; there can be a myriad of other economic, social, familial, and personal reasons why parents would choose to end treatment before completion. It can also be said that all of these other reasons aside, if the treatment was outstanding the parent would most likely be motivated to continue until completion.

One study investigated the reasons for termination of mental health treatment prior to completion of the program. The study was initiated to help families with children who exhibit severe antisocial behavior (Kazdin, 1987; Miller & Prinz, 1990; Patterson, 1986). Using a multiple gating strategy, the sample included 4 to 9 year old boys from 147 families meeting the criteria for treatment. Nine therapists were used in the study consisting of 6 women and 3 men, all possessing master's degrees in clinical or school psychology, as well as 2 years of supervised family experience (Miller & Prinz, 1994). The families were randomly assigned to an EFT (Enhanced Family Treatment) or an SFT (Standard Family Treatment). SFT is a conventional approach to therapy for parent-child interactions. EFT combines the approaches of SFT with an additional focus on issues the parent is confronting. At the outset, the advantages of both approaches were reviewed. If the therapist were to focus only on parent-child interaction, the effectiveness might be reduced because parents' issues would not be resolved.

Conversely, some parents might feel uncomfortable with the focus on themselves (Miller & Prinz, 1990; Wahler & Dumas, 1989). In either of these situations, without knowledge of the parents' feelings, it might be an incentive to discontinue treatment. "Measures included the following: (a) parent and teacher reports on Achenbach's Child Behavior Checklist; (b) parental report of personal adjustment on the Symptom Distress Checklist-90 (SCL-90; Derogatis, Rickels, & Rock, 1976), marital adjustment on the Dyadic Adjustment Scale (DAS; Spanier & Thompson, 1982), and social support on the Arizona Social Support Interview Schedule (ASSIS; Sandler & Barrera, 1984); (c) Hollingshead four-factor socioeconomic status index."

Dropout rates for each program were as follows: 29.2% for EFT and 46.7% for SFT. Dropouts were interviewed by phone to determine their reasons for discontinuation of treatment. Situational factors were cited for the reason more frequently for SFT (73%) versus EFT (47%), but dissatisfaction with intervention was cited much more frequently for SFT (26%) versus EFT (6%).

These results indicate parents need to be provided with supportive services for personal life issues as well as their socially maladaptive children. Giving parents a way to release frustration is imperative for children to continue to receive treatment. It also lends support to the broader issue that consumer satisfaction does indeed make a difference in child therapy. It is important for parents to be satisfied with the treatment of their children and part of this satisfaction may involve providing services to both children and parents.

Due to the importance of treatment completion, it is necessary to discover the specifics which lead to premature termination. The rate of dropout among families

receiving child therapy has been estimated at 40-60% (Kazdin, 1996a; Wierzbicki & Pekarik, 1993). It is assumed that there are barriers to the continuation of treatment such as socioeconomic status. We have already established that there may be issues related to satisfaction with therapists' services. Do all participants who experience barriers to treatment drop out regardless of the level of satisfaction? Premature termination can be detrimental for the families who need the service but there are broader implications. The accuracy of research is affected by the number of dropouts. Sample size and sample characteristics are altered making generalization difficult. Therefore, it is essential to measure the causes of premature termination of treatment for individual and statistical purposes.

The current research identifies social characteristics contributing to premature termination of treatment. "Socioeconomic status, minority group, high levels of stress and family dysfunction, and difficult living circumstances (e.g., single-parent families) are among the salient factors" (Armbruster & Kazdin, 1994; Gould, Shaffer, & Kaplan, 1985). Some other less studied issues relating to drop out are "parental stress and life events, parent psychopathology, and severity of child externalizing problems" (Kazdin, A.E., Holland, L., & Crowley, M., 1997). However, several of these characteristics may produce more than one barrier to receiving treatment, so it is difficult to determine what factor of socioeconomic status, for example, is actually causing the decline. Other factors unrelated to situation may also be involved, such as the quality of therapy and the client-therapist relationship.

The study of premature termination of therapy was designed to test three main hypotheses using a barriers to treatment model which emphasizes not only personal

factors but also the experience in the therapeutic setting: “(1) Family, parent and child factors identified at intake assessment would predict dropping out of treatment, (2) barriers to treatment would predict the rate of drop out, (3) barriers to treatment will increase the rate of drop out incrementally beyond the more fundamental causes of drop out mentioned earlier” (Kazdin, A. E., Holland, L., Crowley, M., 1997). Of the 242 individuals who participated in the study, 146 completed the program, and 96 dropped out of the study.

The findings indicated that perceived barriers to treatment, both fundamental and those related to treatment, perceptions of the treatment process itself, and relationship of parent with therapist all influenced the premature termination of therapy. These results should be tempered with some limitations. The sample included children who were referred to the program and included only cognitive and cognitive-behavioral clinics so diversity in treatment cannot be generalized. Also, responses were collected subsequent to treatment; some recall bias may have influenced results. Finally, it has been established that parents make the decision to continue in treatment, but this ignores the possibility that the child may influence directly or indirectly the final decision of the parent to continue or terminate treatment.

Effects of Managed Care on Studies of Satisfaction

The advent of managed care has brought a new fervor for satisfaction measures in the field of behavioral therapy. Consumers and providers of care need feedback on the effectiveness of therapy. Mental health professionals can point to studies that demonstrate that treatment is beneficial but these studies are increasingly being challenged for more relevant data (e.g. Casey & Berman, 1985; Smith, Glass, & Miller,

1980; Weisz, Weiss, Alicke, & Klotz, 1987). Most study results are based on lab findings, not those done in the field. Mental health researchers have been challenged on the applicability of lab settings to those in the field (see Hoagwood, Hibbs, Brent, & Jenson). This has provided a heightened awareness of the need to obtain results from those in therapy. The few studies that *have been* conducted provide little consistency in results for adults or children and adolescents (Weisz, Weiss, & Donenberg).

A field based study must take into account constructs of interest and methods. Constructs of interest refer to what is being measured and methods refer to how it is being measured. Most evaluations of effectiveness use pathology change as a barometer. “Clinical effectiveness is determined by (a) the extent to which symptoms decrease or functioning increases, and (b) whether scores for a treated group are significantly better than scores for a comparison group (Lambert, Salzer, & Bickman, 1998).” Some obstacles to getting at the information required are cost, data collection methods, and the factors to measure pathology change. Behavioral Health Organizations find that satisfaction and improvement are the best indicators of effectiveness and because most of the studies would be funded by the BHO’s, the choice of method would also be determined by the BHO.

Unfortunately, BHO’s are not only collecting the data to provide measurements of satisfaction to consumers but also to classify what should and should not be considered a mental illness in need of treatment. A mental illness by definition may be difficult to organize, categorize, and classify. The DSM, for this reason, is in a state of continual change. It should be mentioned as well that some therapists resist the quantification of success rates. This is a healing profession and as with any medical doctor, diagnosis and

treatment may take extended periods of time. But if a physical illness cannot be identified, doctors do not give up; they continue searching for a cause based upon symptoms. It is no different for therapists and why should it be?

The problem with using satisfaction as the primary determinant of pathology change is the lack of correlation between various studies that link satisfaction with pathology. Some studies show moderate correlation, yet most do not demonstrate consistent results (Attkisson & Zwick, 1982). Additionally, the level of agreement between ratings by therapist and client have varied greatly (Weiss, Rabinowitz, & Spiro, 1996). Also, self report questionnaires must be differentiated for parent and child. The reason for this inconsistency has been cited as method variance.

A study was designed to compare the effectiveness of traditional therapy services with a quasi-experimental continuum of care design. In designing the study, a multitrait-multimethod model was used to achieve consistent results.

Informants included the client, a parent, and a trained interviewer, with additional data from teachers and providers. For the present study, only adolescent clients were used because younger children did not fill out self-reports...In the present study, we examine relationships among pathology change (as reported by adolescents, parents, and trained interviewers), client satisfaction (as reported by adolescents and their parents), and perceived improvement (reported by parents).

Three hypotheses were tested: (1) Using different informants would provide a greater match to the data than using a shared method, (2) pathology change and satisfaction measures should be kept separate, (3) parent reporting of perceived improvement will be more indicative

of satisfaction than pathology change (Lambert, Salzer, & Bickman, 1998).

The results of the study support the first two hypotheses. It is necessary to distinguish between symptom change and satisfaction as well as maintaining the distinctions between the responses of the interviewer, the parent, and the adolescent. The third hypothesis needs to be retested. Reports of parent perceived improvement were measures more so of satisfaction than pathology change. On the contrary, reports from adolescents of satisfaction were highly correlated with pathology change.

Collaboration and Family Satisfaction

Keeping in mind that the relationship of parent with therapist influences premature termination of therapy, one study by DeChillo, Koren, and Schultz examined the relationship between the collaboration of caregivers and mental health care professionals and family satisfaction. DeChillo, et al (1994) noted earlier studies that found family members of individuals with mental health issues unhappy with the services provided to them by professionals in the field. The dissatisfied family members felt as though they were being blamed for the problems of the individual in need of services and that their needs were not being met by the mental health care providers (Grunebaum, 1984; Hatfield, 1982; McElroy, 1987; Spaniol, Jung, Zipple, & Fitzgerald, 1987). In order to investigate the numerous reports of family dissatisfaction, DeChillo et al pointed to four studies (Benheim & Switalski, 1988; McElroy, 1987; Smets, 1982; Spaniol, Zipple, & Fitzgerald, 1984) which explored the relationship of mental health care professionals and family members of mental health patients. It was discovered that there were two main factors contributing to family dissatisfaction: the lack of family involvement in the treatment of the patient and disagreement between the health care

professional and family members on what treatment was best for the individual in need (DeChillo, Koren, & Schultz, 1994). Studies suggest that patient outcome could be improved if there was collaboration among the patient's family members and those in charge of providing mental health care (DeChillo, Koren, & Schultz, 1994).

DeChillo et al (1994) define collaboration as "two or more parties working together in pursuit of a common goal." In 1993, DeChillo conducted a survey of family members of adult patients on a psychiatric unit to appraise collaboration. Studying the contributing factors of collaboration and their effect on the patient's outcome, DeChillo found the strongest predictor of collaboration was the attitude of the social worker on the issue of family involvement (DeChillo, 1993). One point that was stressed in DeChillo's prior study was the importance of the identification of a common goal between family members of the patient and the social worker (DeChillo, 1993).

In their recent study, DeChillo et al (1994) investigated elements of collaboration by having 455 family members of children with emotional disorders complete a questionnaire intended to measure eight components of collaboration: (1) joint decision making and planning; (2) locating and developing services; (3) funding services; (4) evaluation based on feedback from family; (5) conveying a caring attitude; (6) sharing information; (7) recognizing the family as a resource, and (8) recognizing limits of the family. Caregivers were given one of two questionnaires (one asked the family member to complete the form with regard to a professional in the field he or she felt was easiest to work with; the other was to be filled out with regard to a professional the caregiver felt was most difficult to work with). The study recognized four key elements of collaboration: supportive relationships, which refers to the feeling of partnership between

the professional and caregiver; practical service arrangements, meaning the financial aspects and the coordination and development of services; information sharing, such as the explanations and communication efforts put forth by the professional to the family member; and a flexible, shared approach to measuring failure or success, the professional taking seriously and working with feedback from the caregivers (DeChillo et al, 1994). DeChillo et al found “a very strong relationship between satisfaction and collaboration” (1994). This is consistent with prior research revealing the importance of congruence between therapist and family members of the patient.

Parental Satisfaction with Inpatient Mental Health Services

In an effort to address what is considered to be a neglected area of research, a study was conducted at a state psychiatric center in Staten Island, New York concerning parental satisfaction with mental health care services (Byalin, 1993). Using the Client Satisfaction Questionnaire (CSQ), which can be measured on a standardized scale (Attkisson & Zwick, 1980), parents of recently discharged children and adolescents between the ages of 12 and 18 were asked to give their responses with regard to the services their children received while they were patients in the psychiatric center (Byalin, 1993).

Byalin (1993) points out that there has been little attention and priority given to parental and family satisfaction with services in the mental health field with the exception of a noteworthy 1978 study in which Woodward and Associates researched consumer satisfaction with brief family therapy (Woodward, Santa-Barbara, Levin, & Epstein, 1978). Byalin’s study was originally created to monitor and evaluate the quality of this particular psychiatric facility. The thought behind this was that parental

empowerment, which the program was targeted to increase, would equate to parental satisfaction; therefore, parental satisfaction was used as the measurement tool (Byalin, 1990).

With 53 questionnaires sent out and only 15 questionnaires returned completed, the research in many ways was a disappointment. However, some promising ideas were discovered. For example, Byalin notes that the questionnaire used for this study would be a successful tool in providing ongoing evaluations of the program. It was also found that for the most part, parents were satisfied with services with the exception of one Area--parents indicated from the responses on the questionnaire that they were unhappy that certain issues, not considered to be psychiatric, were not taken care of before their child was discharged. This led to the conclusion that staff members may be less responsive or less attuned to consumer needs they do not consider to be "psychiatric" (Byalin, 1993).

Prior research has provided some general conclusions that can be summarized. Parental satisfaction is a crucial factor in preventing premature termination of child psychotherapy. Many variables can influence satisfaction including situational status, parent-therapist relationship, and contact within the behavioral treatment setting. Most studies are either based on pre or post assessments, but not both. Future studies need to focus on pre-treatment measurements of expectations, as well as post treatment satisfaction.

The results of Reynolds, Ogiba, and Chambliss' (1998) study indicated that therapists and adult clients were largely congruent in their goal objectives. However, Soley, Hooper, Marshall, and Chambliss (1998), using a similar between subject design,

found a disparity between therapist expectations and the expectations of the parents of child clients. In the Soley et al study, parents of child clients rated normalization more highly than therapists. This suggests the possibility that therapists may have misperceived the objectives of parents bringing their children to therapy. Nonetheless, a problem with the Soley et al study was that its between subject analyses included a scaling confound; between group difference in use of the rating system may have produced misleading findings.

In order to remedy this problem, the original Soley et al study was reanalyzed using a within subject design. Using the within subject design we were able to assess the relative treatment priority for the different constituencies.

Method

Therapists from a variety of local treatment centers were asked to record their responses on a questionnaire with 29 (Likert-format) items concerning parent's attitudes when bringing their child to therapy (See appendix A). The therapists were asked to answer the questions as they believed a parent would answer it when bringing his or her child to therapy.

A sample of parents, solicited from child treatment sites, were asked to complete the same questionnaire as the therapists. Additional participating parents were contacted at a local fast food restaurant on the weekend while they were watching their children play in the play area.

Responses were obtained from 12 therapists, consisting of 7 males and 5 females, and 67 parents, consisting of 44 females and 22 males. The mean age of the therapists was 40.92 years, with an average of 14.42 years of experience in the mental health care

field. The mean age of the participating parents was 39.45 years. Approximately 67% of the therapists and 80% of the parents contacted actually completed the questionnaire.

Both the therapists and the parents were from the suburbs of Philadelphia.

Thirty percent of the participating parents reported actually having a child in therapy. The remaining parents answered the questionnaire hypothetically, based on what their expectations would be if their child ever needed therapy.

Previous research, conducted by Reynolds et al (1998), had therapists of adult clients, as well as adult clients, complete a questionnaire with 27 Likert-format items concerning clients' attitudes toward therapy (see Appendix B). Therapists were randomly selected from telephone directories, while prospective psychotherapy clients were obtained by anonymously surveying undergraduates at a small Liberal Arts college. Responses were obtained from 7 therapists of adult clients and 42 prospective clients.

A separate demographic questionnaire was provided for the therapists and for the parents of prospective and actual clients, as well as for the adult client sample. The therapist of child client's demographic questionnaire consisted of 8 items, while the therapists of adult client's demographic questionnaire consisted of 6 items. The parent's demographic questionnaire consisted of 6 items, while the adult client's questionnaire consisted of 4 items. All respondents were assured of anonymity.

Results

Relevant items were totaled for each subject, yielding five sub-scale scores assessing the treatment objectives: Normalization, Analysis, Decision-making, Behavior Change, and Affect Change. Within subject t-tests were used to compare the five sub-scale scores of all respondents considered together, the therapists, parents of child clients, and adult

clients.

When all subjects were considered together, behavior change scores were significantly higher than all of the other sub-scale scores. Affect change and decision-making assistance were rated comparably; scores for both were significantly higher than those for analysis ($p < .05$) and normalization ($p < .001$). Analysis scores were significantly higher than those for normalization ($p < .05$).

Among therapists, t-tests showed results largely paralleling those of the total sample. Behavior change scores were significantly higher than all others, but more similar to affect change ($p < .01$) and decision-making ($p < .01$) than analysis ($p < .001$) and normalization ($p < .001$). Affect change and decision-making were rated similarly, and both had higher scores than analysis and normalization. Analysis was rated higher than normalization ($p < .05$).

Like the sample of therapists, within subject t-tests for parents of child clients yielded results similar to those of the entire sample. Behavior change was rated more highly than all other sub-scales, including decision-making ($p < .04$), normalization ($p < .001$), and analysis ($p < .001$), but was similar to affect change. Affect change was rated as more of a priority than both analysis ($p < .001$) and normalization ($p < .001$), while decision-making assistance was considered more important to parents than analysis ($p < .003$) and normalization ($p < .002$). Normalization and analysis were rated comparably.

Comparison of adult clients' sub-scale scores showed no significant differences among them (see table).

Table 1
Goal Objectives Means and Standard Deviations

	Normalization	Analysis	Decision-Making	Behavior Change	Affect Change
All Subject	12.97 (3.82) N=126	13.56 (3.64) N=125	14.24 (3.79) N=127	15.05 (3.68) N=125	14.38 (3.90) N=125
Therapists N=18	11.44 (3.01)	12.67 (3.66)	14.39 (3.68)	16.67 (2.87)	14.89 (3.34)
Parents of Child Clients N=63	14.31 (3.34)	14.33 (3.78)	15.44 (3.52)	16.37 (2.73)	16.02 (3.23)
Adult Clients N=43	11.58 (4.12)	12.79 (3.21)	12.35 (3.54)	12.40 (3.81)	11.74 (3.65)

Discussion

Within subject analyses showed the relative importance of therapeutic objectives of the various respondent groups. When all respondents were combined, what emerged was a view of therapy focused first on behavior change, next on affective improvement, and help with decision-making, with less emphasis on understanding causes of problems and less still on normalizing experiences.

Among therapists, behavior change was the clear treatment priority. Analysis was seen as less important than affect change and decision-making assistance. Therapists saw normalization as least important of all the objectives. The parents of child clients saw behavior change as having priority over the other goal objectives--just as the therapists had rated it. Affect change and decision making, although more important to parents than normalization and analysis, was rated as less important than behavior change. From this information, it would seem that therapists are quite in tune with the parents of their child clients.

Adult clients seemed to value all of the five therapy objectives equally. Unlike therapists and parents of child clients, they appeared to see normalization and causal understanding (analysis) as being as important as symptom reduction and decision-making assistance.

It is possible that today's therapists may spend too much time on the behavior symptom relief of adult clients in an attempt to over correct for all of the years that therapy focused only on the analysis and normalization aspects. It is important for therapist, especially when dealing with adult clients, to avoid overcorrection and allow some time to downplay the need for behavior change.

Future studies are needed with a larger sample of therapists and parents of children who are actual, rather than prospective clients. Few of the parents who completed questionnaires (30%) had actual first-hand experience with therapy. In addition, results from this study analyzed pre-treatment expectations; future studies will need to evaluate the same sample for pre and post-treatment outcomes.

Appendix A

Parent Attitude Questionnaire (PAQ)

Please share your reasons for bringing your child to treatment by responding to the following items. (1=not at all 2=somewhat 3=very true 4=extremely true)

1. ___ I hope therapy will increase my child's understanding of why he/she does things.
2. ___ I hope my child discovers that his/her feelings aren't unusual.
3. ___ I hope my child will feel less anxious and depressed.
4. ___ I want my child to make the right choice about what to do.
5. ___ I want my child to find out if he/she has a serious problem.
6. ___ I want my child to know the reasons behind his/her actions.
7. ___ I hope my child learns how to be more effective in relating to others.
8. ___ I want my child to have more confidence that he/she can do things.
9. ___ I want my child to learn how to make better decisions.
10. ___ I would like my child to understand how the past influences him/her.
11. ___ I would like my child to learn to be more assertive.
12. ___ I want my child to feel more positive about himself/herself.
13. ___ I want my child to learn how to handle situations more competently.
14. ___ I hope my child learns to be more honest with himself/herself.
15. ___ I would like my child to learn how to relax and have more fun.
16. ___ I want my child to learn whether others have problems like his/hers.
17. ___ I want my child to learn how to prevent becoming depressed.
18. ___ I would like my child to receive practical advice about a choice he/she is facing.
19. ___ I hope my child figures out what makes him/her tick.
20. ___ I want my child to get along better with his/her family.
21. ___ I would like my child to know that he/she is not the only one with this problem.
22. ___ I would like my child to receive help deciding what to do with his/her life.
23. ___ I want my child to learn how to keep from getting too angry.
24. ___ I want my child to figure out the course that will make him/her most happy.
25. ___ I want my child to find out if his/her problems are common.
26. ___ I would like my child to receive medication to help him/her function better.
27. ___ I hope therapy will increase my understanding of why he/she does things.
28. ___ I hope to learn how to respond more effectively to my child.
29. Has your child actually been in therapy? _____

Parents' Motivations for Seeking Treatment for their Children

The Parent Attitude Questionnaire (PAQ) was developed to assess the reasons parents seek the help of a psychotherapist for their child. It was designed to measure five different motivations for entering treatment, and is intended for use in both research and as a tool for clinicians in deciding how to structure their treatment.

The following shows the five factors and the items with the highest loadings on each of these factors. The 25 scale items were selected from an original pool of 40, on the basis of a factor analysis conducted on data from a sample of 140 therapy clients obtained over a period of four years.

Factor analytic research can be used to collapse scales, reducing items without any significant loss of information. Shortened forms impose less of a burden on the recipients, and are therefore expected to yield higher response rates. On-site evaluation provides clinicians with immediate feedback on their clients' expectations of treatment.

Normalization

I want my child to find out if she/he has a serious problem.

I hope my child discovers that his/her feelings aren't very unusual.

I want to my child to learn whether others have problems like his/hers.

I would like my child to know that he/she is not the only one going through this.

I want my child to find out if his/her problems are common.

Analysis

I hope therapy will increase my child's understanding of why he/she does things.

I want my child to know the reasons behind his/her actions.

I would like my child to understand how the past influences him/her.

I hope my child figures out what makes him/her tick.

Feelings

I hope my child will feel less anxious and depressed.

I want my child to feel more positive about himself/herself.

I would like my child to learn to how to relax and have more fun.

I want my child to learn how to prevent becoming depressed.

I want my child to learn how to keep from getting too angry.

Decision-Making

I want my child to make the right choice about what to do.

I want my child to learn how to make better decisions.

I would like my child to receive help deciding what to do with his/her life.

I would like my child to receive practical advice about a choice he/she is facing.

I want my child to figure out the course that will make him/her most happy.

Behavior Skills

I want my child to learn how to handle situations more competently.

I want my child to have more confidence that he/she can do things.

I would like my child to learn to be more assertive.

I hope my child learns how to be more effective in relating to others.

I want my child to learn to get along better with his/her family.

Appendix B

Client Attitude Questionnaire (CAQ)

Please share your reasons for coming to counseling by responding to the following items.

(1=not at all 2=somewhat 3=very true 4=extremely true)

1. ___ I hope therapy will increase my understanding of why I do things.
2. ___ I hope to discover that my feelings aren't unusual.
3. ___ I hope to feel less anxious and depressed.
4. ___ I want to be able to make the right choice about what to do.
5. ___ I want to find out if I have a serious problem.
6. ___ I want to know the reasons behind my actions and feelings.
7. ___ I hope to learn how to be more effective in relating to others.
8. ___ I want to have more confidence that I can do things.
9. ___ I need help in learning how to make better decisions.
10. ___ I need help in understanding how my childhood influenced me.
11. ___ I need help in learning how to be more assertive.
12. ___ I want to feel more positive about myself.
13. ___ I want to learn how to handle situations more competently.
14. ___ I hope to learn how to be more honest with myself.
15. ___ I need to learn how to relax and have more fun.
16. ___ I want to learn whether other people have problems like mine.
17. ___ I want to learn how to prevent becoming depressed.
18. ___ I need some practical advice about a choice I'm facing.
19. ___ I hope to figure out what makes me tick.
20. ___ I want to learn how to get along better with my family now.
21. ___ I need to know if I'm the only one going through this.
22. ___ I need help in deciding what to do with my life.
23. ___ I need to learn how to keep from getting too angry.
24. ___ I want to figure out the course that will make me most happy.
25. ___ I want to find out if my problems are common.
26. ___ I need to receive medication to help me function better.
27. ___ Other things I hope to get from treatment: _____

Clients' Motivations for Seeking Treatment

The Client Attitude Questionnaire (CAQ) was developed to assess the reasons clients seek the help of a psychotherapist. It was designed to measure five different motivations for entering treatment, and is intended for use in both research and as a tool for clinicians in deciding how to structure their treatment.

The following shows the five factors and the items with the highest loadings on each of these factors. The 25 scale items were selected from an original pool of 40, on the basis of a factor analysis conducted on data from a sample of 140 therapy clients obtained over a period of four years.

Factor analytic research can be used to collapse scales, reducing items without any significant loss of information. Shortened forms impose less of a burden on the recipients, and are therefore expected to yield higher response rates. On-site evaluation provides clinicians with immediate feedback on their clients' expectations of treatment.

Normalization

I want to find out if I have a serious problem.

I hope to discover that my feelings aren't very unusual.

I want to learn whether other people have problems like mine.

I need to know if I'm the only one going through this.

I want to find out if my problems are common.

Analysis

I hope therapy will increase my understanding of why I do things.

I want to know the reasons behind my actions and feelings.

I need help in understanding how my childhood influenced me.

I hope to figure out what makes me tick.

I hope to learn to be more honest with myself.

Feelings

I hope to feel less anxious and depressed.

I want to feel more positive about myself.

I need to learn how to relax and have more fun.

I want to learn how to prevent becoming depressed.

I need to learn how to keep from getting too angry.

Decision-Making

I want to be able to make the right choice about what to do.

I need help in learning how to make better decisions.

I need help in deciding what to do with my life.

I need some practical advice about a choice I'm facing.

I want to figure out the course that will make me most happy.

Behavior Skills

I hope to learn how to handle situations more competently.

I want to have more confidence that I can do things.

I need help in learning how to be more assertive.

I hope to learn how to be more effective in relating to others.

I want to learn how to get along better with my family now.

REFERENCES

- Abidin, R. R. (1990). *Parenting Stress Index clinical manual*. Charlottesville, VA: Pediatric Psychology Press.
- Achenbach, T.M. (1991). *Manual for the Child Behavior Checklist/4-18 and 1991 Profile*. Burlington: University of Vermont, Department of Psychiatry.
- Achenbach, T.M., & Edelbrock, C. (1987). *Manual for the Youth Self-Report and Profile*. Burlington: University of Vermont, Department of Psychiatry.
- Achenbach, T.M., McConaughy, S.H., & Howell, C.T. (1987). Child/adolescent behavioral and emotional problems: Implications of cross-informant correlations for situational specificity. *Psychological Bulletin*, 101, 213-232.
- American Educational Research Association, American Psychological Association, and National Council on Measurement in Education (1985). *Standards for educational and psychological testing*. Washington, D.C.: American Psychological Association.
- American Psychiatric Association. (1987). *Diagnostic and statistical manual of mental disorders* (3rd ed., rev.) Washington, D.C.: Author.
- American Psychiatric Association. (1987). *Diagnostic and statistical manual of mental disorders* (3rd ed., rev.). Washington, DC: Author.
- Ankuta, G.Y., & Abeles, N. (1993). Client satisfaction, clinical significance, and meaningful change in psychotherapy. *Professional Psychology: Research and Practice*, 24 (1), 70-74.
- Armbruster, P., & Fallon, T. (1994). Clinical, sociodemographic, and systems risk factors for attrition in a children's mental health clinic. *American Journal of Orthopsychiatry*, 64, 577-585.
- Armbruster, P., & Kazdin, A. E. (1994). Attrition in child psychotherapy. In T. H. Ollendick & R. J. Prinz (Eds.), *Advances in clinical child psychology* (Vol.16, pp.81-108). New York: Plenum.
- Attkisson, C.C., & Zwick, R. (1982). The client satisfaction questionnaire: Psychometric properties and correlations with service utilization and psychotherapy outcome. *Evaluation and Program Planning*, 5, 233-237.
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personal and Social Psychology*, 51, 1173-1182.

- Benheim, K. F., & Switalski, T. (1988). Mental health staff and patient's relatives: How they view each other. *Hospital and Community Psychiatry, 39*, 63-68.
- Bickman, L. (1996a). A continuum of care: More is not always better. *American Psychologist, 51* (7), 689-701.
- Bickman, L. (1996b). Methodological issues in evaluating mental health services [Special issue]. *Evaluation and Program Planning, 19* (2).
- Bickman, L. (1997). Reactions to Fort Bragg findings: New directions for mental health services research. *American Psychologist, 52*, 562-565.
- Bickman, L., Guthrie, P.R., Foster, E.M., Lambert, E.W., Summerfelt, W.T., Breda, C.S., & Heflinger, C.A. (1995). *Evaluating managed mental health services: The Fort Bragg experiment*. New York: Plenum.
- Bilbrey, J., & Bilbrey, P. (1995, July/August). Judging, trusting, and utilizing outcomes data: A survey of behavioral health care payers. *Behavioral Health Care Tomorrow*, pp. 62-65.
- Brannan, A.M., Heflinger, C.A., & Bickman, L. (1997). The Caregiver Strain Questionnaire: Measuring the impact of living with a child with serious emotional disturbance. *Journal of Emotional and Behavioral Disorders*.
- Breckler, S. I. (1990). Applications of covariance structure modeling in psychology: Cause for concern? *Psychological Bulletin, 107*, 260-273.
- Brewin, C. R., Andrews, B., & Gotlib, I. H. (1993). Psychopathology and early experience: A reappraisal of retrospective reports. *Psychological Bulletin, 113*, 82-98.
- Byalin, K., KED, J., & Lehmann, S. (1985). Designing and evaluating intervention strategies for deinstitutionalized mental patients. *International Review of Applied Psychology, 34*, 381-390.
- Byalin, K. (1993). Assessing Parental Satisfaction with children's Mental Health Services. *Evaluation and Program Planning, 16*, 69-71.
- Byalin, K. (1990). Parent empowerment: A Treatment strategy for hospitalized adolescents. *Hospital and Community Psychiatry, 41*, 89-90.
- Byrne, B. M. (1989). *A Primer of LISREL: Basic applications and programming for confirmatory factor analytic models*. New York: Springer Verlag.
- Byrne, B. M. (1994). *Structural equation modeling with EQS and EQS/ Windows*. Thousand Oaks, CA: Sage.

- Byrne, B. M., & Goffin, R. D. (1993). Modeling MTMM data from additive and multiplicative covariance structures: An audit of construct validity concordance. *Multivariate Behavioral Research, 28*, 67-96.
- Campbell, D.T., & Fiske, D. W. (1959). Convergent and discriminant validation by the multi-trait multi-method matrix. *Psychological Bulletin, 56*, 81-105.
- Campbell, J., Ho, L., Evenson, R. C., & Bluebird, G. (1996, February). *Consumer satisfaction and treatment outcomes*. Paper presented at the Sixth Annual National Conference on State Mental Health Agency Services Research and Program Evaluation, Arlington, VA.
- Casey, R. J., & Berman, J.S. (1985). The outcome of psychotherapy with children. *Psychological Bulletin, 98* (2), 388-400.
- Cauce, A. M., Dyck, D., Hendryx, M. Srebnik, D. Caverly, S., & Setevenson, J. (1996). *Evaluation of SSB6547 Progress Report: Substituting outcomes for process regulations*. Report prepared for the Washington State Division of Mental Health.
- Chambers, W. J., Puig-Antich, J., Hirsch, M., Paez, P., Ambrosini, P. J., Tabrizi, M. A., & Davies, M. (1985). The assessment of affective disorders in children and adolescents by semistructured interview: Test-retest reliability. *Archives of General Psychiatry, 42*, 696-702.
- Cronbach, L.J., & Furby, L. (1970). How should we measure "change"--or should we? *Psychological Bulletin, 74*, 68-80.
- Daniels, A., Kramer, T. L., & Mahesh, N.M. (1995, July/August). Quality indicators measured by behavioral group practices. *Behavioral Health Care Tomorrow*. Pp.55-56.
- Deane, F. P. (1993). Client satisfaction with psychotherapy in two outpatient clinics in New Zealand. *Evaluation and Program Planning, 16*, 87-94.
- DeChillo, N. (1993). Collaboration of social workers and families of patients with mental illness. *Families in Society, 74*, 104-115.
- DeChillo, N., Koren, P. E., & Schultz, K. H. (1994). From paternalism to partnership: Family and professional collaboration in children's mental health. *American Journal of Orthopsychiatry, 64*(4), 564-576.
- Denner, B., & Halperin, . (1974). Measuring consumer satisfaction in a community outpost. *American Journal of Community Psychology, 2*, 13-22.
- Derogatis, L.R., Rickels, K., & Rock, A (1976). The SCL-90 and the MMPI: Validation of a new self-report scale. *British Journal of Psychiatry, 128*, 280-289.

- Dumas, J.E. (1989). Let's not forget context in behavioral assessment. *Behavioral Assessment, 11*, 231-247.
- Dumas, J.E. (1992). Conduct disorder. In S.M. Turner, K.S. Calhoun, & H.E. Adams (Eds.), *Handbook of clinical behavior therapy* (2nd ed., pp. 285-316). New York: Wiley.
- Edwards, D. W., Yarvis, R. M., Mueller, D. P., & Langsley, D. G. (1978). Does patient satisfaction correlate with success? *Hospital and Community Psychiatry, 29* (3), 188-190.
- Eisen, S. V. (1996, August). Client satisfaction and clinical outcomes: Do we need to measure both? *Behavioral Healthcare Tomorrow*, pp. 71-73.
- Eyberg, S.M., & Johnson, S.M. (1974). Assessment of behavior modification with families: Contingency contracting and order of treated problems. *Journal of Consulting and Clinical Psychology, 42*, 594-606.
- Fauber., R., & Kendall, P.C. (1992). Children and families: Integrating the focus of interventions. *Journal of Psychotherapy Integration, 2*, 107-124.
- Firestone, P., & Witt, J.E. (1982). Characteristics of families completing and prematurely discontinuing a behavioral parent-training program. *Journal of Pediatric Psychology, 7*, 209-222.
- Forehanc, R., Middlebrook, J., Rogers, T., & Steffe, M. (1983). Dropping out of parent training. *Behavior Research and Therapy, 21*, 663-668.
- Garfield, S. L., Prager, R. A., & Bergin, A. E. (1971). Evaluation of outcome in psychotherapy. *Journal of Consulting and Clinical psychology, 37* (3), 307-313.
- Gilligan, J.F., & Wilderman, M.A. (1977). An economical rural mental health consumer satisfaction evaluation. *Community Mental Health Journal, 13*, 31-36.
- Goode, Erica E. (May 1993). Does Psychotherapy work? *U.S. News and World Report, 4*, 56+.
- Gould, M.S., Shaffer, D., & Kaplan, D. (1985). The characteristics of dropouts from a child psychiatry clinic. *Journal of the American Academy of Child Psychiatry, 24*, 316-328.
- Goyne, J. & Ladoux, P. (1973). Patients' opinions of outpatient clinic services. *Hospital and Community Psychiatry, 24*, 627-628.
- Grunebaum, H. (1984). Comments on Turkelson's "Schizophrenia and the family: II. Adverse effects of family therapy." *Family Processes, 23*, 421-428.

- Hatfield, A. B. (1982). Commentary: Therapists and families: Worlds apart. *Hospital and Community Psychiatry, 33*, 513.
- Hatfield, A.B. (1987) Families as caregivers: A historical perspective. In A.B. Hatfield & H.P. Lefley (Eds.), *Families of the mentally ill: Coping and adaptation*. New York: The Guilford Press.
- Hoagwood, K., Hibbs, E., Brent, D., & Jensen, P. (1995). Introduction to the special edition: Efficacy and effectiveness studies of child and adolescent psychotherapy. *Journal of Consulting and Clinical Psychology, 63*, 683-687.
- Hodges, K. (1990). *The Child and Adolescent Functional Assessment Scale (CAFAS)*. Unpublished manuscript.
- Hodges, K., & Gust, J. (1995). Measures of impairment for children and adolescents. *Journal of Mental Health Administration, 22(4)*, 403-413.
- Hodges, K., Kline, J., Stern, L., Cytryn, L., & McKnew, D. (1982). The development of a child assessment interview for research and clinical use. *Journal of Abnormal Child Psychology, 10*, 173-189.
- Hodges, K., McKnew, D., Cytryn, L., Stern, L., & Kline, J. (1982). The Child Assessment Schedule (CAS) Diagnostic Interview: A report on reliability and validity. *Journal of the American Academy of Child and Adolescent Psychiatry, 21*, 468-473.
- Holcomb, W.R., Adams, N.A. Ponder, H.M., & Reitz, R. (1989). The development an construction of a consumer satisfaction questionnaire for psychiatric inpatients. *Evaluation and Program Planning, 12*, 189-194.
- Holden, D.F., & Lewine, R.R.J. (1982). How families evaluate mental health professionals, resources, and effects of illness. *Schizophrenia Bulletin, 8*, 636-633.
- Hollingshead, A. B., & Redlich, F. C. (1958). *Social class and mental illness*. New York: Wiley.
- Horne, A.M., & Patterson, G.R. (1980). Working with parents of aggressive children. In R.R. Abidin (Ed), *Parent education and intervention handbook*. Springfield, Il: Charles C. Thomas.
- Howard, K. I., Krause, M. S., & Orlinsky, D. E. (1986). The attrition dilemma: Toward a new strategy for psychotherapy research. *Journal of Consulting and Clinical Psychology, 54*, 106-110.
- Jacobson, N. S., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and*

Clinical Psychology, 59 (1), 12 - 19.

- Kaplan, E. L., & Meier, P. (1958). Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association*, 53, 457-481.
- Kazdin, A. E. (1996b). Problem solving and parent management in treating aggressive and antisocial behavior. In E. D. Hibbs & P.S Jensen (Eds.), *Psychosocial treatments for child and adolescent disorders: Empirically based strategies for clinical practice* (pp.377-408). Washington, DC: American Psychological Association.
- Kazdin, A. E. (1989). Hospitalization of antisocial children: Clinical course, follow-up status, and predictors of outcome. *Advances in Behaviour Research and Therapy*, 11, 1-67.
- Kazdin, A. E. (1992). *Research design in clinical psychology* (2nd ed.). Needham Heights, MA: Allyn & Bacon.
- Kazdin, A. E. (1994). Psychotherapy for children and adolescents. In A. E. Bergin & S. L. Garfield (Eds.), *Handbook of psychotherapy and behavior change*. New York: Wiley.
- Kazdin, A. E. (1995). Child, parent, and family dysfunction as predictors of outcome in cognitive-behavioral treatment of antisocial children. *Behaviour Research and Therapy*, 33, 271-281.
- Kazdin, A. E. (1996a). Dropping out of child therapy: Issues for research and implications for practice. *Clinical Child Psychology and Psychiatry*, 1, 133-156.
- Kazdin, A. E., & Crowley, M. (in press). Moderators of treatment outcome in cognitively based treatment of antisocial behavior. *Cognitive Therapy and Research*.
- Kazdin, A. E., & Mazurick, J. L. (1994). Dropping out of child psychotherapy: Distinguishing early and late dropouts over the course of treatment. *Journal of Consulting and Clinical Psychology*, 62, 1069-1074.
- Kazdin, A. E., Holland, L., & Breton, S. (1991). *Barriers to Participation in Treatment Scale-Parent and Therapist Versions*. New Haven, CT: Yale University.
- Kazdin, A. E., Holland, L., & Crowley, M. (1997). Family experience of barriers to treatment and premature termination from child therapy. *Journal of Consulting and Clinical Psychology*, 65, 453-463.
- Kazdin, A. E., Mazurick, J. L., & Bass, D. (1993). Risk for attrition in treatment of antisocial children and families. *Journal of Clinical Child Psychology* 22, 2-16.
- Kazdin, A. E., Siegel, T., & Bass, D. (1990). Drawing upon clinical practice to inform

- research on child and adolescent psychotherapy. *Professional Psychology: Research and Practice*, 21, 189-198.
- Kazdin, A. E., Siegel, T., & Bass, D. (1992). Cognitive problem-solving skills training and parent management training in the treatment of antisocial behavior in children. *Journal of Consulting and Clinical Psychology*, 60, 733-747.
- Kazdin, A.E. (1990). Premature termination from treatment among children referred for antisocial behavior. *Journal of Child Psychology and Psychiatry*, 31, 415-425.
- Kazdin, A.E. (1990). Treatment of antisocial behavior in children: Current status and future directions. *Psychological Bulletin*, 102, 187-203.
- Kazdin, A.E., Siegel, T.C., & Bass, D. (1992). Cognitive problem-solving skills training and parent management training in the treatment of antisocial behavior in children. *Journal of Consulting and Clinical Psychology*, 60, 733-747.
- Kendall, P.C., & Morris, R. (1991). Child therapy: Issues and recommendations. *Journal of Consulting and Clinical Psychology*, 59, 777-784.
- Kreisman, D.E., & Joy, V.D. (1974). Family response to the mental illness of a relative: A review of the literature. *Schizophrenia Bulletin*, 1, 34-57.
- Lambert, W., Salzer, Mark S., & Bickman, Leonard. (1998). Clinical outcome, consumer satisfaction, and ad hoc ratings of improvement in children's mental health. *Journal of Consulting and Clinical Psychology*, 66, 270-279.
- Larsen, D.L. Attkisson, C.C. Harreaves W.A., & Nguyen, T.D. (1979) Assessment of client/patient satisfaction: Development of a general scale. *Evaluation and Program Planning*, 2, 197-207.
- Lebow, J. (1982). Consumer satisfaction with mental health treatment. *Psychological Bulletin*, 91, 244-259.
- Lebow, J.L. (1983). Research assessing consumer satisfaction with mental health treatment: A review of findings. *Evaluation and Program Planning*, 6, 211-236.
- LeVois, M., Nguyen, T & Attkisson, C. (1981). Artifact in client satisfaction assessment: Experience in community mental health settings. *Evaluation and Program Planning*, 4, 139-150.
- Lieberman, H.J., Nigro, J., Tremabath, P., Tenczynski, D., & Scher, S. (1988). Community mental health systems generate new chronic: The study of a pure case. *Administration in Mental Health*, 15, 139-156.
- McElroy, E. M. (1987). The beat of a different drummer. In A. B. Hatfield & H. P.

- Lefley (eds.), *Families of the mentally ill: Coping and adaptation* (pp. 225-243). New York: Guilford Press.
- McLellan, A. T., & Hunkeler, E. (in press). Is patient satisfaction related to patient outcome in alcohol and drug abuse treatment? *Psychiatric Services*.
- McMahon, R., Forehand, R., Griest, D., & Wells, K. (1981). Who drops out of therapy during parent behavioral training? *Behavioral Counseling Quarterly*, 1, 79-85.
- Mental health: Does therapy help? (1995, November). *Consumer Reports*, pp.734-739.
- Miller, G.E., & Prinz, R.J. (1990). The enhancement of social learning family interventions for childhood conduct disorder. *Psychological Bulletin*, 108, 291-307.
- Miller, G.E., & Prinz, R.J., (1994). Family based treatment for childhood antisocial behavior: Experimental influences on dropout and engagement. *Journal of Consulting and Clinical Psychology*, 62, 645-650.
- Minsky, S. (1997, February). *The relationship between a self-report quality of life instrument and a consumer satisfaction survey*. Paper presented at the Seventh Annual National Conference on State Mental Health Agency Services Research and Program Evaluation, Arlington, VA.
- Newman, F. L. (1994). Criteria for selecting statistical procedures for progress and outcome assessment. In M. E. Maruish (Ed.). *Use of psychological testing for treatment planning and outcome assessment* (pp.111-134). Hillsdale, NJ: Erlbaum.
- Nguyen, T.D., Attkisson, C.C., & Stegner, B.L. (1983). Assessment of patient satisfaction: Development and refinement of service evaluation questionnaire. *Evaluation and Program Planning*, 6, 299-313.
- Northrup, D. A. (1995). *The relationship between adolescent and parent reports about family issues*. Unpublished doctoral dissertation. Vanderbilt University, Nashville, TN.
- Patterson, G.R. (1986). Performance models for antisocial boys. *American Psychologist*, 41, 432-444.
- Patterson, G.R., Reid, J.B., & Dishion, T.J. (1992). *Antisocial boys*. Eugene, OR: Castalia.
- Pekarik, G., & Wolff, C. B. (1996). Relationship of satisfaction to symptom change, follow-up adjustment, and clinical significance. *Professional Psychology: Research and Practice*, 27 (2), 202-208.
- Prinz, R. J., & Miller, G.E. (1994). Family-based treatment for childhood antisocial behavior: Experimental influences on dropout and engagement. *Journal of Consulting*

and *Clinical Psychology*, 62, 645-650.

- Reynolds, K., Ogiba, S., & Chambliss, C. (1998) Educating therapists in training about clients' expectations of treatment. *Resources in Education*. ERIC/CASS. CGO28449.
- Ries, R. K., Jaffe, C., Comtois, K. A., & Kitchell, M. (1996). *Patient satisfaction and outcome with outpatient integrated dual diagnosis treatment*. Unpublished manuscript.
- Robins, L. N. (1981). Epidemiological approaches to natural history research: Antisocial disorders in children. *Journal of the American Academy of Child Psychiatry* 20, 566-680.
- Ross, E. C. (1997). Managed behavioral health care premises, accountable systems of care, and AMBHA's PERMS. *Evaluation Review*, 21, 318-321.
- Rutter, M. (1987). Psychosocial resilience and protective mechanisms. *American Journal of Orthopsychiatry* 57, 316-331.
- Salzer, M. S. (1997). Consumer empowerment in mental health organizations: Concepts, benefits, and impediments. *Administration and Policy in Mental Health*, 24, 425-434.
- Sandler, I.N., & Barrera, M. (1984). Toward a multimethod approach to assessing the effects of social support. *American Journal of Community Psychology*, 12, 37-52.
- Santisteban, D. A., Szapocznik, J., Perez-Vidal, A., Kurtines, W. H., Murray, E. J., & LaPerriere, A. (1996). Efficacy of intervention for engaging youth and families into treatment and some variables that may contribute to differential effectiveness. *Journal of Family Psychology*, 10, 35-44.
- Sarkis, L., Dlugacz, D.P., & Wilner, M. (1987). A unique urban state hospital: Partial-hospital programs reduce full-time bed occupancy. *International Journal of Partial Hospitalization*, 4, 157-165.
- Schainblatt, A.H. (1980). What happens to clients? *Community Mental Health Journal*, 16, 330-342.
- Seligman, M. E. P. (1995). The effectiveness of psychotherapy: The *Consumer Reports* study. *American Psychologist*, 50(12), 965-974.
- Silver, L. B., & Silver, B. J. (1983). Clinical practice of child psychiatry: A survey. *Journal of the American Academy of Child Psychiatry*, 22, 573-579.
- Smets, A. C. (1982). Family and staff attitude toward family involvement in the treatment of hospitalized chronic patients. *Hospital and Community Psychiatry*, 33, 573-575.
- Smith, M.L., Glass, G. V., & Miller, T. I. (1980). *Psychotherapy*. Baltimore, MD: Johns

Hopkins University Press.

Sorenson, J., Kantor, L., Margolis, R., & Galano, J. (1979). The extent, nature and utility of evaluating consumer satisfaction in community mental health centers. *American Journal of Community Psychology*, 7, 329-337.

Soley, G., Hooper, Y., Marshall, R., & Chambliss, C. (1998). A comparison of parents' and therapists' expectations of child psychotherapy. *Unpublished manuscript*.

Spanier, G.B., & Thompson, L. (1982). A confirmatory analysis of the Dyadic Adjustment Scale. *Journal of Marriage and the Family*, 44, 731-738.

Spaniol, L., Jung, H., Zipple, A., & Fitzgerald, S. (1987). Families as a resource in rehabilitation of the severely psychiatrically disabled. In A. B. Hatfield & H. P. Lefley (eds.), *Families of the mentally ill: Coping and adaptation* (pp. 167-190). New York: Guilford Press.

Spaniol, L., Zipple, A., & Fitzgerald, S. (1984). How professionals can share power with families: Practical approaches to working with families of the mentally ill. *Psychosocial Rehabilitation Journal*, 8(2), 77-84.

Szapocznik, J., Perez-Vidal, A., Brickman, A., Foote, F. H., Santisteban, D. A., Hervis, O., & Kurtines, W. H. (1988). Engaging adolescent drug abusers and their families into treatment: A strategic structural systems approach. *Journal of Consulting and Clinical Psychology*, 56, 552-557.

Tanner, B.A., & Stacy, W. (1981). Factors influencing client satisfaction with mental health services: A review of quantitative research. *Evaluation and Program Planning*, 4, 279-286.

Tanner, B.A., & Stacy, W. (1985). A validity scale for the sharp consumer satisfaction scales. *Evaluation and Program Planning*, 8, 147-153.

Tarling, R., & Perry, J. (1985). Statistical methods in criminological prediction. In D. P. Farrington & R. Tarling (Eds.), *Prediction in criminology* (pp. 210-231). Albany: State University of New York Press.

Trabin, T., Freeman, M.A., & Pallak, M.S. (Eds.). (1995). *Inside outcomes: The national review of behavioral health care outcome programs* (Revised and expanded edition). Tiburon, CA: Centralink Publications.

U.S. Congress, Office of Technology Assessment. (1991). *Adolescent health* (OTA-H-468). Washington, DC: U.S. Government Printing Office.

U.S. Congress, Office of Technology Assessment. (1986). *Children's mental health: Problems and services-A background paper* (OTA-BP-H-33). Washington, DC: U.S.

Government Printing Office.

- Wechsler, D. (1974). *Manual for the Wechsler Intelligence Scale for Children-Revised*. New York: Psychological Corporation.
- Weiss, I., Rabinowitz, J., & Spiro, S. (1996). Agreement between therapists and clients in evaluating therapy and its outcomes: Literature review. *Administration and Policy in Mental Health*, 23, 493-511.
- Weisz, J. R., Donenberg, G. R., Han, S. S., & Weiss, B. (1995). Bridging the gap between laboratory and clinic in child and adolescent psychotherapy. *Journal of Consulting and Clinical Psychology*, 63 (5), 688-701.
- Weisz, J. R., Weiss, B., & Donenberg, G. R. (1992). The lab versus the clinic: Effects of child and adolescent psychotherapy. *American Psychologist*, 47, 1578-1585.
- Weisz, J. R., Weiss, B., Alicke, M.D., & Klutz, M. L. (1987). Effectiveness of psychotherapy with children and adolescents: A meta-analysis for clinicians. *Journal of Consulting and Clinical Psychology*, 55 (4), 542-549.
- Weisz, J.R., Weiss, B., & Langmeyer, D.B. (1987). Giving up on child psychotherapy: Who drops out? *Journal of Consulting and Clinical Psychology*, 55, 916-918.
- Wierzbicki, M., & Pekarik, G. (1993). A meta-analysis of psychotherapy dropout. *Professional Psychology: Research and Practice*, 24, 190-195.
- Williams, B. (1994). Patient satisfaction: A valid concept? *Social Science and Medicine*, 38 (4), 509-516.
- Woodard, S., Santa-Barbara, J., Levin, S., & Epstein, N. (1978). Aspects of consumer satisfaction with brief family therapy. *Family Process*, 17, 399-407.



U.S. Department of Education
Office of Educational Research and Improvement (OERI)
Educational Resources Information Center (ERIC)



REPRODUCTION RELEASE

(Specific Document)

I. DOCUMENT IDENTIFICATION:

Title: <i>Training Therapists about Client Expectations of Psychotherapy</i>	
Author(s): <i>Soley, G., Marshall, R., + Chambliss, C.</i>	
Corporate Source: <i>Ursinus College</i>	Publication Date: <i>1998</i>


II. REPRODUCTION RELEASE:

In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, *Resources in Education* (RIE), are usually made available to users in microfiche, reproduced paper copy, and electronic/optical media, and sold through the ERIC Document Reproduction Service (EDRS) or other ERIC vendors. Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to the document.

If permission is granted to reproduce and disseminate the identified document, please CHECK ONE of the following two options and sign at the bottom of the page.

The sample sticker shown below will be affixed to all Level 1 documents

The sample sticker shown below will be affixed to all Level 2 documents



PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY

Sample

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN OTHER THAN PAPER COPY HAS BEEN GRANTED BY

Sample

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)



Check here
For Level 1 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical) and paper copy.

Check here
For Level 2 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical), but not in paper copy.

Level 1

Level 2

Documents will be processed as indicated provided reproduction quality permits. If permission to reproduce is granted, but neither box is checked, documents will be processed at Level 1.

"I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate this document as indicated above. Reproduction from the ERIC microfiche or electronic/optical media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries."

Sign here → please

Signature: <i>Catherine Chambliss</i>	Printed Name/Position/Title: <i>Catherine Chambliss, Ph.D., Chair, Psychology</i>
Organization/Address: <i>Dept. of Psychology Ursinus College Collegeville, PA 19426</i>	Telephone: <i>(610) 409 3000</i>
	FAX: <i>(610) 489 0627</i>
	E-Mail Address: <i>cchambliss@acad.ursinus.edu</i>
	Date: <i>8/28/98</i>

III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of the document from another source, please provide the following information regarding the availability of the document. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:
Address:
Price:

IV. REFERRAL OF ERIC TO COPYRIGHT/REPRODUCTION RIGHTS HOLDER:

If the right to grant reproduction release is held by someone other than the addressee, please provide the appropriate name and address:

Name:
Address:

V. WHERE TO SEND THIS FORM:

Send this form to the following ERIC Clearinghouse:

However, if solicited by the ERIC Facility, or if making an unsolicited contribution to ERIC, return this form (and the document being contributed) to:

ERIC Processing and Reference Facility
1301 Piccard Drive, Suite 100
Rockville, Maryland 20850-4305

Telephone: 301-258-5500

FAX: 301-948-3695

Toll Free: 800-799-3742

e-mail: ericfac@lnet.ed.gov