

DOCUMENT RESUME

ED 462 258

SE 060 565

TITLE Physics 30: Grade 12 Diploma Examination, June 1997.  
INSTITUTION Alberta Dept. of Education, Edmonton. Student Evaluation Branch.  
PUB DATE 1997-06-00  
NOTE 43p.  
AVAILABLE FROM Alberta Education, Student Evaluation Branch, 11160 Jasper Avenue, Edmonton, Alberta T5K 0L2, Canada.  
PUB TYPE Tests/Questionnaires (160)  
EDRS PRICE MF01/PC02 Plus Postage.  
DESCRIPTORS Foreign Countries; \*Grade 12; High Schools; \*Physics; \*Science Tests; Scientific Concepts  
IDENTIFIERS \*Alberta Grade Twelve Diploma Examinations

ABSTRACT

This document is a Physics 30 Grade 12 Diploma Examination from Alberta Education. It is a 2.5 hour closed-book examination consisting of 37 multiple-choice and 12 numerical-response questions of equal value worth 70% of the examination and two written-response questions of equal value worth 30% of the examination. The exam contains sets of related questions that may contain multiple-choice and/or numerical-response and/or written response questions. The exam booklet also contains a tear-out data sheet, a Periodic Table of Elements, and blank perforated pages for rough work. (JRH)

Reproductions supplied by EDRS are the best that can be made from the original document.

June 1997



# Physics 30

## Grade 12 Diploma Examination

PERMISSION TO REPRODUCE AND  
DISSEMINATE THIS MATERIAL  
HAS BEEN GRANTED BY

*C. Andrews*

TO THE EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC)

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

This document has been reproduced as  
received from the person or organization  
originating it.

Minor changes have been made to  
improve reproduction quality.

Points of view or opinions stated in this  
document do not necessarily represent  
official OERI position or policy.

BEST COPY AVAILABLE

Alberta  
EDUCATION

590565

Copyright 1997, the Crown in Right of Alberta, as represented by the Minister of Education, Alberta Education, Student Evaluation Branch, 11160 Jasper Avenue, Edmonton, Alberta T5K 0L2. All rights reserved. Additional copies may be purchased from the Learning Resources Distributing Centre.

**Special permission** is granted to **Alberta educators only** to reproduce, for educational purposes and on a non-profit basis, parts of this examination that do **not** contain excerpted material **only after the administration of this examination**.

Excerpted material in this examination **shall not** be reproduced without the written permission of the original publisher (see credits page, where applicable).

*June 1997*

# *Physics 30*

## *Grade 12 Diploma Examination*

### *Description*

Time: 2.5 h. You may take an additional 0.5 h to complete the examination.

Total possible marks: 70

This is a **closed-book** examination consisting of

- 37 multiple-choice and 12 numerical-response questions, of equal value, worth 70% of the examination
- 2 written-response questions, worth a total 30% of the examination

This examination contains sets of related questions. A set of questions may contain multiple-choice and/or numerical-response and/or written-response questions.

A tear-out data sheet is included near the back of this booklet. A Periodic Table of the Elements is also provided.

The blank perforated pages at the back of this booklet may be torn out and used for your rough work. No marks will be given for work done on the tear-out pages.

### *Instructions*

- Fill in the information required on the answer sheet and the examination booklet as directed by the presiding examiner.
- You are expected to provide your own scientific calculator.
- Use only an HB pencil for the machine-scored answer sheet.
- If you wish to change an answer, erase **all** traces of your first answer.
- Consider all numbers used in the examination to be the result of a measurement or observation.
- Do not fold the answer sheet.
- The presiding examiner will collect your answer sheet and examination booklet and send them to Alberta Education.
- Read each question carefully.
- Now turn this page and read the detailed instructions for answering machine-scored and written-response questions.





1. The physical quantity that can have the same unit as impulse is
  - A. force
  - B. work
  - C. power
  - D. momentum

*Use the following information to answer the next two questions.*

A 5.00 kg object is dropped from a height above the ground. When the object is 4.00 m from the ground, it has a speed of 9.00 m/s. The potential energy of the object is chosen to be zero at ground level and the effects of air resistance are ignored.

2. What is the total mechanical energy of the falling object?
  - A. 6.30 J
  - B. 196 J
  - C. 202 J
  - D. 399 J

### **Numerical Response**

*Use your recorded answer for **Multiple Choice 2** to solve **Numerical Response 1**.*

- 1.** The object was dropped from an initial height of \_\_\_\_\_ m above the ground.  
(Round and record your answer to three digits.)

3. A space shuttle astronaut has a mass of 110 kg with her space suit on. She is on a space walk and picks up a full can of spray with a mass of 20 kg. Relative to the space shuttle, she is at rest. She then holds the can directly in front of her centre of mass to avoid rotation and releases 3.0 kg of spray at a speed of 15 m/s. Her speed, relative to the space shuttle, when she has stopped spraying is approximately
- A. 0.35 m/s
  - B. 0.41 m/s
  - C. 2.3 m/s
  - D. 2.5 m/s

*Use the following information to answer the next question.*

A popular game of young children is to shuffle across a carpet with stocking feet and then touch a friend. The spark that can be generated is caused by a charge buildup from the friction of the socks on the carpet.

4. Two friends, Sam and Jeff, shuffled on a carpet and obtained approximately the same negative charge. They then stood shoulder to shoulder without touching. A third friend, Cale, who was not charged, touched Jeff on the shoulder farthest from Sam. What is the nature of the final charges on the three boys?
- A. Sam, Jeff, and Cale are all negatively charged.
  - B. Jeff and Cale are uncharged, and Sam is negatively charged.
  - C. Sam and Cale are negatively charged, and Jeff is positively charged.
  - D. Sam is negatively charged, and Jeff and Cale are positively charged.
- 
5. Three pithballs hang in an isolated container. Ball **X** has a charge of  $1.0 \times 10^{-9}$  C, and balls **Y** and **Z** are neutral. Ball **X** is brought momentarily into contact with ball **Y**, then separated. Ball **Y** is then brought momentarily into contact with ball **Z**, then separated. When placed 1.0 m apart, balls **X** and **Z** will now exert a force on each other of magnitude
- A.  $1.0 \times 10^{-9}$  N
  - B.  $1.1 \times 10^{-9}$  N
  - C.  $2.2 \times 10^{-9}$  N
  - D.  $9.0 \times 10^{-9}$  N




### Numerical Response

2. Two charged bodies exert electrostatic forces on each other of magnitude  $1.11 \times 10^{-4}$  N. If the magnitude of each charge is doubled and the distance separating them is doubled, then the magnitude of the electrostatic force, expressed in scientific notation, is  $b \times 10^{-w}$  N. The value of  $b$  is \_\_\_\_\_. (Round and record your answer to three digits.)

### Numerical Response


3. In moving an electric charge of 4.00 C from point X to point Y, 15.0 J of work is done. The potential difference between X and Y, in volts, is \_\_\_\_\_. (Round and record your answer to three digits.)

Use the following information to answer the next question.



6. Which of the following statements best describes the diagram above?
- A. Circuit Y dissipates more power than does circuit X.
  - B. The current in circuit Y is larger than the current in circuit X.
  - C. The current in circuit Y is the same as the current in circuit X.
  - D. The current in circuit Y is smaller than the current in circuit X.

Use the following information to answer the next question.



7. An electron is placed at point  $P$ . It will accelerate toward region
- A. I
  - B. II
  - C. III
  - D. IV
- 
8. The volt is the SI unit of potential difference. An equivalent SI unit may be written as
- A. J/A
  - B. J/C
  - C. N/C
  - D. A/ $\Omega$

9. Which of the following is a definition of **conventional** direct current?
- A. A movement of negative charge in one direction only
  - B. A movement of positive charge in one direction only
  - C. A shift of negative charge that reaches a peak in the forward direction before reversing and reaching a peak in the reverse direction
  - D. A shift of positive charge that reaches a peak in the forward direction before reversing and reaching a peak in the reverse direction

*Use the following information to answer the next two questions.*

*Torpedo occidentalis* is a large electric fish that uses electricity in attack and defense. A typical individual fish is capable of producing potential differences of up to 220 V and of generating pulses of 15.0 A current through its seawater environment. Pulses are typically  $2.00 \times 10^{-3}$  s in duration.


10. The total charge transferred by the fish in one of these pulses is
- A.  $3.00 \times 10^{-2}$  C
  - B.  $4.40 \times 10^{-1}$  C
  - C.  $3.00 \times 10^3$  C
  - D.  $3.30 \times 10^3$  C

### **Numerical Response**

4. The maximum electrical work done during one pulse is \_\_\_\_\_ J.  
(Round and record your answer to three digits.)

11. If the resistance of a circuit is halved and the voltage applied to the circuit is doubled, then the current in the circuit is
- A. the same
  - B. quartered
  - C. doubled
  - D. quadrupled

Use the following information to answer the next four questions.



12. When the switch is closed, the above circuit can be correctly described as
- A. two series lights, in series with the two outlets
  - B. two parallel lights, in series with the two outlets
  - C. two series lights, in parallel with the two outlets
  - D. two parallel lights, in parallel with the two outlets

13. A  $1.00 \times 10^3$  W toaster is plugged into one outlet of the circuit and switched on. Both lights are on. The **maximum** power rating for a kettle that could be plugged into the other outlet and switched on without burning out the fuse is
- A.  $7.00 \times 10^2$  W
  - B.  $8.00 \times 10^2$  W
  - C.  $1.00 \times 10^3$  W
  - D.  $1.50 \times 10^3$  W


### Numerical Response

5. When a  $1.00 \times 10^3$  W toaster is plugged into one of the outlets, the current in the toaster is \_\_\_\_\_ A.  
(Round and record your answer to three digits.)

### Numerical Response

6. At a rate of  $6.71\text{¢}/(\text{kW}\cdot\text{h})$ , the cost of operating the  $1.00 \times 10^3$  W toaster for 1.10 minutes a day for 30 days is \_\_\_\_\_ ¢.  
(Round and record your answer to three digits.)

Use the following information to answer the next three questions.



14. Why do ions of only a certain speed pass through the velocity selection chamber undeflected?
- A. Only these ions possess the charge needed to be undeflected by the fields.
  - B. The electric field strength is the same as the magnetic field strength.
  - C. Ions travelling at other speeds have insufficient  $E_k$  to pass through the chamber.
  - D. The net deflecting force, from the electric and magnetic fields, is zero for only these ions.

Use the following additional information to answer the next two questions.

The biochemist has the spectrometer set as follows:

- Velocity selection chamber:  $|\vec{E}| = 2.17 \times 10^4 \text{ V/m}$   
 $B_{\perp} = 9.00 \times 10^{-3} \text{ T}$
- Ion separation chamber:  $B_{\perp} = 1.40 \text{ T}$   
deflecting radius = 1.00 m

At these settings, an ion is detected. The biochemist expects the ion to be one of the ions listed below. The mass corresponding to each ion is given.

$\text{Cr}^{2+}$	$8.64 \times 10^{-26} \text{ kg}$
$\text{Cd}^{2+}$	$1.86 \times 10^{-25} \text{ kg}$
$\text{Hg}^{2+}$	$3.33 \times 10^{-25} \text{ kg}$
$\text{Pb}^{2+}$	$3.44 \times 10^{-25} \text{ kg}$


15. Which of the above pollutants is detected by the spectrometer?
- A.  $\text{Cr}^{2+}$
  - B.  $\text{Cd}^{2+}$
  - C.  $\text{Hg}^{2+}$
  - D.  $\text{Pb}^{2+}$

Use the following additional information to answer the next question.

In the ion generation and acceleration chamber, atoms in the sample are ionized by bombarding them with electrons to remove outermost electrons. The biochemist must have the accelerating voltage in the electron gun set high enough to ensure ionization of the particles.

16. Which of the following physical principles must be used to calculate the value of the accelerating voltage in the electron gun?
- A. Ohm's law
  - B. Coulomb's law
  - C. Conservation of energy
  - D. Conservation of momentum

Use the following information to answer the next two questions.



17. If the maximum recommended input current for the motor is 300 A, the approximate time at which shutdown will occur if the motor is using 150 A is
- A. never
  - B. 4.00 min
  - C. 12.00 min
  - D. immediately
18. If the same motor shuts down at 8.00 min, the current before shutdown is approximately
- A. 150 A
  - B. 200 A
  - C. 600 A
  - D. 900 A




Use the following information to answer the next question.

In many electrically powered passenger trains, the input voltage  $V_i$  from the power supply is not the same as the operating voltage  $V_o$  of the electrical circuitry of the train.

Examples:

	$V_i$	$V_o$
England	750	1 500
English Channel	25 000	1 500
Belgium	3 000	1 500
France	50 000	1 500

The diagram below is a partial schematic of the electrical circuitry of an electric train.



19. The component labelled A in the diagram is **most likely** a


- A. battery
- B. resistor
- C. generator
- D. transformer

**Numerical Response**


7. A typical television set requires  $2.00 \times 10^4$  V AC for its operation. Since a television is plugged into a standard 110 V outlet, the voltage must be increased. If the ideal transformer used to increase the voltage has  $1.87 \times 10^4$  turns of wire on the secondary coil, then the number of turns of wire that must be placed on the primary coil, expressed in scientific notation, is  $b \times 10^w$ . The value of  $b$  is \_\_\_\_\_.  
(Round and record your answer to three digits.)

20. The diagrams below show the direction of a magnetic field relative to a set of coordinate axes. A negatively charged particle travels across the page in the positive  $x$  direction. The magnetic configuration that will cause the particle to bend in the positive  $z$  direction is


A.




B.



C.



D.



### Numerical Response

8. Northern lights are often observed in Alberta skies. The most common colour, green, has a wavelength of 558 nm. When a collision occurs between energetic electrons and oxygen atoms in the upper atmosphere, the oxygen atoms are excited. To cause the most common colour of northern lights, the electrons must be travelling with a minimum speed, expressed in scientific notation, of  $b \times 10^w$  m/s. The value of  $b$  is \_\_\_\_\_.  
(Round and record your answer to three digits.)

21. Accelerating charges generate
- A. electric waves
  - B. magnetic waves
  - C. longitudinal waves
  - D. electromagnetic waves

### Numerical Response

9. If a photon of electromagnetic radiation has a frequency of  $1.09 \times 10^{17}$  Hz, it has a wavelength, expressed in scientific notation, of  $b \times 10^{-w}$  m. The value of  $b$  is \_\_\_\_\_.  
(Round and record your answer to three digits.)

## Numerical Response

10. An explosion that produces a flash of light occurs at a distance of 6.06 km from a group of people. The minimum possible time, expressed in scientific notation, that elapses before the people can see the explosion is  $a.bc \times 10^{-d}$  s. The values of  $a$ ,  $b$ ,  $c$ , and  $d$  are \_\_\_\_\_.  
(Record your answer as    .)


22. Which of the following sets of electromagnetic radiations is arranged in order of increasing photon frequency?
- A. Gamma rays, ultraviolet radiation, radio waves
  - B. Radio waves, ultraviolet radiation, gamma rays
  - C. Gamma rays, radio waves, ultraviolet radiation
  - D. Radio waves, gamma rays, ultraviolet radiation

Use the following information to answer the next three questions.

A cyclotron is a particle accelerator used to investigate subatomic structure. Magnetic fields are used to control the path of charged particles within a cyclotron.

23. The radius of the path followed by charged particles moving perpendicularly through the magnetic field of a cyclotron could be reduced by
- A. increasing the strength of the magnetic field
  - B. using particles with a smaller charge
  - C. increasing the speed of the particles
  - D. using particles with a greater mass
24. The period  $T$  for a particle of charge  $q$  in a magnetic field of strength  $B$  is
- A.  $\frac{2\pi m}{qB}$
  - B.  $\frac{\pi m}{qB}$
  - C.  $\frac{qB}{2\pi}$
  - D.  $\frac{qB}{\pi m}$
25. An alpha particle travels in a direction perpendicular to a magnetic field of strength 1.6 T. If the alpha particle experiences a force of magnitude  $1.1 \times 10^{-13}$  N, then its measured speed will be
- A.  $2.1 \times 10^{-7}$  m/s
  - B.  $4.3 \times 10^{-7}$  m/s
  - C.  $2.1 \times 10^5$  m/s
  - D.  $4.3 \times 10^5$  m/s

Use the following information to answer the next question.



### Numerical Response


- 11.** A mass with a +1.00 elementary charge placed between the plates will experience an electric force, expressed in scientific notation, of magnitude  $b \times 10^{-w}$  N. The value of  $b$  is \_\_\_\_\_.  
(Round and record your answer to three digits.)
- \_\_\_\_\_

- 26.** X-rays may be focused using
- A. magnetic fields
  - B. electric fields
  - C. either electric or magnetic fields
  - D. neither electric nor magnetic fields
- 27.** In a photoelectric experiment, the maximum kinetic energy of photoelectrons does **not** depend on the
- A. work function of the emitting material
  - B. wavelength of the incident light
  - C. intensity of the incident light
  - D. energy of an incident photon

28. Copper has a work function of 4.46 eV. What is the maximum kinetic energy of the ejected electrons if the metal is illuminated by light with a wavelength of 450 nm?
- A.  $2.72 \times 10^{-19}$  J
  - B.  $4.42 \times 10^{-19}$  J
  - C.  $7.14 \times 10^{-19}$  J
  - D. 0 J, because no electrons are ejected
29. Louis de Broglie proposed that
- A. the energy absorbed by an atom is the same as the energy released by an atom
  - B. if light has particle properties, then particles have wave properties
  - C. the intensity of light controls the current in the photoelectric effect
  - D. energy and mass are related
30. A burglar knows that an alarm in a certain museum makes use of the photoelectric effect. Ultraviolet light shines on a photocell with a work function of 5.01 eV. Any break in the light will set the alarm off. The burglar realizes that if he shines his own ultraviolet light source at the photocell, he can ensure that there is no break in the light and that the alarm will not be set off. He obtains an ultraviolet light source with a frequency of  $1.13 \times 10^{15}$  Hz. Will he be successful in his burglary attempt and why?
- A. No, because the frequency of the burglar's light is too low for the photocell to function.
  - B. No, because the frequency of the burglar's light is too high for the photocell to function.
  - C. Yes, because the frequency of the burglar's light is low enough for the photocell to function.
  - D. Yes, because the frequency of the burglar's light is high enough for the photocell to function.

Use the following information to answer the next two questions.

Robert Millikan showed experimentally that Einstein's photoelectric equation  $E_{k_{\max}} = hf - W$  was valid. Using a variety of cathode materials, he measured the maximum kinetic energy of photoelectrons while varying the light frequency. The graph shown is typical for a particular cathode. The dotted line is an extrapolation (extension) of the experimental data.



The letters  $w$ ,  $x$ ,  $y$ , and  $z$  represent experimental or extrapolated data.

31. The value for Planck's constant could be determined with the expression

- A.  $\frac{w}{z - y}$
- B.  $\frac{w}{z}$
- C.  $\frac{w(z - y)}{2}$
- D.  $-\frac{y}{x}$



32. The work function of the cathode material is equal to the expression

A.  $\frac{w-x}{z}$

B.  $\frac{w}{z}$

C.  $-x$

D.  $y$

---

33. An atom has energy states  $E_1 = -4.8$  eV,  $E_2 = -2.4$  eV,  $E_3 = -1.2$  eV,  $E_4 = -0.80$  eV, and  $E_5 = -0.40$  eV. The wavelength of emitted light when an electron in the atom makes the transition  $E_4$  to  $E_1$  is

A.  $2.6 \times 10^{-7}$  m

B.  $3.1 \times 10^{-7}$  m

C.  $1.6 \times 10^{-6}$  m





D.  $5.0 \times 10^{-6}$  m

*Use the following information to answer the next question.*


In December 1994, research physicists in Darmstadt, Germany, announced that they had detected three atoms of a new element. With 111 protons and 161 neutrons, this lab-made element had the highest atomic number known to that date. To create element 111, the physicists bombarded bismuth atoms, which have 83 protons, with a beam of nickel atoms, which contain 28 protons. Signals of the three atoms of element 111 appeared for less than two-thousandths of a second. The atoms then decayed into lighter elements and alpha particles. One of the isotopes produced in the decay was element 107 with a mass number of 264. This isotope had never previously been observed.

**Note:** Because neither element 111 nor element 107 had been officially named, element 111 was referred to as X and element 107 was referred to as Y.

34. The overall nuclear equation for this decay reaction is



Use the following information to answer the next three questions.



### Half-Life of Selected Isotopes

Element	Isotope	Half-life	Radiation produced
hydrogen	${}^3_1\text{H}$	12.3 a	$\beta$
carbon	${}^{14}_6\text{C}$	5715 a	$\beta$
iodine	${}^{131}_{53}\text{I}$	8.04 d	$\beta$
lead	${}^{212}_{82}\text{Pb}$	10.6 h	$\beta$
polonium	${}^{194}_{84}\text{Po}$	0.7 s	$\alpha$
polonium	${}^{210}_{84}\text{Po}$	138 d	$\alpha$
uranium	${}^{227}_{92}\text{U}$	1.1 min	$\alpha$
uranium	${}^{235}_{92}\text{U}$	$7.04 \times 10^8$ a	$\alpha$
uranium	${}^{238}_{92}\text{U}$	$4.46 \times 10^9$ a	$\alpha$
plutonium	${}^{236}_{94}\text{Pu}$	2.87 a	$\alpha$
plutonium	${}^{242}_{94}\text{Pu}$	$3.76 \times 10^5$ a	$\alpha$

**Legend:** a = annum = year

35. Given the specifications of this smoke alarm, which of the following isotopes could be used as a radioactive source?

- A.  ${}^3_1\text{H}$
- B.  ${}^{14}_6\text{C}$
- C.  ${}^{194}_{84}\text{Po}$
- D.  ${}^{236}_{94}\text{Pu}$

36. The product of the alpha decay of  ${}^{238}_{92}\text{U}$  is

- A.  ${}^{234}_{90}\text{Th}$
- B.  ${}^{232}_{90}\text{Th}$
- C.  ${}^{232}_{92}\text{U}$
- D.  ${}^{234}_{90}\text{U}$

### Numerical Response

12. Tritium ( ${}^3_1\text{H}$ ), an isotope of hydrogen, was once used in some watches to produce a fluorescent glow. Assuming that the brightness of the glow is proportional to the amount of tritium present, the length of time it would take for the watch to reach  $\frac{1}{4}$  of its original brightness is \_\_\_\_\_ years. (Round and record your answer to three digits.)

\_\_\_\_\_

37. To calculate the amount of energy given off during a fusion reaction, the equation that should be used is

- A.  $E = hf$
- B.  $E = \frac{1}{2}mv^2$
- C.  $E = mc^2$
- D.  $E = \frac{h}{t}$

**Written Response — 11 marks**

- 1.** An astronaut has just landed on an unknown, uninhabited planet and has to send some information about the planet back to Earth. Assume the astronaut has all of the equipment needed to perform the necessary experiments.

Using physics concepts as well as any related formulas, describe procedures that could be used in order to:


- **measure** the magnitude and direction of the gravitational field at the astronaut's location on the unknown planet
- determine whether or not there is an electric field at the location and, if there is, to determine its **magnitude** and **direction**
- determine whether or not there is a magnetic field at the location and, if there is, to determine its **direction**

**Note:** A maximum of 8 marks will be awarded for the physics used to solve this problem. A maximum of 3 marks will be awarded for the effective communication of your response.

Use the following information to answer written-response question 2.

Computer-Generated Data						
time (s)	2.00 kg			1.00 kg		
	$p_x$ kg·m/s	$p_y$ kg·m/s	$ \vec{p} $ kg·m/s	$p_x$ kg·m/s	$p_y$ kg·m/s	$ \vec{p} $ kg·m/s
0.000	18.0	0.00	18.0	0.00	4.00	4.00
0.020	18.0	0.00	18.0	0.00	4.00	4.00
0.040	18.0	0.00	18.0	0.00	4.00	4.00
0.060	18.0	0.00	18.0	0.00	4.00	4.00
0.080	18.0	0.00	18.0	0.00	4.00	4.00
0.100	18.0	0.00	18.0	0.00	4.00	4.00
0.120	18.0	0.00	18.0	0.00	4.00	4.00
0.140	18.0	0.00	18.0	0.00	4.00	4.00
0.160	18.0	0.00	18.0	0.00	4.00	4.00
0.180	12.5	7.31	14.5	5.48	-3.31	6.41
0.200	12.5	7.32	14.5	5.47	-3.32	6.40
0.220	12.5	7.32	14.5	5.47	-3.32	6.40
0.240	12.5	7.32	14.5	5.47	-3.32	6.40
0.260	12.5	7.32	14.5	5.47	-3.32	6.40
0.280	12.5	7.32	14.5	5.47	-3.32	6.40
0.300	12.5	7.32	14.5	5.47	-3.32	6.40
0.320	12.5	7.32	14.5	5.47	-3.32	6.40
0.340	12.5	7.32	14.5	5.47	-3.32	6.40
0.360	12.5	7.32	14.5	5.47	-3.32	6.40
0.380	12.5	7.32	14.5	5.47	-3.32	6.40
0.400	12.5	7.32	14.5	5.47	-3.32	6.40

**Note:**  $p_x$  and  $p_y$  are the  $x$  and  $y$  components of a momentum vector  $\vec{p}$ .




**Written Response — 10 marks**

2. A 2.00 kg ball and a 1.00 kg ball collide. Their original directions of motion are as indicated in the diagram. A computer program that simulates this collision generated the data on the previous page.
- Indicate **on the diagram** the approximate direction of motion for each ball after collision.
  - Determine the speed of each ball before **and** after the collision.
  - Determine the angle between the balls **after** the collision.

*(parts d. and e. are on the next page)*

d. Show that the total momentum before the collision is equal to the total momentum after the collision.

e. How much kinetic energy is lost as a result of this collision?



*You have now completed the examination.  
If you have time, you may wish to check your answers.*

## PHYSICS DATA SHEETS

### CONSTANTS

#### Gravity, Electricity, and Magnetism

Acceleration Due to Gravity or Gravitational Field Near Earth .....	$a_g$ or $g = 9.81 \text{ m/s}^2$ or $9.81 \text{ N/kg}$
Gravitational Constant .....	$G = 6.67 \times 10^{-11} \text{ N}\cdot\text{m}^2/\text{kg}^2$
Mass of Earth .....	$M_e = 5.98 \times 10^{24} \text{ kg}$
Radius of Earth .....	$R_e = 6.37 \times 10^6 \text{ m}$
Coulomb's Law Constant .....	$k = 8.99 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2$
Electron Volt .....	$1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$
Elementary Charge .....	$e = 1.60 \times 10^{-19} \text{ C}$
Index of Refraction of Air .....	$n = 1.00$
Speed of Light in Vacuum .....	$c = 3.00 \times 10^8 \text{ m/s}$

#### Atomic Physics

Energy of an Electron in the 1st Bohr Orbit of Hydrogen .....	$E_1 = -2.18 \times 10^{-18} \text{ J}$ or $-13.6 \text{ eV}$
Planck's Constant .....	$h = 6.63 \times 10^{-34} \text{ J}\cdot\text{s}$
Radius of 1st Bohr Orbit of Hydrogen .....	$r_1 = 5.29 \times 10^{-11} \text{ m}$
Rydberg's Constant for Hydrogen .....	$R_H = 1.10 \times 10^7/\text{m}$

#### Particles

	Rest Mass	Charge
Alpha Particle .....	$m_\alpha = 6.65 \times 10^{-27} \text{ kg}$	$\alpha^{2+}$
Electron .....	$m_e = 9.11 \times 10^{-31} \text{ kg}$	$e^-$
Neutron .....	$m_n = 1.67 \times 10^{-27} \text{ kg}$	$n^0$
Proton .....	$m_p = 1.67 \times 10^{-27} \text{ kg}$	$p^+$

#### Trigonometry and Vectors

For any Vector  $\vec{R}$

$$R = \sqrt{R_x^2 + R_y^2}$$

$$\tan \theta = \frac{R_y}{R_x}$$

$$R_x = R \cos \theta$$

$$R_y = R \sin \theta$$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

#### Prefixes Used With SI Units

Prefix	Symbol	Exponential Value	Prefix	Symbol	Exponential Value
pico .....	p	$10^{-12}$	tera .....	T	$10^{12}$
nano .....	n	$10^{-9}$	giga .....	G	$10^9$
micro .....	$\mu$	$10^{-6}$	mega .....	M	$10^6$
milli .....	m	$10^{-3}$	kilo .....	k	$10^3$
centi .....	c	$10^{-2}$	hecto .....	h	$10^2$
deci .....	d	$10^{-1}$	deka .....	da	$10^1$



## EQUATIONS

### Kinematics

$$\bar{v}_{\text{ave}} = \frac{\bar{d}}{t}$$

$$\bar{a} = \frac{\bar{v}_f - \bar{v}_i}{t}$$

$$\bar{d} = \bar{v}_f t - \frac{1}{2} \bar{a} t^2$$

$$\bar{d} = \left( \frac{\bar{v}_f + \bar{v}_i}{2} \right) t$$

$$\bar{d} = \bar{v}_i t + \frac{1}{2} \bar{a} t^2$$

$$v_f^2 = v_i^2 + 2ad$$

### Dynamics

$$\bar{F} = m\bar{a}$$

$$F_g = \frac{Gm_1 m_2}{r^2}$$

$$\bar{F} \Delta t = m \Delta \bar{v}$$

$$g = \frac{Gm_1}{r^2}$$

$$\bar{F}_g = m\bar{g}$$

$$F_c = \frac{mv^2}{r}$$

$$F_f = \mu F_N$$

$$F_c = \frac{4\pi^2 mr}{T^2}$$

$$\bar{F}_s = -k\bar{x}$$

### Momentum and Energy

$$\bar{p} = m\bar{v}$$

$$E_k = \frac{1}{2} mv^2$$

$$W = Fd$$

$$E_p = mgh$$

$$W = \Delta E = Fd \cos \theta$$

$$P = \frac{W}{t} = \frac{\Delta E}{t}$$

$$E_p = \frac{1}{2} kx^2$$

### Waves and Light

$$T = 2\pi \sqrt{\frac{m}{k}}$$

$$T = 2\pi \sqrt{\frac{l}{g}}$$

$$T = \frac{1}{f}$$

$$\lambda = \frac{d \sin \theta}{n}$$

$$v = f\lambda$$

$$m = \frac{h_i}{h_o} = \frac{-d_i}{d_o}$$

$$\frac{\lambda_1}{2} = l; \frac{\lambda_1}{4} = l$$

$$\frac{1}{f} = \frac{1}{d_o} + \frac{1}{d_i}$$

### Atomic Physics

$$hf = E_k + W_{\text{max}}$$

$$\frac{1}{\lambda} = R_H \left( \frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

$$W = hf_0$$

$$E_n = \frac{1}{n^2} E_1$$

$$E_{k_{\text{max}}} = qV_{\text{stop}}$$

$$r_n = n^2 r_1$$

$$E = hf = \frac{hc}{\lambda}$$

$$N = N_0 \left( \frac{1}{2} \right)^n$$

### Quantum Mechanics and Nuclear Physics

$$E = mc^2$$

$$p = \frac{h}{\lambda}$$

$$p = \frac{hf}{c}; E = pc$$

### Electricity and Magnetism

$$F_e = \frac{kq_1 q_2}{r^2}$$

$$V = IR$$

$$|\bar{E}| = \frac{kq_1}{r^2}$$

$$P = IV$$

$$\bar{E} = \frac{\bar{F}_e}{q}$$

$$I = \frac{q}{t}$$

$$|\bar{E}| = \frac{V}{d}$$

$$F_m = I\bar{B}_{\perp}$$

$$V = \frac{\Delta E}{q}$$

$$F_m = qv\bar{B}_{\perp}$$

$$R = R_1 + R_2 + R_3$$

$$V = l v \bar{B}_{\perp}$$

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$$

$$\frac{N_p}{N_s} = \frac{V_p}{V_s} = \frac{I_s}{I_p}$$

$$I_{\text{eff}} = 0.707 I_{\text{max}}$$

$$V_{\text{eff}} = 0.707 V_{\text{max}}$$

# Periodic Table of the Elements

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
IA	IIA	IIIB	IVB	VB	VIB	VII B	VIII B	VIIIB	VIIIB	IB	IIB	IIIA	IVA	VA	VIA	VIIA	VIIIA or O
<b>1</b> H 1.01 hydrogen	<b>2</b> He 4.00 helium																
<b>3</b> Li 6.94 lithium	<b>4</b> Be 9.01 beryllium	<b>5</b> B 10.81 boron	<b>6</b> C 12.01 carbon	<b>7</b> N 14.01 nitrogen	<b>8</b> O 16.00 oxygen	<b>9</b> F 19.00 fluorine	<b>10</b> Ne 20.17 neon	<b>11</b> Na 22.99 sodium	<b>12</b> Mg 24.31 magnesium	<b>13</b> Al 26.98 aluminum	<b>14</b> Si 28.09 silicon	<b>15</b> P 30.97 phosphorus	<b>16</b> S 32.06 sulphur	<b>17</b> Cl 35.45 chlorine	<b>18</b> Ar 39.95 argon		
<b>19</b> K 39.10 potassium	<b>20</b> Ca 40.08 calcium	<b>21</b> Sc 44.96 scandium	<b>22</b> Ti 47.90 titanium	<b>23</b> V 50.94 vanadium	<b>24</b> Cr 52.00 chromium	<b>25</b> Mn 54.94 manganese	<b>26</b> Fe 55.85 iron	<b>27</b> Co 58.93 cobalt	<b>28</b> Ni 58.71 nickel	<b>29</b> Cu 63.55 copper	<b>30</b> Zn 65.38 zinc	<b>31</b> Ga 69.72 gallium	<b>32</b> Ge 72.59 germanium	<b>33</b> As 74.92 arsenic	<b>34</b> Se 78.96 selenium	<b>35</b> Br 79.90 bromine	<b>36</b> Kr 83.80 krypton
<b>37</b> Rb 85.47 rubidium	<b>38</b> Sr 87.62 strontium	<b>39</b> Y 88.91 yttrium	<b>40</b> Zr 91.22 zirconium	<b>41</b> Nb 92.91 niobium	<b>42</b> Mo 95.94 molybdenum	<b>43</b> Tc (98.91) technetium	<b>44</b> Ru 101.07 ruthenium	<b>45</b> Rh 102.91 rhodium	<b>46</b> Pd 106.40 palladium	<b>47</b> Ag 107.87 silver	<b>48</b> Cd 112.41 cadmium	<b>49</b> In 114.82 indium	<b>50</b> Sn 118.69 tin	<b>51</b> Sb 121.75 antimony	<b>52</b> Te 127.60 tellurium	<b>53</b> I 126.90 iodine	<b>54</b> Xe 131.30 xenon
<b>55</b> Cs 132.91 cesium	<b>56</b> Ba 137.33 barium	<b>57-71</b> Fr (223.02) francium	<b>72</b> Hf 178.49 hafnium	<b>73</b> Ta 180.95 tantalum	<b>74</b> W 183.85 tungsten	<b>75</b> Re 186.21 rhenium	<b>76</b> Os 190.20 osmium	<b>77</b> Ir 192.22 iridium	<b>78</b> Pt 195.09 platinum	<b>79</b> Au 196.97 gold	<b>80</b> Hg 200.59 mercury	<b>81</b> Tl 204.37 thallium	<b>82</b> Pb 207.19 lead	<b>83</b> Bi 208.98 bismuth	<b>84</b> Po (208.98) polonium	<b>85</b> At (209.98) astatine	<b>86</b> Rn (222.02) radon
<b>87</b> Fr (223.02) francium	<b>88</b> Ra (226.03) radium	<b>89-103</b> Unp (266.11) unnilpentium	<b>104</b> Unq (266.11) unnilquadium	<b>105</b> Unp (262.11) unnilpentium	<b>106</b> Unh (263.12) unnilhexium	<b>107</b> Uns (262.12) unnilseptium	<b>108</b> Uno (265) unniloctium	<b>109</b> Une (266) unnilennium	<b>63</b> Eu 151.96 europium	<b>64</b> Gd 157.25 gadolinium	<b>65</b> Tb 158.93 terbium	<b>66</b> Dy 162.50 dysprosium	<b>67</b> Ho 164.93 holmium	<b>68</b> Er 167.26 erbium	<b>69</b> Tm 168.93 thulium	<b>70</b> Yb 173.04 ytterbium	<b>71</b> Lu 174.97 lutetium
<b>89</b> Ac (227.03) actinium	<b>90</b> Th (232.04) thorium	<b>91</b> Pa (231.04) protactinium	<b>92</b> U 238.03 uranium	<b>93</b> Np (237.05) neptunium	<b>94</b> Pu (244.06) plutonium	<b>95</b> Am (243.06) americium	<b>96</b> Cm (247.07) curium	<b>97</b> Bk (247.07) berkelium	<b>98</b> Cf (242.06) californium	<b>99</b> Es (252.08) einsteinium	<b>100</b> Fm (257.10) fermium	<b>101</b> Md (258.10) mendelevium	<b>102</b> No (259.10) nobelium	<b>103</b> Lr (260.11) lawrencium			

**Key**

Atomic number → **3** Li → Symbol

Atomic molar mass → **6.94**


Name → lithium

Based on  $^{12}\text{C}$

( ) Indicates mass of the most stable isotope

*No marks will be given for work done on this page.*

*Fold and tear along perforation.*



*No marks will be given for work done on this page.*

*Fold and tear along perforation.*

mm

10

20

30

40

50

60

70

80

90

100

110

120

130

140

150

160

170

180


190

200

210

*No marks will be given for work done on this page.*

*Fold and tear along perforation.*



*No marks will be given for work done on this page.*





**U.S. DEPARTMENT OF EDUCATION**  
*Office of Educational Research and Improvement (OERI)*  
*Educational Resources Information Center (ERIC)*



## NOTICE

### REPRODUCTION BASIS

This document is covered by a signed "Reproduction Release (Blanket)" form (on file within the ERIC system), encompassing all or classes of documents from its source organization and, therefore, does not require a "Specific Document" Release form.

This document is Federally-funded, or carries its own permission to reproduce, or is otherwise in the public domain and, therefore, may be reproduced by ERIC without a signed Reproduction Release form (either "Specific Document" or "Blanket").