

DOCUMENT RESUME

ED 474 680

CG 032 291

AUTHOR Goggin, K.; Metcalf, K.; Wise, D.; Kennedy, S.; Murray, T.; Burgess, D.; Reese-Smith, J.; Terhune, N.; Broadus, K.; Downes, A.; Buckendahl, H.

TITLE Youth-Initiated HIV Risk and Substance Use Prevention Program.

PUB DATE 2002-08-00

NOTE 20p.; Paper presented at the Annual Conference of the American Psychological Association (110th, Chicago, IL, August 22-25, 2002).

PUB TYPE Reports - Evaluative (142) -- Speeches/Meeting Papers (150)

EDRS PRICE EDRS Price MF01/PC01 Plus Postage.

DESCRIPTORS Acquired Immunodeficiency Syndrome; Adolescent Attitudes; Condoms; Drinking; Drug Use; Feasibility Studies; Inner City; Peer Influence; *Prevention; *Program Effectiveness; *Program Evaluation; *Substance Abuse; *Youth Programs

IDENTIFIERS *Risk Taking Behavior

ABSTRACT

This study evaluates the first year of a novel HIV and substance use prevention program for inner city youth (Offering New Youth eXperiences--ONYX). Baseline and follow-up measures of knowledge, attitudes, and risk behaviors were administered seven months apart to 441 youth participating in the ONYX program. Youth (n=71) who provided data at both baseline and follow-up assessments are reported on here. Significant reduction in risk behavior was not observed, however improvement in other outcomes was noted. Youth's knowledge about HIV significantly increased from baseline to follow-up. Further, increases in condom use self-efficacy, perceived harm from drug use, and resistance of peer pressure to drink alcohol were observed. Overall, the results of this feasibility phase suggest that the design is possible and the program may be effective in producing some of the desired outcomes. (Author)

Reproductions supplied by EDRS are the best that can be made
from the original document.

A Youth-Initiated HIV Risk and Substance Use Prevention Program

by

K. Goggin
 K. Metcalf
 D. Wise
 S. Kennedy
 T. Murray
 D. Burgess
 J. Reese-Smith
 N. Terhune
 K. Broadus
 A. Downes
 H. Buckendahl

U.S. DEPARTMENT OF EDUCATION
 Office of Educational Research and Improvement
 EDUCATIONAL RESOURCES INFORMATION
 CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE AND
 DISSEMINATE THIS MATERIAL HAS
 BEEN GRANTED BY

K. GOGGIN

TO THE EDUCATIONAL RESOURCES
 INFORMATION CENTER (ERIC)

1

Abstract: A Youth-initiated HIV Risk and Substance Use Prevention Program

This study evaluates the first year of a novel HIV and substance use prevention program for inner city youth (Offering New Youth eXperiences – ONYX). Baseline and follow-up measures of knowledge, attitudes, and risk behaviors were administered seven months apart to 441 youth participating in the ONYX program. Youth (n=71) who provided data at both baseline and follow-up assessments are reported on here. Significant reductions in risk behavior were not observed, however improvement in other outcomes were noted. Youth's knowledge about HIV significantly increased from baseline to follow-up [$t(68)=3.23, p<0.05$]. Further, increases in condom use self-efficacy ($X^2=5.08, p<0.05$), perceived harm from drug use ($X^2=4.06, p<0.05$), and resistance of peer pressure to drink alcohol ($X^2=5.00, p<0.05$) were observed. Overall, the results of this feasibility phase suggest that the design is possible and the program may be effective in producing some of the desired outcomes.

Goggin, K., Metcalf, K., Wise, D., Kennedy, S., Murray, T., Burgess, D., Reese-Smith, J., Terhune, N., Broadus, K., Downes, A., & Buckendahl, H. (2001). *A Youth-initiated HIV Risk and Substance Use Prevention Program*. Under review for the 109th Annual Meeting of the American Psychological Association.

A Youth-initiated HIV Risk and Substance Use Prevention Program

In 1999, African American (AA) adolescents (age 13-19) accounted for 56% of all new HIV diagnoses and rates of HIV infection continue to rise among AA youth despite declining rates in other populations (CDC, 2001). Evidence is emerging to suggest that youth who engage in risky sexual behaviors also use substances. For example, AA males (age 15-18) who engage in risky sexual behaviors are also likely to report alcohol and other drug use behaviors (Newman & Zimmerman, 2000). Similarly, adolescent female marijuana users report a greater number of lifetime sexual partners and are more likely to have had an STD (Boyer, 2000).

While prevention programs specifically targeting AA youth are emerging (e.g., Jemmott, Jemmott & Fong, 1998), few are community-based programs that involve youth at every level of the program and even fewer simultaneously target both substance use and HIV risk behaviors. This paper reports the results of an evaluation of the feasibility phase of the Offering New Youth eXperiences (ONYX) program which is a community-based youth-initiated combined substance use and HIV risk behavior reduction program implemented by the Kansas City Free Health Clinic.

The ONYX Program

Working from a Diffusions of Innovations (Rogers, 2002) framework, the ONYX

program involves youth at all levels of the program, from the development of prevention materials and activities, to the distribution of grant funds. The process is guided by a group of 10-12 student ambassadors who are recruited from the community sites. These youth receive training in substance use/abuse and HIV risk reduction, epidemiology, leadership and speaking skills, and grantsmanship. Ambassadors then develop a request for combined substance use and HIV risk reduction grant applications from their peers and develop a plan for informing their peers about the program and the availability of up to \$2000 for youth-initiated prevention programs. The ambassadors then host a conference to educate their peers about substance use and HIV risk behaviors and teach them how to write grant proposals. Two months later the ambassadors review the grants and decide which to fund. During the year that is reported on here, \$25,815.22 were distributed to support youth-initiated prevention activities.


Participants were 441 youth enrolled in one of the alternative schools or church based youth groups that participated in ONYX. Baseline and follow-up measures of HIV/AIDS knowledge, attitudes, and risk behaviors were administered seven months apart. Due to the tremendous dropout rate at the sites, only 71 youth that completed baseline and follow-up questionnaires during the feasibility phase are reported on here. Further, the control group was dropped as it was inappropriate in terms of risk behaviors and age. At baseline, participants averaged 15 years of age ($SD = 2.4$, range 11-20), most identified as African American (86%) and approximately half were females (53%).

Sexual Behaviors, Self-Efficacy, and HIV Knowledge

Although a reduction in risky sexual behaviors (i.e., sex without a condom) was observed from baseline ($M = 4.2$, $SD = 8.6$) to follow-up ($M = 2.4$, $SD = 4.4$), it did not reach statistical significance [$t(37) = 1.03$, $p > 0.05$].


However, participants did demonstrate increased confidence in their ability to get their partner to use a condom at follow-up. Four percent of the participants strongly disagreed with the statement “*I can get my partner to use a condom, even if he or she does not want to*” at follow-up compared to 15% at baseline ($\chi^2 = 5.08$, $p < 0.05$).

I can get my partner to use a condom even if he or she does not want to



Youth's knowledge about HIV also significantly increased from baseline to follow-up [t(68) = 3.23, p < 0.05].

HIV is present in blood, semen, vaginal fluid & breast milk (percentage)




Alcohol Use and Self-Efficacy

Alcohol use did not decrease at follow-up. Specifically, 23% reported alcohol consumption an average of 11 days during the month prior to the follow-up assessment (SD = 12.2, range 1-30). Furthermore, the percentage of participants who consumed 5 or more alcoholic beverages increased from baseline (7%) to follow-up (13%) with youth reporting binge drinking on an average of 11 days during the past month (SD = 7.9, range 1-21).

However, respondents' attitudes and perceptions of alcohol use did change favorably. For example, 86% of the participants at follow-up believed that drinking alcohol would probably or definitely *not* make them more popular with their peers, compared to 70% at baseline ($\chi^2 = 5.00, p < 0.05$).

I believe if I drink alcohol I will be more accepted by my peers



Participants' perception of the physical risks associated with alcohol use increased at follow-up with more than half the sample (58%) reporting that there was a moderate to great physical risk associated with having 4 or more alcoholic beverages once or twice a week as compared to 38% at baseline ($\chi^2 = 4.06, p < 0.05$)


Marijuana Use and Self-Efficacy

The proportion of participants who used marijuana one month prior to the follow-up assessment decreased slightly as compared to baseline (28% and 34%, respectively).

However, the average number of days of marijuana use remained constant from baseline to follow-up ($M = 14$ days, $SD = 12.5$, range 1-30).

There was a trend towards improved perception of physical risks associated with marijuana from baseline to follow-up. Specifically, more than half of the sample (60%) believed that there was a moderate to great risk involved with smoking marijuana once a month or more, compared to 43% of participants at baseline ($\chi^2 = 3.67$, $p = .07$).

How much do people risk harming themselves when they smoke marijuana once or twice a week.



Conclusion

Significant changes in self-efficacy and knowledge related to HIV and substance use prevention was observed among the students we were able to assess at both baseline and follow-up, but not behavior change. Additional procedures will be implemented to reduce drop-outs in the future, however this may remain a limitation of the evaluation of the program due to the transient nature of the youth at the community sites. Results also suggest that to accomplish behavior change, that a longer duration of intervention and more technical assistance to youth to strengthen their prevention efforts may be required. Interpretation of these preliminary results is also limited by the lack of an appropriate control group, as factors like maturation may have had a significant impact on the risk behaviors being studied. Overall, the results of this feasibility phase suggest that the design is possible and the program may be effective in reducing most of the desired outcomes.

References

- Boyer, C., Shafer, M., Wibbelsman, C., Seeberg, D., Teitle, E., Lovell, N. (2000). Associations of sociodemographic, psychosocial, and behavioral factors with sexual risk and sexually transmitted diseases in teen clinic patients. *Journal of Adolescent Health, 27*, 102-111.
- Centers for Disease Control and Prevention (2001). Young people at risk: HIV/AIDS among American's youth. National Center for HIV, STD, and TB Prevention. www.cdc.gov/hiv/pubs/facts/youth.htm.
- Jemmott, J.B., Jemmott, L.S., & Fong, G.T. (1998). Abstinence and safer sex HIV risk-reduction interventions for African American adolescents. *Journal of the American Medical Association, 279* (19), 1529-1536.
- Newman, P. & Zimmerman, M. (2000). Gender differences in HIV-related sexual risk behavior among urban African American youth: a multivariate approach. *AIDS Education and Prevention, 12* (4), 308-325.
- Rogers, E.M. (2002). Diffusion of preventive innovations. *Addictive Behaviors, 27* (6), 989-993.

Abstract: A Youth-initiated HIV Risk and Substance Use Prevention Program

This study evaluates the first year of a novel HIV and substance use prevention program for inner city youth (Offering New Youth eXperiences – ONYX). Baseline and follow-up measures of knowledge, attitudes, and risk behaviors were administered seven months apart to 441 youth participating in the ONYX program. Youth (n=71) who provided data at both baseline and follow-up assessments are reported on here. Significant reductions in risk behavior were not observed, however improvement in other outcomes were noted. Youth's knowledge about HIV significantly increased from baseline to follow-up [$t(68)=3.23$, $p<0.05$]. Further, increases in condom use self-efficacy ($X^2=5.08$, $p<0.05$), perceived harm from drug use ($X^2=4.06$, $p<0.05$), and resistance of peer pressure to drink alcohol ($X^2=5.00$, $p<0.05$) were observed. Overall, the results of this feasibility phase suggest that the design is possible and the program may be effective in producing some of the desired outcomes.

Goggin, K., Metcalf, K., Wise, D., Kennedy, S., Murray, T., Burgess, D., Reese-Smith, J., Terhune, N., Broadus, K., Downes, A., & Buckendahl, H. (2001). *A Youth-initiated HIV Risk and Substance Use Prevention Program*. Under review for the 109th Annual Meeting of the American Psychological Association.

A Youth-initiated HIV Risk and Substance Use Prevention Program

In 1999, African American (AA) adolescents (age 13-19) accounted for 56% of all new HIV diagnoses and rates of HIV infection continue to rise among AA youth despite declining rates in other populations (CDC, 2001). Evidence is emerging to suggest that youth who engage in risky sexual behaviors also use substances. For example, AA males (age 15-18) who engage in risky sexual behaviors are also likely to report alcohol and other drug use behaviors (Newman & Zimmerman, 2000). Similarly, adolescent female marijuana users report a greater number of lifetime sexual partners and are more likely to have had an STD (Boyer, 2000).

While prevention programs specifically targeting AA youth are emerging (e.g., Jemmott, Jemmott & Fong, 1998), few are community-based programs that involve youth at every level of the program and even fewer simultaneously target both substance use and HIV risk behaviors. This paper reports the results of an evaluation of the feasibility phase of the Offering New Youth eXperiences (ONYX) program which is a community-based youth-initiated combined substance use and HIV risk behavior reduction program implemented by the Kansas City Free Health Clinic.

The ONYX Program

Working from a Diffusions of Innovations (Rogers, 2002) framework, the ONYX

program involves youth at all levels of the program, from the development of prevention materials and activities, to the distribution of grant funds. The process is guided by a group of 10-12 student ambassadors who are recruited from the community sites. These youth receive training in substance use/abuse and HIV risk reduction, epidemiology, leadership and speaking skills, and grantsmanship. Ambassadors then develop a request for combined substance use and HIV risk reduction grant applications from their peers and develop a plan for informing their peers about the program and the availability of up to \$2000 for youth-initiated prevention programs. The ambassadors then host a conference to educate their peers about substance use and HIV risk behaviors and teach them how to write grant proposals. Two months later the ambassadors review the grants and decide which to fund. During the year that is reported on here, \$25,815.22 were distributed to support youth-initiated prevention activities.


Participants were 441 youth enrolled in one of the alternative schools or church based youth groups that participated in ONYX. Baseline and follow-up measures of HIV/AIDS knowledge, attitudes, and risk behaviors were administered seven months apart. Due to the tremendous dropout rate at the sites, only 71 youth that completed baseline and follow-up questionnaires during the feasibility phase are reported on here. Further, the control group was dropped as it was inappropriate in terms of risk behaviors and age. At baseline, participants averaged 15 years of age ($SD = 2.4$, range 11-20), most identified as African American (86%) and approximately half were females (53%).

Sexual Behaviors, Self-Efficacy, and HIV Knowledge

Although a reduction in risky sexual behaviors (i.e., sex without a condom) was observed from baseline ($M = 4.2$, $SD = 8.6$) to follow-up ($M = 2.4$, $SD = 4.4$), it did not reach statistical significance [$t(37) = 1.03$, $p > 0.05$].


However, participants did demonstrate increased confidence in their ability to get their partner to use a condom at follow-up. Four percent of the participants strongly disagreed with the statement “*I can get my partner to use a condom, even if he or she does not want to*” at follow-up compared to 15% at baseline ($\chi^2 = 5.08$, $p < 0.05$).

I can get my partner to use a condom even if he or she does not want to



Youth's knowledge about HIV also significantly increased from baseline to follow-up [t(68) = 3.23, p < 0.05].

HIV is present in blood, semen, vaginal fluid & breast milk (percentage)




Alcohol Use and Self-Efficacy

Alcohol use did not decrease at follow-up. Specifically, 23% reported alcohol consumption an average of 11 days during the month prior to the follow-up assessment (SD = 12.2, range 1-30). Furthermore, the percentage of participants who consumed 5 or more alcoholic beverages increased from baseline (7%) to follow-up (13%) with youth reporting binge drinking on an average of 11 days during the past month (SD = 7.9, range 1-21).

However, respondents' attitudes and perceptions of alcohol use did change favorably. For example, 86% of the participants at follow-up believed that drinking alcohol would probably or definitely *not* make them more popular with their peers, compared to 70% at baseline ($\chi^2 = 5.00, p < 0.05$).

I believe if I drink alcohol I will be more accepted by my peers



Participants' perception of the physical risks associated with alcohol use increased at follow-up with more than half the sample (58%) reporting that there was a moderate to great physical risk associated with having 4 or more alcoholic beverages once or twice a week as compared to 38% at baseline ($\chi^2 = 4.06, p < 0.05$).


Marijuana Use and Self-Efficacy

The proportion of participants who used marijuana one month prior to the follow-up assessment decreased slightly as compared to baseline (28% and 34%, respectively).

However, the average number of days of marijuana use remained constant from baseline to follow-up ($M = 14$ days, $SD = 12.5$, range 1-30).

There was a trend towards improved perception of physical risks associated with marijuana from baseline to follow-up. Specifically, more than half of the sample (60%) believed that there was a moderate to great risk involved with smoking marijuana once a month or more, compared to 43% of participants at baseline ($\chi^2 = 3.67$, $p = .07$).

How much do people risk harming themselves when they smoke marijuana once or twice a week.




Conclusion

Significant changes in self-efficacy and knowledge related to HIV and substance use prevention was observed among the students we were able to assess at both baseline and follow-up, but not behavior change. Additional procedures will be implemented to reduce drop-outs in the future, however this may remain a limitation of the evaluation of the program due to the transient nature of the youth at the community sites. Results also suggest that to accomplish behavior change, that a longer duration of intervention and more technical assistance to youth to strengthen their prevention efforts may be required. Interpretation of these preliminary results is also limited by the lack of an appropriate control group, as factors like maturation may have had a significant impact on the risk behaviors being studied. Overall, the results of this feasibility phase suggest that the design is possible and the program may be effective in reducing most of the desired outcomes.

References

- Boyer, C., Shafer, M., Wibbelsman, C., Seeberg, D., Teitle, E., Lovell, N. (2000). Associations of sociodemographic, psychosocial, and behavioral factors with sexual risk and sexually transmitted diseases in teen clinic patients. *Journal of Adolescent Health, 27*, 102-111.
- Centers for Disease Control and Prevention (2001). Young people at risk: HIV/AIDS among American's youth. National Center for HIV, STD, and TB Prevention. www.cdc.gov/hiv/pubs/facts/youth.htm.
- Jemmott, J.B., Jemmott, L.S., & Fong, G.T. (1998). Abstinence and safer sex HIV risk-reduction interventions for African American adolescents. *Journal of the American Medical Association, 279* (19), 1529-1536.
- Newman, P. & Zimmerman, M. (2000). Gender differences in HIV-related sexual risk behavior among urban African American youth: a multivariate approach. *AIDS Education and Prevention, 12* (4), 308-325.
- Rogers, E.M. (2002). Diffusion of preventive innovations. *Addictive Behaviors, 27* (6), 989-993.



U.S. Department of Education
Office of Educational Research and Improvement (OERI)
National Library of Education (NLE)
Educational Resources Information Center (ERIC)



REPRODUCTION RELEASE

(Specific Document)

I. DOCUMENT IDENTIFICATION:

Title: A Youth-initiated HIV Risk and Substance Use Prevention Program	
Author(s): Goggin, K Metcalf, K Wise, D Kennedy, S Murray, T, Bugess, D, Reese-Smith, J Tepune, N	
Corporate Source: University of Missouri Kansas City	Publication Date: 2001

II. REPRODUCTION RELEASE:

In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, *Resources in Education* (RIE), are usually made available to users in microfiche, reproduced paper copy, and electronic media, and sold through the ERIC Document Reproduction Service (EDRS). Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to each document.

If permission is granted to reproduce and disseminate the identified documents, please CHECK ONE of the following three options and sign at the bottom of the page.

<p>The sample sticker shown below will be affixed to all Level 1 documents</p> <div style="border: 1px solid black; padding: 5px;"> <p>PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY -</p> <p><u>Dr. Kathy Goggin</u></p> <hr/> <p>TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)</p> </div> <p>1</p> <div style="text-align: center;"> <p>Level 1</p> <input checked="" type="checkbox"/> </div>	<p>The sample sticker shown below will be affixed to all Level 2A documents</p> <div style="border: 1px solid black; padding: 5px;"> <p>PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN MICROFICHE, AND IN ELECTRONIC MEDIA FOR ERIC COLLECTION SUBSCRIBERS ONLY, HAS BEEN GRANTED BY</p> <p>_____</p> <hr/> <p>TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)</p> </div> <p>2A</p> <div style="text-align: center;"> <p>Level 2A</p> <input type="checkbox"/> </div>	<p>The sample sticker shown below will be affixed to all Level 2B documents</p> <div style="border: 1px solid black; padding: 5px;"> <p>PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN MICROFICHE ONLY HAS BEEN GRANTED BY</p> <p>_____</p> <hr/> <p>TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)</p> </div> <p>2B</p> <div style="text-align: center;"> <p>Level 2B</p> <input type="checkbox"/> </div>
---	--	--

Check here for Level 1 release, permitting reproduction and dissemination in microfiche or other ERIC archival media (e.g., electronic) and paper copy.

Check here for Level 2A release, permitting reproduction and dissemination in microfiche and in electronic media for ERIC archival collection subscribers only

Check here for Level 2B release, permitting reproduction and dissemination in microfiche only

Documents will be processed as indicated provided reproduction quality permits. If permission to reproduce is granted, but no box is checked, documents will be processed at Level 1.

I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate these documents as indicated above. Reproduction from the ERIC microfiche or electronic media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries.

Sign here, please →

Signature: <u>Kathy Goggin</u>	Printed Name/Position/Title: Dr. Kathy Goggin Associate Professor	
Organization/Address: 4825 Troost, suite 111-D	Telephone: 816-235-1059	FAX: 816 235 - 1062
	E-Mail Address: goggink@umkc.edu	Date: 03-10-03



III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of these documents from another source, please provide the following information regarding the availability of these documents. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:
Address:
Price:

IV. REFERRAL OF ERIC TO COPYRIGHT/REPRODUCTION RIGHTS HOLDER:

If the right to grant this reproduction release is held by someone other than the addressee, please provide the appropriate name and address:

Name:
Address:

V. WHERE TO SEND THIS FORM:

Send this form to the following ERIC Clearinghouse:	ERIC Counseling & Student Services University of North Carolina at Greensboro 201 Ferguson Building PO Box 26171 Greensboro, NC 27402-6171
---	---